


Library
National Wetlands Research Center
U. S. Fish and Wildlife Service
700 Cajandome Boulevard
Lafayette, La. 70506

**Biological Services Program
and
Division of Ecological Services**

FWS/OBS-82/10.16
JULY 1982

**HABITAT SUITABILITY INDEX MODELS:
LARGEMOUTH BASS**



Fish and Wildlife Service

S. Department of the Interior

SK
361
.U54
no. 82-
10.16

The Biological Services Program was established within the U.S. Fish and Wildlife Service to supply scientific information and methodologies on key environmental issues that impact fish and wildlife resources and their supporting ecosystems. The mission of the program is as follows:

- To strengthen the Fish and Wildlife Service in its role as a primary source of information on national fish and wildlife resources, particularly in respect to environmental impact assessment.
- To gather, analyze, and present information that will aid decisionmakers in the identification and resolution of problems associated with major changes in land and water use.
- To provide better ecological information and evaluation for Department of the Interior development programs, such as those relating to energy development.

Information developed by the Biological Services Program is intended for use in the planning and decisionmaking process to prevent or minimize the impact of development on fish and wildlife. Research activities and technical assistance services are based on an analysis of the issues, a determination of the decisionmakers involved and their information needs, and an evaluation of the state of the art to identify information gaps and to determine priorities. This is a strategy that will ensure that the products produced and disseminated are timely and useful.

Projects have been initiated in the following areas: coal extraction and conversion; power plants; geothermal, mineral and oil shale development; water resource analysis, including stream alterations and western water allocation; coastal ecosystems and Outer Continental Shelf development; and systems inventory, including National Wetland Inventory, habitat classification and analysis, and information transfer.

The Biological Services Program consists of the Office of Biological Services in Washington, D.C., which is responsible for overall planning and management; National Teams, which provide the Program's central scientific and technical expertise and arrange for contracting biological services studies with states, universities, consulting firms, and others; Regional Staffs, who provide a link to problems at the operating level; and staffs at certain Fish and Wildlife Service research facilities, who conduct in-house research studies.

HABITAT SUITABILITY INDEX MODELS: LARGEMOUTH BASS

by

Robert J. Stuber
Habitat Evaluation Procedures Group
Western Energy and Land Use Team
U.S. Fish and Wildlife Service
Drake Creekside Building One
2625 Redwing Road
Fort Collins, CO 80526

Glen Gebhart
and
O. Eugene Maughan
Oklahoma Cooperative Fishery Research Unit
Oklahoma State University
Stillwater, OK 74074

Western Energy and Land Use Team
Office of Biological Services
Fish and Wildlife Service
U.S. Department of the Interior
Washington, DC 20240

This report should be cited as:

Stuber, R. J., G. Gebhart, and O. E. Maughan. 1982. Habitat suitability index models: Largemouth bass. U.S. Dept. Int. Fish Wildl. Serv. FWS/OBS-82/10.16. 32 pp.

PREFACE

The habitat use information and Habitat Suitability Index (HSI) models presented in this document are an aid for impact assessment and habitat management activities. Literature concerning a species' habitat requirements and preferences is reviewed and then synthesized into HSI models, which are scaled to produce an index between 0 (unsuitable habitat) and 1 (optimal habitat). Assumptions used to transform habitat use information into these mathematical models are noted, and guidelines for model application are described. Any models found in the literature which can also be used to calculate an HSI are cited, and simplified HSI models, based on what the authors believe to be the most important characteristics for this species, are presented.

Use of the models presented in this publication for impact assessment requires the setting of clear study objectives and may require modification of the models to meet those objectives. Methods for reducing model complexity and recommended measurement techniques for model variables are presented in Terrell et al. (in press).¹ A discussion of HSI model building techniques, including the component approach, is presented in U.S. Fish and Wildlife Service (1981).²

The HSI models presented herein are complex hypotheses of species-habitat relationships, not statements of proven cause and effect relationships. Results of model performance tests, when available, are referenced; however, models that have demonstrated reliability in specific situations may prove unreliable in others. For this reason, the U.S. Fish and Wildlife Service encourages model users to send comments and suggestions that might help us increase the utility and effectiveness of this habitat-based approach to fish and wildlife planning. Please send comments to:

Habitat Evaluation Procedures Group
Western Energy and Land Use Team
U.S. Fish and Wildlife Service
2625 Redwing Road
Ft. Collins, CO 80526

¹Terrell, J. W., T. E. McMahon, P. D. Inskip, R. F. Raleigh, and K. W. Williamson (in press). Habitat Suitability Index Models: Appendix A. Guidelines for riverine and lacustrine applications of fish HSI models with the habitat evaluation procedures. U.S. Dept. Int. Fish Wildl. Serv. FWS/OBS 82-10.A.

²U.S. Fish and Wildlife Service. 1981. Standards for the development of Habitat Suitability Index models. 103 ESM. U.S. Dept. Int. Fish Wildl. Serv., Div. Ecol. Serv.

CONTENTS

	<u>Page</u>
PREFACE	iii
ACKNOWLEDGMENTS	v
HABITAT USE INFORMATION	1
General	1
Age, Growth, and Food	1
Reproduction	1
Specific Habitat Requirements	2
HABITAT SUITABILITY INDEX (HSI) MODELS	4
Model Applicability	4
Model Description - Riverine	5
Model Description - Lacustrine	5
Suitability Index (SI) Graphs for Model Variables	8
Riverine Model	16
Lacustrine Model	17
Interpreting Model Outputs	22
ADDITIONAL HABITAT MODELS	22
Model 1	22
Model 2	27
Model 3	27
REFERENCES	27

ACKNOWLEDGMENTS

We would like to thank James Clugston, George Elliot, Jim Winter, William Reynolds, Martha Casterlin, Robert Summerfelt, and Richard Anderson for reviewing the manuscript. Their review contributions are gratefully acknowledged, but the authors accept full responsibility for the contents of the document. Word processing was provided by Dora Ibarra and Carolyn Gulzow. Cover illustration is from Freshwater Fishes of Canada, Bulletin 184, Fisheries Research Board of Canada, by W. B. Scott and E. J. Crossman.

LARGEMOUTH BASS (Micropterus salmoides)

HABITAT USE INFORMATION

General

The largemouth bass (Micropterus salmoides) is native to the eastern United States, excluding the Northeastern States (MacCrimmon and Robbins 1975), and has been introduced throughout the United States (Robbins and MacCrimmon 1974). Two subspecies are recognized, the northern subspecies, M. salmoides salmoides, and the Florida subspecies, M. salmoides floridanus (Bailey and Hubbs 1949; Hart 1952; Ramsey 1975).

Age, Growth, and Food

The maximum known age of largemouth bass is 15 years, and the normal rate of growth for adult largemouth is approximately 454 g per year (Bennett 1937; Anderson 1975). Largemouth bass mature and spawn as early as age I near the southern limit of their range (Morgan 1958; Clugston 1964; La Faunce et al. 1964; Smitherman 1975). Maturity is delayed among more northern populations (Eipper and Regier 1962; Bennett 1971; Carlander 1977). In Canada, maturity is reached in 3-4 years for males and 4-5 years for females (Scott and Crossman 1973).

Largemouth bass fry feed mainly on microcrustaceans and small insects, juveniles consume mostly insects and small fish, and adults feed primarily on fish and crayfish (Emig 1966; Zweiacker and Summerfelt 1974; Carlander 1977). Adults often feed near vegetation within shallow areas. Largemouth bass feeding intensity is bimodal, with peaks in the early morning and late evening (Snow 1971; Olmstead 1974).

Reproduction

Spawning typically begins in the spring when the water temperature reaches 12.0-15.5° C (Mraz 1964; Miller and Kramer 1971). Spawning has been recorded between 11.5 and 29.0° C, but most occurs between 16 and 22° C (Kramer and Smith 1960; Badenhuizen 1969; Carlson and Hale 1972). Incubation time ranges from 2 to 7 days, depending on water temperature (Kramer and Smith 1960; Badenhuizen 1969). The Florida subspecies nests and spawns earlier than the northern subspecies and at a 1 to 3° C lower temperature (Bottroff 1967).

A gravel substrate is preferred for spawning (Newell 1960; Robinson 1961; Mraz 1964), but largemouth bass will nest on a wide variety of other

substrates, including vegetation, roots, sand, mud, and cobble (Harlan and Speaker 1956; Mraz and Cooper 1957; Mraz et al. 1961). Nests are constructed by the male at water depths averaging 0.3-0.9 m, with depths ranging from about 0.15 m to 7.5 m (Swingle and Smith 1950; Harlan and Speaker 1956; Mraz 1964; Clugston 1966; Allan and Romero 1975). Nests have been found as deep as 8.23 m in a reservoir where depth increased during the spawning period (Miller and Kramer 1971).

Specific Habitat Requirements

Lacustrine environments are the preferred habitat of largemouth bass (Emig 1966; Scott and Crossman 1973). Optimal conditions are lakes with extensive ($\geq 25\%$ of surface area) shallow (≤ 6 m depth) areas to support submergent vegetation, yet deep enough (3-15 m mean depth) to successfully overwinter bass (Robbins and MacCrimmon 1974; Carlander 1977; Winter 1977). Thus, it is assumed that 40-60% of the lake area should be > 6 m depth to provide optimal overwintering habitat in northern latitudes.

Optimal riverine habitat for largemouth bass is characterized by large, slow moving rivers or pools of streams with soft bottoms, some aquatic vegetation, and relatively clear water (Trautman 1957; Larimore and Smith 1963; Scott and Crossman 1973). First and second order streams are generally poor habitat (Anderson, personal communication). Deacon (1961) reported that largemouth bass abundance increased in rivers during dry years when the flow was reduced and the water pooled. Thus, it is assumed that a river with a high percent ($\geq 60\%$) of pool and backwater area is optimal. Also, largemouth bass prefer low gradient (≤ 1 m/km) streams; abundance declines as gradient increases toward headwater areas (Finnell et al. 1956; Trautman 1957; Moyle and Nichols 1973). It is assumed that gradients > 4 m/km would be unsuitable.

Growth of largemouth bass is reduced at dissolved oxygen levels < 8 mg/l, and a substantial reduction occurs below 4 mg/l (Stewart et al. 1967). Distress may be evident at 5 mg/l (Katz et al. 1959; Whitmore et al. 1960; Dahlberg et al. 1968; Petit 1973). Levels below 1.0 mg/l are considered lethal (Moss and Scott 1961; Mohler 1966; Petit 1973).

Standing crops of black basses (*Micropterus* spp.) are positively correlated with total dissolved solid levels of 100-350 ppm (Jenkins 1976).

Largemouth bass are considered intolerant of suspended solids (turbidity) and sediment (Muncy et al. 1979). High levels of suspended solids may interfere with reproductive processes and reduce growth. Bulkley (1975) felt that high levels of suspended solids reduced the available food for largemouth bass to the extent that growth of maturing fish was reduced enough that they were physically incapable of reproduction. Buck (1956a; 1956b) found that the greatest survival and growth of largemouth bass occurred in ponds with turbidities (suspended solids) < 25 ppm. Growth was intermediate in ponds with 25-100 ppm turbidity levels and lowest in ponds with turbidities > 100 ppm. Also, no young-of-the-year bass were found in the ponds with high turbidities, while they were recovered from the ponds with low and intermediate turbidities. Thus, optimum suspended solid levels are assumed to be 5-25 ppm, and levels < 5 ppm indicate low productivity.

Largemouth bass require a pH between 5 and 10 for successful reproduction (Swingle 1956; Buck and Thoits 1970). Using Stroud's (1967) criteria for freshwater fish, optimal pH range is 6.5-8.5. Largemouth bass can tolerate short term exposures to pH levels of 3.9 and 10.5 (Calabrese 1969).

Adult. Adult largemouth bass are most abundant in areas with vegetation (Jenkins et al. 1952; Miller 1975) and other forms of cover (e.g., logs, brush, and debris). Optimal cover corresponds to 40-60% of the pool or littoral area; too much cover may reduce prey availability (Saiki and Tash 1979). Adults are most abundant in areas of low current velocity, based on catch data of Kallemyn and Novotny (1977) from the Missouri River. Optimal current velocities are ≤ 6 cm/sec, and velocities > 20 cm/sec are unsuitable based on probability of use curves developed by Hardin and Bovee (1978). Increased water levels in reservoirs may reduce prey availability due to increased cover for prey species. Stable to decreased water levels concentrated prey, which increased feeding and growth rates of adult bass (Heman et al. 1969). Thus, stable to slightly negative midsummer fluctuations (0-3m) are considered optimal for adult largemouth bass.

Optimal temperatures for growth of adult bass range from 24-30° C (Mohler 1966; Coutant 1975; Brungs and Jones 1977; Carlander 1977; Venables et al. 1978). Very little growth occurs below 15° C (Mohler 1966) or above 36° C (Carlander 1977). Salinity levels above 4 ppt cause sharp declines in abundance (Tebo and McCoy 1964). Kilby (1955) found no bass in Mississippi River coastal marshes with salinities above 11.8 ppt, whereas Bailey et al. (1954) reported finding adult largemouth in water with salinities as high as 24.4 ppt in the Escambia River, Alabama and Florida.

Embryo. Optimal spawning substrate is gravel (Newell 1960; Robinson 1961), but other substrates, such as vegetation, roots, sand, and mud, are suitable (Harlan and Speaker 1956; Mraz and Cooper 1957; Mraz et al. 1961). Silty, mucky bottoms are unsuitable (Robinson 1961). Exposed, shallow water (< 1.5 m) nests are vulnerable to destruction by wave action (Miller and Kramer 1971). Boulders, irregular bottoms, and other forms of shelter may protect these nests. Water velocities as low as 40 cm/sec may result in mortality of embryos (Dudley 1969), and Hardin and Bovee (1978) reported that velocities > 10 cm/sec were avoided by the species. Stable water levels during spawning are optimal; drawdowns often result in poor survival (Jester et al. 1969). Since largemouth bass spawn at depths ranging from 0.15 m to 7.5 m, it is assumed that drawdowns > 7.5 m are unsuitable for successful embryonic development during spawning.

Optimal temperatures for successful spawning and incubation are 20-21° C (Clugston 1966; Badenhuizen 1969), with a range of 13-26° C (Carr 1942; Kelley 1968). Survival is very low at temperatures > 30 ° C (Strawn 1961; Kelley 1968; Badenhuizen 1969; McCormick and Wegner 1981) and < 10 ° C (Kramer and Smith 1960). Survival of embryos is impaired at salinities > 1.5 ppt, and zero at levels > 10.5 ppt (Tebo and McCoy 1964).

Fry. Optimal current velocities for fry are < 4 cm/sec (Hardin and Bovee 1978), and fry cannot tolerate current velocities > 27 cm/sec (MacLeod 1967; Laurence 1972). Cover, in the form of flooded terrestrial vegetation, is an

important requirement for fry habitat suitability, because the amount of cover has been positively correlated to number of fry (Aggus and Elliot 1975). However, too much cover constitutes poor spawning habitat (R. O. Anderson, personal communication). Thus, it is assumed that optimal pools or littoral areas contain 40-80% cover. Also, stable to increased summer water level is optimal, because it increases cover availability. It is assumed that decreasing (> 1 m) water levels would be suboptimal because fry would be more susceptible to predation with the decrease in available cover.

Optimal temperatures for fry growth are 27-30° C. Little growth occurs below 15° C or above 32° C (Strawn 1961). The growth rate of fry declined at salinities > 1.66 ppt and was zero at 6 ppt (Tebo and McCoy 1964).

Juvenile. Specific habitat requirements of juveniles are presumed to be similar to those of adult largemouth bass.

HABITAT SUITABILITY INDEX (HSI) MODELS

Model Applicability

Geographic area. The models provided are applicable throughout the 48 conterminous States. The standard of comparison for each individual variable suitability index is the optimal value of the variable that occurs anywhere within this region. Therefore, the models will never provide an HSI of 1.0 when applied to northern waters where temperature related variables do not reach the optimal values found in the southern United States.

Season. The models provide a rating for a body of water based on its ability to support a reproducing population of largemouth bass through all seasons of the year.

Cover types. The models are applicable in riverine, lacustrine, palustrine, and estuarine habitats, as described by Cowardin et al. (1979).

Minimum habitat area. Minimum habitat area is defined as the minimum area of contiguous suitable habitat that is required for a species to live and reproduce. No attempt has been made to establish a minimum habitat size for largemouth bass.

Verification level. The acceptable output for each of these largemouth bass models is an index between 0 and 1 which the author believes has a positive relationship to carrying capacity. In order to verify that the model output is acceptable, the author developed sample data sets from which HSI's were calculated. These sample data sets and their relationship to model verification will be discussed in greater detail following presentation of the model.

Model Description - Riverine

The structure of the riverine HSI model is depicted graphically in Figure 1.

Food component. Percent bottom cover (V_3 , V_4) is assumed to be important because bottom cover provides habitat for aquatic insects, crayfish, and forage fish, which are the predominant food items of largemouth bass. Percent pool and backwater area (V_1) is included to quantify the amount of food habitat.

Cover component. Percent bottom cover (V_3 , V_4) is included because largemouth bass are most abundant in areas with cover. Percent pool and backwater area (V_1) quantifies the amount of cover habitat. Water level fluctuation (V_{16} , V_{18}) is assumed to be important because the amount of available cover is dependent on fluctuations.

Water quality component. The water quality component is limited to dissolved oxygen (V_6), pH range (V_7), temperature (V_8 , V_{10}), turbidity (V_{11}), and salinity (V_{12} , V_{13}) measurements. These parameters have been shown to affect growth or survival. Variables related to temperature and oxygen are assumed to be limiting when they reach near lethal levels. Toxic substances are not considered in this model.

Reproductive component. Temperature (V_9) and salinity (V_{14}) during spawning and embryonic development describe water quality conditions which affect reproduction. Maximum water level fluctuation (V_{17}) is included because optimal development and survival is dependent on stable water levels during spawning. Current velocity (V_{20}) is important because embryos require areas of little or no velocity. Percent pool and backwater area (V_1) quantifies the amount of low velocity spawning areas.

Other component. The variables which are in the other component are those which also describe habitat suitability for the largemouth bass, yet are not specifically related to the life requisite components already presented. Stream gradient (V_{22}) is included because largemouth bass prefer slow moving streams. Current velocity (V_{19} , V_{21}) is an alternative way of describing the habitat suitability of a riverine environment, because of the positive relationship between gradient and current velocity.

Model Description - Lacustrine

Lacustrine model structure is depicted graphically in Figure 2.




Figure 1. Tree diagram illustrating conceptual structure of riverine model for largemouth bass.




Figure 2. Tree diagram illustrating conceptual structure of lacustrine model for largemouth bass.

Food component. Average TDS (V_5) is included because there is a positive correlation between standing crops of black bass and TDS levels, presumably due to the greater amount of food organisms produced at higher TDS levels.

Cover component. Percent bottom cover (V_3, V_4) is included because largemouth bass are most abundant in areas with cover. Percent lacustrine area ≤ 6 m depth (V_2) quantifies the amount of cover habitat. It also serves to quantify the amount of overwintering habitat in northern latitudes. Water level fluctuation (V_{16}, V_{18}) is assumed to be important because the amount of available cover is dependent on fluctuations.

Water quality component. Same explanation as presented in the riverine model description.


Reproduction component. Temperature (V_9) and salinity (V_{14}) for spawning and embryonic development describe water quality conditions which affect reproduction. Substrate (V_{15}) is important for spawning success. Water level fluctuation (V_{17}) is included because optimal development and survival of embryos is dependent on stable water levels during spawning. Percent lacustrine area ≤ 6 m depth (V_2) quantifies the amount of spawning habitat.

Suitability Index (SI) Graphs for Model Variables

This section contains suitability index graphs for the 22 variables described above and equations for combining selected variable indices into a species HSI using the component approach. The "R" and "L" refer to riverine and lacustrine habitat variables, respectively.

<u>Habitat</u>	<u>Variable</u>	
R	V_1	Percent pool and back-water area during average summer flow.

Suitability Graph




L V₂

Percent lacustrine area \leq 6 m depth.


Note: A regional consideration is made for bodies of water in more southern latitudes where overwintering requirements are not as rigorous.

- A. Northern latitudes
- B. Southern latitudes




R,L V₃

Percent bottom cover (e.g., aquatic vegetation, logs, and debris) within pools, backwaters, or littoral areas during summer. (Adult, Juvenile)




R,L V₄

Percent bottom cover (e.g., aquatic vegetation, logs, and debris) within pools, backwaters, or littoral areas during summer. (Fry)




L V₅ Average TDS concentration during growing season when carbonate-bicarbonate > sulfate-chloride ionic concentration. If sulfate-chloride concentration exceeds carbonate-bicarbonate, reduce SI rating for TDS by 0.2.




R,L V₆ Minimum dissolved oxygen levels during midsummer within pools or littoral areas.

A) Frequently < 2 mg/l
 B) Usually ≥ 2 and < 5 mg/l
 C) Usually ≥ 5 and < 8 mg/l
 D) Often > 8 mg/l




R,L V₇ pH range during growing season.


A) < 5.0 or > 10.0
 B) ≥ 5 and < 6.5 or > 8.5 and ≤ 10.0
 C) 6.5-8.5




R,L V_8 Average water temperature within pools, backwaters, or littoral areas during the growing season. (Adult, Juvenile)



R,L V_9 Average weekly mean temperature within pools or littoral areas during spawning and incubation. (Embryo)




R,L V_{10} Average water temperature within pools, backwaters, or littoral areas during the growing season. (Fry)




R,L V_{11} Maximum monthly average turbidity (suspended solids) during growing season.


A) 5-25 ppm
 B) > 25 and ≤ 100 ppm
 C) < 5 ppm, > 100 ppm




R,L V_{12} Maximum salinity during summer. (Adult, Juvenile)



R,L V_{13} Maximum salinity during summer. (Fry)




R,L V₁₄ Maximum salinity during spawning and incubation. (Embryo)




R,L V₁₅ Substrate composition within riverine pools and backwaters or lacustrine littoral areas. (Embryo)


- A) Boulders and bedrock predominate ($\geq 50\%$)
- B) Sand (0.062-2.0 mm) predominates
- C) Silt and clay (0.0-0.004 mm) predominate
- D) Gravel (0.2-6.4 cm) predominates




R,L V₁₆ Average water level fluctuation during growing season. (Adult, Juvenile)




R,L V_{17} Maximum water level fluctuation during spawning. (Embryo)




R,L V_{18} Average water level fluctuation during growing season. (Fry)




R V_{19} Average current velocity at 0.6 depth during summer. (Adult, Juvenile)




R V_{20} Maximum current velocity at 0.8 depth within pools or backwaters during spawning (May-June). (Embryo)



R V_{21} Average current velocity at 0.6 depth during summer. (Fry)



R V_{22} Stream gradient within representative reach.



Riverine Model

These equations utilize the life requisite approach and consist of five components: food, cover, water quality, reproduction, and other.

Food (C_F).

$$C_F = \left[V_1 \times \frac{(V_3 + V_4)}{2} \right]^{1/2}$$

Cover (C_C).

$$C_C = \left[V_1 \times \frac{(V_3 + V_4)}{2} \times \frac{(V_{16} + V_{18})}{2} \right]^{1/3}$$

Water Quality (C_{WQ}).

If V_{12} and $V_{13} = 1.0$,

$$C_{WQ} = \frac{2V_6 + V_7 + 2V_8 + V_{10} + V_{11}}{7}$$

If V_{12} or $V_{13} < 1.0$,

$$C_{WQ} = \frac{2V_6 + V_7 + 2V_8 + V_{10} + V_{11} + \frac{(V_{12} + V_{13})}{2}}{8}$$

If V_6 , V_8 , or $V_{10} \leq 0.4$, C_{WQ} equals the lowest of the following: V_6 , V_8 , V_{10} , or the above equation.

Reproduction (C_R).

If $V_{14} = 1.0$,

$$C_R = (V_1 \times V_9 \times V_{15} \times V_{17} \times V_{20})^{1/5}$$

If $V_{14} < 1.0$,

$$C_R = (V_1 \times V_9 \times V_{14} \times V_{15} \times V_{17} \times V_{20})^{1/6}$$

Other (C_{OT}).

Note: Since there is a correlation between stream gradient and current velocity, the user has two options for the "other" component:

A) $C_{OT} = \frac{V_{19} + V_{21}}{2}$, or

B) $C_{OT} = V_{22}$

HSI determination.

$$HSI = (C_F \times C_C \times C_{WQ} \times C_R \times C_{OT})^{1/5}$$

If C_{WQ} or C_R is ≤ 0.4 , then the HSI equals the lowest of the following: C_{WQ} , C_R or the above equation.

Lacustrine Model

These equations utilize the life requisite approach and consist of four components: food, cover, water quality, and reproduction.

Food (C_F).

$$C_F = V_5$$

Cover (C_C).

$$C_C = \left[V_2 \times \frac{(V_3 + V_4)}{2} \times V_{16} \times V_{18} \right]^{1/4}$$

Water Quality (C_{WQ}).

Same as the riverine habitat suitability index equations for water quality.

Reproduction (C_R).

If $V_{14} = 1.0$,

$$C_R = (V_2 \times V_9 \times V_{15} \times V_{17})^{1/4}$$

If $V_{14} < 1.0$,

$$C_R = (V_2 \times V_9 \times V_{14} \times V_{15} \times V_{17})^{1/5}$$

HSI determination.

$$HSI = (C_F \times C_C \times C_{WQ} \times C_R)^{1/4}$$

If C_{WQ} or C_R is ≤ 0.4 , then the HSI equals the lowest of the following: C_{WQ} ; C_R ; or the above equation.

Sources of data and assumptions made in developing the suitability indices are presented in Table 1.

Table 1. Data sources and assumptions for largemouth bass suitability indices.

	Variable and source	Assumption
V ₁	Trautman 1957 Deacon 1961 Larimore and Smith 1963 Branson 1967 Scott and Crossman 1973 Funk 1975	Largemouth bass typically inhabit pool and backwater areas of streams; optimal habitat consists of at least 60% pool/backwater area.
V ₂	Robbins and MacCrimmon 1974 Carlander 1977 Winter 1977	Cover adequate to support large populations can be provided by greater than 25% area \leq 6 m depth. However, lacustrine habitats in northern latitudes need to be deep enough to successfully overwinter bass.
V ₃	Jenkins et al. 1952 Miller 1975 Saiki and Tash 1979	Adult largemouth bass are most abundant in areas which contain cover; too much cover may reduce prey availability.
V ₄	Kramer and Smith 1960 Newell 1960 Aggus and Elliot 1975 Anderson 1981	The amount of cover has been positively correlated with the number of fry. Too much cover constitutes poor spawning and rearing habitat.
V ₅	Jenkins 1976	Total dissolved solids (TDS) levels correlated with high standing crops are optimal; those correlated with lower standing crops are suboptimal. The data used to develop this curve are primarily from southeastern reservoirs.
V ₆	Katz et al. 1959 Whitmore et al. 1960 Moss and Scott 1961 Mohler 1966 Stewart et al. 1967 Dahlberg et al. 1968 Petit 1973	Dissolved oxygen levels where growth is not impaired are optimal, those where growth is reduced are suboptimal, and those where death may occur are unsuitable.
V ₇	Swingle 1956 Stroud 1967 (FW) ^a Calabrese 1969 Buck and Thoits 1970	Optimal pH range is presumably the same as those for all freshwater fish. Levels which impair growth of largemouth bass are suboptimal; those which can result in death are unsuitable.

Table 1. (continued).

Variable and source	Assumption
V ₈ Hart 1952 Johnson and Charlton 1960 Mohler 1966 Coutant 1975 Venables et al. 1978	Temperatures where growth of adult and juvenile largemouth bass is highest are optimal, those where growth is reduced are suboptimal, and those where little or no growth occurs are unsuitable.
V ₉ Carr 1942 Kramer and Smith 1960 Strawn 1961 Clugston 1966 Kelley 1968 Badenhuisen 1969	Temperatures which result in greatest survival of embryos are optimal, those where survival is reduced are suboptimal, and those where little or no survival occurs are unsuitable.
V ₁₀ Strawn 1961	Same assumption as V ₉ , only applied to fry life stage.
V ₁₁ Buck 1956a; 1956b Bulkley 1975 Muncy et al. 1979	Turbidity (suspended solid) levels where the greatest survival and growth occur are optimal. Levels which reduce growth and interfere with reproductive processes are suboptimal.
V ₁₂ Bailey et al. 1954 Tebo and McCoy 1964	Salinity levels where adult and juvenile largemouth bass are most abundant are optimal. Those where abundance declines are suboptimal to unsuitable.
V ₁₃ Tebo and McCoy 1964	Salinity levels where the growth rate of fry are not impaired are optimal, those where growth rates decline are sub-optimal, and those where no growth occurs are unsuitable.
V ₁₄ Tebo and McCoy 1964	Salinity levels which do not affect embryonic survival are optimal, those where survival is impaired are sub-optimal, and levels where no survival occurs are unsuitable.
V ₁₅ Harlan and Speaker 1956 Mraz and Cooper 1957 Newell 1960 Mraz et al. 1961 Robinson 1961 Mraz 1964	The substrate type on which most spawning takes place is optimal; other substrate types are suboptimal.

Table 1. (concluded).

Variable and source	Assumption
V ₁₆ Heman et al. 1969	Water level fluctuations which concentrate prey and lead to increases in growth rates of adult and juvenile largemouth bass are optimal; those which reduce prey availability are suboptimal.
V ₁₇ Harlan and Speaker 1956 Mraz 1964 Clugston 1966 Jester et al. 1969 Allan and Romero 1975	Fluctuations in water level which do not affect survival of embryos are optimal; those fluctuations which exceed the average depth of nests (and reduce survival) are suboptimal to unsuitable.
V ₁₈ Aggus and Elliot 1975	Water level fluctuations which lead to increased cover availability for fry are optimal. Those which decrease the amount of cover are suboptimal.
V ₁₉ Bailey et al. 1954 Kallemeyn and Novotny 1977 Hardin and Bovee 1978	Current velocities where abundance of adult and juvenile largemouth bass is greatest are optimal; those where abundance declines are suboptimal to unsuitable.
V ₂₀ Deacon 1961 Dudley 1969	Velocities which do not impair embryonic survival are optimal, those which reduce survival are suboptimal to unsuitable.
V ₂₁ MacLeod 1967 Laurence 1972 Hardin and Bovee 1978	Same assumption as V ₁₉ , only applicable to fry life stage.
V ₂₂ Finnell et al. 1956 Trautman 1957 Moyle and Nichols 1973	Gradients where species abundance is greatest are optimal; those which lead to a decline in abundance are suboptimal to unsuitable.

^a(FW) = Freshwater fish; remaining citations are largemouth bass data.

Sample data sets from which HSI's have been generated using the riverine HSI equations are presented in Table 2. Similar sets using the lacustrine HSI equations are given in Table 3. The data sets are not actual field measurements, but represent combinations of variable values we believe could occur in a riverine or lacustrine habitat. The HSI's calculated from the data reflect what we believe carrying capacity trends would be in riverine and lacustrine habitats with the listed characteristics. Accuracy of the models in predicting population trends has not been tested.

Interpreting Model Outputs

The largemouth bass HSI determined by use of these models will not necessarily represent the population of largemouth bass in the study area. Habitats with an HSI of 0 may contain some largemouth bass; habitats with a high HSI may contain few. This is because the population of a study area of a stream or lake does not totally depend on the habitat variables, as is assumed by the model. If the models are a good representation of largemouth bass habitat, then in riverine and lacustrine environments where largemouth bass population levels are due primarily to habitat related factors, the models should be positively correlated to the long term average population levels. However, this has not been tested. The proper interpretation of the HSI is one of comparison. If two riverine or lacustrine habitats have different HSI's, the one with the higher HSI should have the potential to support more largemouth bass than the one with the lower HSI, given the model assumptions have not been violated.

This model does not specifically address the effects of wind induced turbulence on bass reproductive success, but wave destruction of nests may be locally important (Miller and Kramer 1971; Summerfelt 1975). The direction of prevailing winds, surrounding topography, lake morphometry, and the placement of objects which might provide shelter for nests (e.g., boulders and ledges) may be relevant criteria in specific cases.

ADDITIONAL HABITAT MODELS

Model 1

Assuming water quality is adequate, optimal riverine habitat for largemouth bass may be characterized as follows: large, low (≤ 1 m/km) gradient streams; abundant (40-80% of pool and backwater area) cover in the form of aquatic vegetation, brush, logs, or other cover items; warm (24-30° C) mid-summer water temperatures; low (< 25 ppm) turbidity; and a predominance (> 60% stream area) of pools.

$$HSI = \frac{\text{Number of above criteria present}}{5}$$

Table 2. Sample data sets using riverine HSI model.

Variable		Data set 1		Data set 2		Data set 3	
		Data	SI	Data	SI	Data	SI
% pool, backwater area	V ₁	20	0.1	40	0.5	70	1.0
% bottom cover- adult, juvenile	V ₂	10	0.4	25	0.7	60	1.0
% bottom cover	V ₄	10	0.2	25	0.6	60	0.8
Dissolved O ₂ (mg/l)	V ₆	3.0	0.4	5.1	0.8	6.4	0.8
pH range	V ₇	7.8	1.0	8.1	1.0	7.5	1.0
Average temperature- adult, juvenile (°C)	V ₈	22	0.8	23	0.9	27	1.0
Average temperature- embryo (°C)	V ₉	19	0.8	22	1.0	24	0.5
Average temperature- fry (°C)	V ₁₀	20	0.4	23	0.7	25	0.9
Maximum turbidity- JTU	V ₁₁	150	0.3	25	1.0	40	0.7
Maximum salinity- adult, juvenile (ppt)	V ₁₂	1.2	1.0	2.1	1.0	0.8	1.0
Maximum salinity- fry (ppt)	V ₁₃	1.2	1.0	2.1	1.0	0.8	1.0
Maximum salinity- embryo (ppt)	V ₁₄	1.2	1.0	2.1	0.9	0.8	1.0
Substrate type- embryo	V ₁₅	silt/clay	0.8	sand	0.5	gravel	1.0

Table 2. (concluded).

Variable	Data set 1		Data set 2		Data set 3		
	Data	SI	Data	SI	Data	SI	
Water level fluctuation-adult, juvenile	V_{16}	-0.5	1.0	0	1.0	0	1.0
Water level fluctuation-embryo (m)	V_{17}	-1.0	0.9	-0.5	0.9	0	1.0
Water level fluctuation-fry (m)	V_{18}	-0.6	1.0	0	1.0	0	1.0
Current velocity-embryo (cm/sec)	V_{20}	0.8	1.0	0.5	1.0	0.2	1.0
Stream gradient (m/km)	V_{22}	2	0.7	1.2	0.9	0.7	1.0
Component SI							
$C_F =$			0.2		0.6		1.0
$C_C =$			0.3		0.7		1.0
$C_{WQ} =$			0.6		0.9		0.9
$C_R =$			0.6		0.8		0.9
$C_{OT} =$			0.7		0.9		1.0
HSI =			0.4		0.8		1.0

Table 3. Sample data sets using lacustrine HSI model.

Variable		Data set 1		Data set 2		Data set 3	
		Data	SI	Data	SI	Data	SI
% lacustrine area ≤ 6 m depth	V ₂	5	0.2	15	0.6	65	0.9
% bottom cover adult, juvenile	V ₃	5	0.3	30	0.8	75	0.8
% bottom cover (fry)	V ₄	5	0.1	30	0.7	75	1.0
Average TDS (ppm)	V ₅	50	0.4	20	0.2	200	1.0
Dissolved O ₂ (mg/l)	V ₆	6.5	0.8	7.0	0.8	6.4	0.8
pH range	V ₇	8.2	1.0	8.7	0.5	7.9	1.0
Average temperature- adult, juvenile (°C)	V ₈	21	0.7	28	1.0	26	1.0
Average temperature- embryo (°C)	V ₉	20	1.0	25	0.3	23	0.8
Average temperature- fry (°C)	V ₁₀	22	0.6	27	1.0	25	0.9
Maximum turbidity (ppm)	V ₁₁	15	1.0	70	0.7	25	1.0
Maximum salinity adult, juvenile (ppt)	V ₁₂	1.3	1.0	1.5	1.0	0.5	1.0
Maximum salinity (ppt)	V ₁₃	1.3	1.0	1.5	1.0	0.5	1.0

Table 3. (concluded).

Variable		Data set 1		Data set 2		Data set 3	
		Data	SI	Data	SI	Data	SI
Maximum salinity (ppt)	V ₁₅	1.3	1.0	1.5	1.0	0.5	1.0
Substrate type-embryo	V ₁₅	sand	0.5	gravel	1.0	silt/clay	0.8
Water level fluctuation-adult, embryo (m)	V ₁₆	-5	0.7	0	1.0	0	1.0
Water level fluctuation-embryo (m)	V ₁₇	+2	0.9	+0.3	1.0	0	1.0
Water level fluctuation-fry (m)	V ₁₈	-5	0.3	0	1.0	0	1.0
Component SI							
C _F =			0.4		0.2		1.0
C _C =			0.3		0.8		0.9
C _{WQ} =			0.8		0.8		0.9
C _R =			0.5		0.7		0.9
HSI =			0.5		0.5		0.9

Model 2

Assuming water quality is adequate, optimal lacustrine habitat for largemouth bass may be characterized as follows: fertile (TDS levels 100-350 ppm) lakes; abundant (40-80% of littoral area) bottom cover; warm (24-30° C) mid-summer water temperatures; and extensive (25-60% for northern latitudes; ≥ 25% for southern latitudes) shallow (≤ 6 m depth) areas.

$$HSI = \frac{\text{Number of above criteria present}}{4}$$

Model 3

Use the regression models for largemouth bass standing crop in reservoirs presented by Aggus and Morais (1979) to calculate an HSI.

REFERENCES

- Aggus, L. R., and G. J. Elliot. 1975. Effects of cover and food on year-class strength of largemouth bass. Pages 317-322 in H. Clepper, ed. Black bass biology and management. Sport Fish. Inst., Washington, D. C.
- Aggus, L. R., and D. I. Morais. 1979. Habitat suitability index equations for reservoirs based on standing crop of fish. Natl. Reservoir Res. Prog. Rep. to U.S. Fish Wildl. Serv., Habitat Evaluation Proj., Ft. Collins, Colorado. 120 pp.
- Allan, R. C., and J. Romero. 1975. Underwater observations of largemouth bass spawning and survival in Lake Mead. Pages 104-112 in H. Clepper, ed. Black bass biology and management. Sport Fish. Inst., Washington, D.C.
- Anderson, R. O. 1975. Factors influencing the quality of largemouth bass fishing. Pages 183-194 in H. Clepper, ed. Black bass Biology and Management. Sport Fish. Inst., Washington, D. C.
- _____. 1981. Personal communication. Missouri Cooperative Fishery Unit, Columbia, Missouri.
- Badenhuizen, T. R. 1969. Effect of incubation temperature on mortality of embryos of the largemouth bass Micropterus salmoides (Lacepede). M.S. Thesis, Cornell Univ., Ithaca, New York. 88 pp.
- Bailey, R. M., and C. L. Hubbs. 1949. The black basses (Micropterus) of Florida, with description of a new species. Univ. Michigan Mus. Zool., Occas. Pap. 516. 43 pp.

- Bailey, R. M., H. E. Winn, and C. L. Smith. 1954. Fishes from the Escambia River, Alabama and Florida, with ecologic and taxonomic notes. Proc. Acad. Nat. Sci., Philadelphia 106:109-164.
- Bennett, G. W. 1937. The growth of the largemouthed black bass, Huro salmoides (Lacepede), in the waters of Wisconsin. Copeia 1937(2):104-118.
- _____. 1971. Management of lakes and ponds. 2nd Ed. Van Nostrand Reinhold Co., New York. 375 pp.
- Bottroff, L. J. 1967. Integradation of Florida bass in San Diego County, California. M.S. Thesis, San Diego State Coll., San Diego, CA. 135 pp.
- Branson, B. A. 1967. Fishes of the Neosho River system in Oklahoma. Am. Midl. Nat. 78:126-154.
- Brungs, W. A., and B. R. Jones. 1977. Temperature criteria for freshwater fish: protocol and procedures. U.S. Environ. Protection Agency, Environ. Res. Lab. Ecol. Res. Ser. EPA-600/3-77-061. 139 pp.
- Buck, D. H. 1956a. Effects of turbidity on fish and fishing. Oklahoma Fish. Res. Lab. Rep. 56. 62 pp.
- _____. 1956b. Effects of turbidity on fish and fishing. Trans. N. Am. Wildl. Conf. 21:249-261.
- Buck, D. H., and C. F. Thoits, III. 1970. Dynamics of one-species populations of fishes in ponds subjected to cropping and additional stocking. Illinois Nat. Hist. Surv. Bull. 30:68-165.
- Bulkley, R. V. 1975. Chemical and physical effects on the centrarchid basses. Pages 286-294 in H. Clepper, ed. Black bass biology and management. Sport Fish. Inst., Washington, D. C.
- Calabrese, A. 1969. Effect of acids and alkalies on survival of bluegills and largemouth bass. U.S. Bur. Sport Fish. Wildl. Tech. Paper 42. 10 pp.
- Carlander, K. D. 1977. Largemouth bass. Pages 200-275 in Handbook of freshwater fishery biology. Iowa State Univ. Press, Ames. Vol. 2.
- Carlson, A. R., and J. G. Hale. 1972. Successful spawning of largemouth bass Micropterus salmoides (Lacepede) under laboratory conditions. Trans. Am. Fish. Soc. 101:539-542.
- Carr, M. H. 1942. The breeding habits, embryology and larval development of the largemouthed black bass in Florida. Proc. New England Zool. Club 20:43-77.

- Clugston, J. P. 1964. Growth of the Florida largemouth bass, Micropterus salmoides floridanus (Lesueur), and the northern largemouth bass, M. s. salmoides (Lacepede), in subtropical Florida. *Trans. Am. Fish. Soc.* 93:146-154.
- _____. 1966. Centrarchid spawning in the Florida everglades. *Q. J. Florida Acad. Sci.* 29:137-144.
- Coutant, C. C. 1975. Responses of bass to natural and artificial temperature regimes. Pages 272-285 in H. Clepper, ed. *Black bass biology and management.* Sport Fish. Inst., Washington, D.C.
- Cowardin, L. M., V. Carter, F. C. Golet, and E. T. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. *USDI Fish Wildl. Serv., FWS/OBS-79/31.* 103 p.
- Dahlberg, M. L., D. L. Shumway, and P. Doudoroff. 1968. Influence of dissolved oxygen and carbon dioxide on swimming performance of largemouth bass and coho salmon. *J. Fish. Res. Board Can.* 25:49-70.
- Deacon, J. E. 1961. Fish populations, following a drought, in the Neosho and Marais des Cygnes Rivers of Kansas. *Univ. Kansas Mus. Nat. Hist. Publ.* 13:359-427.
- Dudley, R. G. 1969. Survival of largemouth bass embryos at low dissolved oxygen concentrations. M.S. Thesis, Cornell Univ., Ithaca, New York. 61 pp.
- Eipper, A. W., and H. A. Regier. 1962. Fish management in New York farm ponds. *Cornell Ext. Bull.* 1089. 40 pp.
- Emig, J. W. 1966. Largemouth bass. Pages 332-353 in A. Calhoun, ed. *Inland fisheries management.* California Fish Game.
- Finnell, J. C., R. M. Jenkins, and G. E. Hall. 1956. The fishery resources of the Little River system, McCurtain County, Oklahoma. *Oklahoma Fish. Res. Lab. Rep.* 55. 82 pp.
- Funk, J. L. 1975. Structure of fish communities in streams which contain bass. Pages 140-153 in H. Clepper, ed. *Black bass biology and management.* Sport Fish. Inst., Washington, D.C.
- Hardin, T., and K. Bovee. 1978. Largemouth bass. Instream Flow Group, U.S. Fish Wildl. Serv., Western Energy and Land Use Team, Ft. Collins, Colorado. Unpublished data.
- Harlan, J. R., and E. B. Speaker. 1956. *Iowa fish and fishing.* 3rd Ed. State of Iowa. 377 pp.
- Hart, J. S. 1952. Geographic variations of some physiological and morphological characters in certain freshwater fish. *Univ. of Toronto Biol. Ser.* 60, *Publ. Ont. Fish. Res. Lab.* 72. 79 pp.


- Heman, M. L., R. S. Campbell, and L. C. Redmond. 1969. Manipulation of fish populations through reservoir drawdown. *Trans. Am. Fish. Soc.* 98:293-304.
- Jenkins, R. M. 1976. Prediction of fish production in Oklahoma reservoirs on the basis of environmental variables. *Ann. Oklahoma Sc.* 5:11-20.
- Jenkins, R. M., E. M. Leonard, and G. E. Hall. 1952. An investigation of the fisheries resources of the Illinois River and pre-impoundment study of Tenkiller Reservoir, Oklahoma. *Oklahoma Fish. Res. Lab. Rep.* 26. 136 pp.
- Jester, D. B., T. M. Moody, C. Sanchez, Jr., and D. E. Jennings. 1969. A study of game fish reproduction and rough fish problems in Elephant Butte Lake. *New Mexico Job Compl. Rep. Fed. Aid. Proj. F-22R-9, Job F-1.* 73 pp. (Cited by Carlander 1977).
- Johnson, M. G., and W. H. Charlton. 1960. Some effects of temperature on the metabolism and activity of largemouth bass, Micropterus salmoides Lacepede. *Prog. Fish. Cult.* 22:155-163.
- Kallemeyn, L. W., and J. F. Novotny. 1977. Fish and fish food organisms in various habitats of the Missouri River in South Dakota, Nebraska, and Iowa. *USDI Fish Wildl. Serv., FWS/OBS-77/25.*
- Katz, M., A. Pritchard, and C. E. Warren. 1959. Ability of some salmoides and a centrarchid to swim in water of reduced oxygen content. *Trans. Am. Fish. Soc.* 88:88-95.
- Kelley, J. W. 1968. Effects of incubation temperature on survival of largemouth bass eggs. *Prog. Fish-Cult.* 30:159-163.
- Kilby, J. D. 1955. The fishes of two gulf coastal marsh areas of Florida. *Tulane studies in Zoology* 2:175-247.
- Kramer, R. H., and L. L. Smith, Jr. 1960. First-year growth of the largemouth bass, Micropterus salmoides (Lacepede), and some related ecological factors. *Trans. Am. Fish. Soc.* 89:222-233.
- La Faunce, D. A., J. B. Kimsey, and H. K. Chadwick. 1964. The fishery at Sutherland Reservoir, San Diego County, California. *California Fish Game* 50:271-291.
- Larimore, R. W., and P. W. Smith. 1963. The fishes of Champaign County, Illinois, as affected by 60 years of stream changes. *Illinois Nat. Hist. Surv. Bull.* 28:299-382.
- Laurence, G. C. 1972. Comparative swimming abilities of fed and starved larval largemouth bass (Micropterus salmoides). *J. Fish. Biol.* 4(1):73-78.
- MacCrimmon, H. R., and W. H. Robbins. 1975. Distribution of the black basses in North America. Pages 56-66 in H. Clepper, ed. *Black bass biology and management.* Sport Fish. Inst., Washington, D.C.

- MacLeod, J. C. 1967. A new apparatus for measuring maximum swimming speeds of small fish. *J. Fish. Res. Board Can.* 24:1241-1252.
- McCormick, J. H., and J. A. Wegner. 1981. Responses of largemouth bass from different latitudes to elevated water temperatures. *Trans. Am. Fish. Soc.* 110:417-429.
- Miller, K. D., and R. H. Kramer. 1971. Spawning and early life history of largemouth bass (*Micropterus salmoides*) in Lake Powell. Pages 73-83 in G. E. Hall. *Reservoir fisheries and limnology*. Am. Fish. Soc. Spec. Publ. 8.
- Miller, R. J. 1975. Comparative behavior of centrarchid basses. Pages 85-94 in H. Clepper, ed. *Black bass biology and management*. Sport Fish. Inst., Washington, D.C.
- Mohler, S. H. 1966. Comparative seasonal growth of the largemouth, spotted and smallmouth bass. M.S. Thesis, Univ. of Missouri, Columbia, Mo. 99 pp.
- Morgan, G. D. 1958. A study of six different pond stocking ratios of largemouth bass, *Micropterus salmoides* (Lacepede), and bluegill, *Lepomis macrochirus* (Rafinesque); and the relation of the chemical, physical, and biological data to pond balance and productivity. *J. Sci. Lab. Denison Univ.* 44:151-202.
- Moss, D. D., and D. C. Scott. 1961. Dissolved oxygen requirements of three species of fish. *Trans. Am. Fish. Soc.* 90:377-393.
- Moyle, P. B., and R. D. Nichols. 1973. Ecology of some native and introduced fishes of the Sierra Nevada foothills in central California. *Copeia* 1973:478-490.
- Mraz, D. 1964. Observations on large and smallmouth bass nesting and early life history. Wisconsin Conserv. Dept., Res. Rep. 11 (Fisheries). 13 pp.
- Mraz, D., and E. L. Cooper. 1957. Reproduction of carp, largemouth bass, bluegills, and black crappies in small rearing ponds. *J. Wildl. Manage.* 21:127-133.
- Mraz, D., S. Kmiotek, and L. Frankenberger. 1961. The largemouth bass, its life history, ecology and management. Wisconsin Conserv. Dept. Publ. 232. 15 pp.
- Muncy, R. J., G. J. Atchison, R. V. Bulkley, B. W. Menzel, L. G. Perry, and R. C. Summerfelt. 1979. Effects of suspended solids and sediment on reproduction and early life of warmwater fishes: a review. U.S. Environmental Protection Agency EPA-600/3-79-042. 101 pp.

- Newell, A. E. 1960. Biological survey of the lakes and ponds in Coos, Grafton and Carroll Counties. New Hampshire Fish Game Surv. Rep. 8a. 297 pp.
- Olmstead, L. L. 1974. The ecology of largemouth bass (Micropterus salmoides) and spotted bass (Micropterus punctulatus) in Lake Fort Smith, Arkansas. Ph.D. Dissertation, Univ. Arkansas, Fayetteville, Ak. 133 p.
- Petit, G. D. 1973. Effects of dissolved oxygen on survival and behavior of selected fishes of western Lake Erie. Ohio Biol. Surv. Bull. 4(4):1-76.
- Ramsey, J. S. 1975. Taxonomic history and systematic relationships among species of Micropterus. Pages 67-75 in H. Clepper, ed. Black bass biology and management. Sport Fish. Inst., Washington, D.C.
- Robbins, W. H., and H. R. MacCrimmon. 1974. The blackbass in America and overseas. Publ. Div., Biomanagement and Research Enterprises, Ontario. 196 pp.
- Robinson, D. W. 1961. Utilization of spawning box by bass. Prog. Fish-Cult. 23:119.
- Saiki, M. K., and J. C. Tash. 1979. Use of cover and dispersal by crayfish to reduce predation by largemouth bass. Pages 44-48 In D. L. Johnson and R. A. Stein, eds. Response of fish to habitat structure in standing water. N. Central Div. Am. Fish. Soc. Spec. Publ. 6.
- Scott, W. B., and E. J. Crossman. 1973. Freshwater fishes of Canada. Fish. Res. Board Can. Bull. 184. pp. 734-740.
- Smitherman, R. O. 1975. Experimental species associations of basses in Alabama ponds. Pages 76-84 in H. Clepper, ed. Black bass biology and management. Sport Fish. Inst., Washington, D.C.
- Snow, H. E. 1971. Harvest and feeding habits of largemouth bass in Murphy Flowage, Wisconsin. Wisc. Dept. Nat. Resour. Tech. Bull. 50. 25 p.
- Stewart, N. E., D. L. Shumway, and P. Doudoroff. 1967. Influence of oxygen concentration on the growth of juvenile largemouth bass. J. Fish. Res. Board Can. 24:475-494.
- Strawn, K. 1961. Growth of largemouth bass fry at various temperatures. Trans. Am. Fish. Soc. 90:334-335.
- Stroud, R. H. 1967. Water quality criteria to protect aquatic life: a summary. Am. Fish. Soc. Spec. Publ. 4:33-37.
- Summerfelt, R. C. 1975. Relationship between weather and year-class strength of largemouth bass. Pages 166-174 in H. Clepper, ed. Black bass biology and management. Sport Fish. Inst., Washington, D.C.
- Swingle, H. S. 1956. Determination of balance in farm fish ponds. Trans. N. Am. Wildl. Conf. 21:298-322.

- Swingle, H. S., and E. V. Smith. 1950. Factors affecting the reproduction of bluegill bream and largemouth black bass in ponds. Agr. Exp. Stn. Alabama Polytechnic Inst. Circ. 87. 8 pp.
- Tebo, L. B., Jr., and E. G. McCoy. 1964. Effect of sea-water concentration on the reproduction and survival of largemouth bass and bluegills. Prog. Fish-Cult. 26:99-106.
- Trautman, M. B. 1957. The fishes of Ohio. Ohio State Univ. Press, Columbus, Oh. 683 pp.
- Venables, B. J., L. D. Fitzpatrick, and W. D. Pearson. 1978. Laboratory measurement of preferred body temperature of adult largemouth bass (Micropterus salmoides). Hydrobiologia 58(1):33-36.
- Whitmore, C. M., C. E. Warren, and P. Doudoroff. 1960. Avoidance reactions of salmonid and centrarchid fishes to low oxygen concentrations. Trans. Am. Fish. Soc. 89:17-26.
- Winter, J. D. 1977. Summer home range movements and habitat use by four largemouth bass in Mary Lake, Minnesota. Trans. Am. Fish. Soc. 106:323-330.
- Zweiacker, P. L., and R. C. Summerfelt. 1974. Seasonal variation in food and diet periodicity in feeding of northern largemouth bass, Micropterus salmoides salmoides (Lacepede), in an Oklahoma reservoir. Proc. Southeastern Assoc. Game and Fish Comm. 27(1973):579-591.

REPORT DOCUMENTATION PAGE		1. REPORT NO. FWS/OBS-82/10.16	2.	3. Recipient's Accession No.
4. Title and Subtitle Habitat Suitability Index Models: Largemouth bass			5. Report Date July 1982	
7. Author(s) Robert J. Stuber, Glen Gebhart and O. Eugene Maughan			6.	
9. Performing Organization Name and Address Habitat Evaluation Procedures Group Western Energy and Land Use Team U.S. Fish and Wildlife Service Drake Creekside Building One 2625 Redwing Road Fort Collins, Colorado 80526			8. Performing Organization Rept. No.	
12. Sponsoring Organization Name and Address Western Energy and Land Use Team Office of Biological Services Fish and Wildlife Service U.S. Department of the Interior Washington, DC 20240			10. Project/Task/Work Unit No.	
15. Supplementary Notes			11. Contract(C) or Grant(G) No. (C) (G)	
16. Abstract (Limit: 200 words) This is one of a series of publications that provide information on the habitat requirements of selected fish and wildlife species. Literature describing the relationship between habitat variables related to life requisites and habitat suitability for the Largemouth bass (<u>Micropterus salmoides</u>) are synthesized. These data are subsequently used to develop Habitat Suitability (HSI) models. The HSI models are designed to provide information that can be used in impact assessment and habitat management.			13. Type of Report & Period Covered	
17. Document Analysis a. Descriptors ·Animal behavior · Mathematical models ·Animal ecology ·Bass ·Fishes ·Habitability b. Identifiers/Open-Ended Terms Largemouth bass Habitat Suitability Index models <u>Micropterus salmoides</u> Habitat requirements Habitat suitability Species-habitat relationships Habitat preference Impact assessment Habitat management c. COSATI Field/Group			14.	
18. Availability Statement Unlimited		19. Security Class (This Report) UNCLASSIFIED		21. No. of Pages ii - v + 32 pp
		20. Security Class (This Page) UNCLASSIFIED		22. Price



REGION 1

Regional Director
 U.S. Fish and Wildlife Service
 Lloyd Five Hundred Building, Suite 1692
 500 N.E. Multnomah Street
 Portland, Oregon 97232

REGION 2

Regional Director
 U.S. Fish and Wildlife Service
 P.O. Box 1306
 Albuquerque, New Mexico 87103

REGION 3

Regional Director
 U.S. Fish and Wildlife Service
 Federal Building, Fort Snelling
 Twin Cities, Minnesota 55111

REGION 4

Regional Director
 U.S. Fish and Wildlife Service
 Richard B. Russell Building
 75 Spring Street, S.W.
 Atlanta, Georgia 30303

REGION 5

Regional Director
 U.S. Fish and Wildlife Service
 One Gateway Center
 Newton Corner, Massachusetts 02158

REGION 6

Regional Director
 U.S. Fish and Wildlife Service
 P.O. Box 25486
 Denver Federal Center
 Denver, Colorado 80225

REGION 7

Regional Director
 U.S. Fish and Wildlife Service
 1011 E. Tudor Road
 Anchorage, Alaska 99503



DEPARTMENT OF THE INTERIOR U.S. FISH AND WILDLIFE SERVICE



As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering the wisest use of our land and water resources, protecting our fish and wildlife, preserving the environmental and cultural values of our national parks and historical places, and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to assure that their development is in the best interests of all our people. The Department also has a major responsibility for American Indian reservation communities and for people who live in island territories under U.S. administration.