

North Coast Hydrologic Region — Table of Contents

North Coast Hydrologic Region.....	NC-1
North Coast Hydrologic Region Summary	NC-1
Current State of the Region.....	NC-1
Setting	NC-1
Watersheds.....	NC-2
Klamath River Subbasin	NC-2
North Coastal Subbasin.....	NC-6
North Coast Rivers Basin.....	NC-8
Russian River Watershed.....	NC-12
Bodega Watershed	NC-12
Groundwater Aquifers.....	NC-12
Aquifer Description	NC-13
Well Infrastructure and Distribution	NC-16
California Statewide Groundwater Elevation Monitoring (CASGEM) Basin	
Prioritization	NC-18
North Coast Hydrologic Region Groundwater Monitoring Efforts	NC-18
Ecosystems.....	NC-20
Flood	NC-21
Recent Tsunamis on the California Coast.....	NC-22
Climate.....	NC-22
Demographics	NC-23
Population	NC-24
Tribal Communities	NC-25
Disadvantaged Communities	NC-27
Land Use Patterns	NC-27
Diversity of Agriculture in North Coast Region.....	NC-29
Regional Resource Management Conditions.....	NC-30
Water in the Environment.....	NC-30
Instream Fisheries Requirements.....	NC-30
Water Supplies	NC-30
Surface Water.....	NC-31
Groundwater	NC-32
Reclaimed Water.....	NC-34
Geysers Recharge Project	NC-34
Imported / Exported Water	NC-35
Water Uses	NC-35
Drinking Water	NC-35
Agricultural Water Use.....	NC-36
Water Conservation Act of 2009 (SB x7-7) Implementation Status.....	NC-37
Water Balance Summaries	NC-38
Project Operations.....	NC-39
Potter Valley Project.....	NC-39
Coyote Valley Dam and Lake Mendocino.....	NC-40
Warm Springs Dam and Lake Sonoma.....	NC-40
Petaluma Aqueduct	NC-40
North Marin Aqueduct.....	NC-41
R.W. Matthews Dam, Ruth Lake and Mad River.....	NC-41

Iron Gate Dam and Klamath River	NC-42
Trinity Dam and Exports from Trinity River to Central Valley	NC-43
USBR Flow Releases to the Trinity and Sacramento Rivers	NC-44
Gravel Mining	NC-45
Statewide Instream Mining (Suction Dredge Mining)	NC-45
Irrigated Lands Program	NC-46
Tribal Water Rights	NC-46
Tribal Water Rights on the Klamath and Trinity Rivers	NC-47
Water Quality	NC-48
Surface Water Quality	NC-48
Groundwater Quality	NC-50
Drinking Water Quality	NC-50
Groundwater Conditions and Issues	NC-51
Groundwater Occurrence and Movement	NC-51
Depth to Groundwater	NC-51
Change in Groundwater Storage	NC-55
Flood Management	NC-55
Flood Hazard Exposure	NC-56
Sea Level Rise	NC-56
Levee and Channel System	NC-57
Levee Performance and Risk Studies	NC-57
Recent Tsunamis on the California Coast	NC-58
Redwood Coast Tsunami Work Group	NC-58
Water Governance	NC-58
AB 2409 (Nestande, 2010)	NC-59
California Water Code Sec 1259.4 AB 2121	NC-59
Fish and Game Code Section 5653	NC-59
Water Code Division 5, Sections 8,000 - 9,651	NC-59
AB 70 (2007) Flood Liability	NC-59
AB 162 (2007) General Plans	NC-59
Fish and Game Code Section 1602 Streambed Alteration Permits	NC-60
Potter Valley Project FERC License	NC-60
Hydropower, a Renewable Energy	NC-61
State Funding Received	NC-61
Flood Governance	NC-61
Groundwater Governance	NC-61
Surface Water Ambient Monitoring Program (SWAMP)	NC-66
Current Relationships with Other Regions and States	NC-66
Klamath Basin	NC-66
Trinity River	NC-67
Potter Valley Project	NC-68
Sonoma-Petaluma Aqueduct	NC-68
Regional Water Planning and Management	NC-68
Integrated Regional Water Management Coordination and Planning	NC-69
Accomplishments	NC-69
Recent Initiatives or Actions to Improve Water Quality	NC-69
Ecosystem Restoration	NC-71
Sage Steppe Ecosystem Restoration	NC-72
Klamath Basin National Wildlife Refuge Complex	NC-72
River Restoration	NC-73
Klamath River	NC-73

Shasta and Scott Rivers.....	NC-73
Salt River	NC-73
Big River.....	NC-74
Salmon Creek.....	NC-74
Russian River	NC-74
Laguna de Santa Rosa.....	NC-75
Mattole River	NC-75
Trinity River Restoration Program	NC-75
Shasta River	NC-75
Challenges.....	NC-76
Flood Challenges	NC-78
Conjunctive Management and Groundwater Storage	NC-78
Conjunctive Management Inventory Results.....	NC-79
Drought Planning	NC-79
Klamath Basin Restoration Agreement Drought Plan 2011	NC-79
Looking to the Future.....	NC-79
Future Conditions.....	NC-79
Future Scenarios.....	NC-79
Water Conservation	NC-80
North Coast Growth Scenarios	NC-80
North Coast 2050 Water Demands	NC-81
Integrated Water Management Plan Summaries.....	NC-82
Resource Management Strategies	NC-83
Regional Resource Management Strategies.....	NC-84
Climate Change.....	NC-85
Observations	NC-85
Projections and Impacts	NC-86
Adaptation.....	NC-88
Mitigation.....	NC-89
References.....	NC-92
References Cited.....	NC-92
Personal Communications.....	NC-110

Tables

PLACEHOLDER Table NC-1 Alluvial Groundwater Basins and Subbasins within the North Coast Hydrologic Region.....	NC-13
PLACEHOLDER Table NC-2 Number of Well Logs by County and Use for the North Coast Hydrologic Region (1977-2010).....	NC-16
PLACEHOLDER Table NC-3 CASGEM Groundwater Basin Prioritization for the North Coast Hydrologic Region.....	NC-18
PLACEHOLDER Table NC-4 Groundwater Level Monitoring Wells by Monitoring Entity in the North Coast Hydrologic Region	NC-19
PLACEHOLDER Table NC-5 Sources of Groundwater Quality Information.....	NC-20
PLACEHOLDER Table NC-6 Federally Recognized Tribes in North Coast Hydrologic Region..	NC-25
PLACEHOLDER Table NC-7 California Native American Tribes (Non-Recognized) in North Coast Hydrologic Region	NC-25
PLACEHOLDER Table NC-8 North Coast Hydrologic Region Average Annual Groundwater Supply by Planning Area and by Type of Use (2005-2010)	NC-33
PLACEHOLDER Table NC-9 North Coast Hydrologic Region Average Annual Groundwater Supply by County and by Type of Use (2005-2010)	NC-33

[PLACEHOLDER Table NC-10](#) Summary of Community Water Systems within the North Coast Hydrologic Region..... NC-36

[PLACEHOLDER Table NC-11](#) North Coast Hydrologic Water Balance Summary, 2001-2010 (Thousand af)..... NC-38

[PLACEHOLDER Table NC-12](#) Trinity River ROD Water Year Types NC-45

[PLACEHOLDER Table NC-13](#) Trinity River Release and Diversion Summary Since 2000..... NC-45

[PLACEHOLDER Table NC-14](#) Summary of Community Drinking Water Systems in the North Coast Hydrologic Region that Rely on One or More Contaminated Groundwater Well that Exceeds a Primary Drinking Water Standard NC-51

[PLACEHOLDER Table NC-15](#) Summary of Contaminants Affecting Community Drinking Water Systems in the North Coast Hydrologic Region NC-51

[PLACEHOLDER Table NC-16](#) North Coast Hydrologic Region Water Management Agencies .. NC-58

[PLACEHOLDER Table NC-17](#) State Funding Received..... NC-61

[PLACEHOLDER Table NC-18](#) Groundwater Management Plans in the North Coast Hydrologic Region..... NC-62

[PLACEHOLDER Table NC-19](#) Assessment for SB 1938 GWMP Required Components, SB 1938 GWMP Voluntary Components, and Bulletin 118-03 Recommended Components..... NC-63

[PLACEHOLDER Table NC-20](#) Factors Contributing to Successful Groundwater Management Plan Implementation in the North Coast Hydrologic Region NC-64

[PLACEHOLDER Table NC-21](#) Factors Limiting Successful Groundwater Management Plan Implementation in the North Coast Hydrologic Region NC-64

[PLACEHOLDER Table NC-22](#) Groundwater Ordinances that Apply to Counties in the North Coast Hydrologic Region..... NC-64

[PLACEHOLDER Table NC-23](#) Groundwater Adjudications in the North Coast Hydrologic Region NC-65

[PLACEHOLDER Table NC-24](#) Conceptual Growth Scenarios..... NC-80

[PLACEHOLDER Table NC-25](#) Growth Scenarios (Urban) – North Coast..... NC-81

[PLACEHOLDER Table NC-26](#) Growth Scenarios (Agricultural) – North Coast..... NC-81

[PLACEHOLDER Table NC-27](#) Resource Management Strategies addressed in IRWMPs in the North Coast Hydrologic Region..... NC-84

Figures

[PLACEHOLDER Figure NC-1](#) North Coast Hydrologic Region NC-1

[PLACEHOLDER Figure NC-2](#) North Coast Watersheds NC-2

[PLACEHOLDER Figure NC-3](#) Alluvial Groundwater Basins and Subbasins within the North Coast Hydrologic Region..... NC-13

[PLACEHOLDER Figure NC-4](#) Number of Well Logs by County and Use for the North Coast Hydrologic Region (1977-2010)..... NC-17

[PLACEHOLDER Figure NC-5](#) Percentage of Well Logs by Use for the North Coast Hydrologic Region (1977-2010)..... NC-17

[PLACEHOLDER Figure NC-6](#) Number of Well Logs Filed per Year by Use for the North Coast Hydrologic Region (1977-2010)..... NC-17

[PLACEHOLDER Figure NC-7](#) CASGEM Groundwater Basin Prioritization for the North Coast Hydrologic Region..... NC-18

[PLACEHOLDER Figure NC-8](#) Monitoring Well Location by Agency, Monitoring Cooperator, and CASGEM Monitoring Entity in the North Coast Hydrologic Region..... NC-19

[PLACEHOLDER Figure NC-9](#) Percentage of Monitoring Wells by Use in the North Coast Hydrologic Region..... NC-19

[PLACEHOLDER Figure NC-10](#) Total Population 1960-2010 (2050), North Coast Hydrologic Region [figure to come]..... NC-25

PLACEHOLDER Figure NC-11 Population Growth Trends, North Coast Hydrologic Region
 [figure to come]..... NC-25

PLACEHOLDER Figure NC-12 Disadvantaged Communities in the North Coast Hydrologic
 Region..... NC-27

PLACEHOLDER Figure NC-13 North Coast Regional Inflows and Outflows in 2010. NC-31

PLACEHOLDER Figure NC-14 Contribution of Groundwater to the North Coast Hydrologic Region
 Water Supply by Planning Area (2005-2010)..... NC-33

PLACEHOLDER Figure NC-15 North Coast Hydrologic Region Annual Groundwater Water Supply
 Trend (2002-2010)..... NC-34

PLACEHOLDER Figure NC-16 North Coast Hydrologic Region Annual Groundwater Supply Trend
 by Type of Use (2002-2010)..... NC-34

PLACEHOLDER Figure NC-17 North Coast Water Balance by Water Year, 2001–2010 NC-38

PLACEHOLDER Figure NC-18 Groundwater Level Trends in Selected Wells in the North Coast
 Hydrologic Region..... NC-55

PLACEHOLDER Figure NC-19 Flood Exposure to the 100-Year Floodplain, North Coast Hydrologic
 Region..... NC-56

PLACEHOLDER Figure NC-20 Flood Exposure to the 500-Year Floodplain, North Coast Hydrologic
 Region..... NC-56

PLACEHOLDER Figure NC-21 Location of Groundwater Management Plans in the North Coast
 Hydrologic Region..... NC-62

PLACEHOLDER Figure NC-22 Groundwater Adjudications in the North Coast
 Hydrologic Region..... NC-65

PLACEHOLDER Figure NC-23 Integrated Regional Management Planning areas in the North Coast
 Hydrologic Region..... NC-69

PLACEHOLDER Figure NC-24 Change in North Coast Agricultural and Urban Water Demands for
 117 Scenarios from 2006-2050 (thousand af /yr) NC-81

PLACEHOLDER Figure NC-25 Energy Intensity of Raw Water Extraction and Conveyance in the
 North Coast Hydrologic Region NC-90

Boxes

PLACEHOLDER Box NC-1 California Statewide Groundwater Elevation Monitoring (CASGEM) Basin
 Prioritization Data Considerations NC-18

PLACEHOLDER Box NC-2 Near Coastal Issues..... NC-22

PLACEHOLDER Box NC-3 Tribal Water Rights on the Klamath and Trinity Rivers NC-47

PLACEHOLDER Box NC-4 Other Groundwater Management Planning Efforts in the North Coast
 Hydrologic Region..... NC-66

PLACEHOLDER Box NC-5 Statewide Conjunctive Management Inventory Effort in California.... NC-79

Acronyms and Abbreviations Used in This Report

AB	Assembly Bill
ABAG	Bay Area Association of Bay Area Governments
ACWA	Association of California Water Agencies
BLM	Bureau of Land Management
BMO	Basin Management Objective
BO	Biological Opinion
CASGEM	California Statewide Groundwater Elevation Monitoring
CDPH	California Department of Public Health
CEQA	California Environmental Quality Act
cfs	Cubic Feet per Second
CIMIS	California Irrigation Management Information System
CVP	Central Valley Project
CVPIA	Central Valley Project Improvement Act
CWA	Clean Water Act
DAC	Disadvantaged Community
DFW	California Department of Fish and Wildlife
DPR	Department of Pesticide Regulation
DWR	California Department of Water Resources
EIS	Environmental Impact Statement
EL	Energy Intensity
EP	Effective Precipitation
FERC	Federal Energy Regulatory Commission
FRG	Fisheries Review Group
GAMA	Groundwater Ambient Monitoring and Assessment
GHG	Greenhouse Gas
GMWP	Groundwater Management Plan
gpm	Gallons per Minute
HBMWD	Humboldt Bay Municipal Water District
HCP	Habitat Conservation Plan
HIP	High Population Scenario
HVT	Hoopa Valley Tribe
KBRA	Klamath Basin Restoration Agreement
KHSA	Klamath Hydroelectric Settlement Agreement
LLNL	Lawrence Livermore National Laboratory
LOP	Low Population Growth Scenario
maf	Million Acre-Feet
DOF	California Department of Finance
mgd	Million Gallons per Day
MHI	Median Household Income
MSL	Mean Sea Level
MWh	Megawatt-Hour

NCIRWMP	North Coast Integrated Regional Water Management Plan
NCRWMG	North Coast Regional Water Management Group
NMFS	National Marine Fisheries Service
NPDES	National Pollutant Discharge Elimination System
NPS	Non-Point-Sources
NWR	National Wildlife Refuge
PA	Planning Area
PCP	Pentachlorophenol
POA	Plan of Action for Phase II
RCD	Resource Conservation District
RCTWG	Redwood Coast Tsunami Work Group
ROD	Record of Decision
RPA	Reasonable and Prudent Alternative
RPS	Renewables Portfolio Standard
RVCWD	Redwood Valley County Water District
RVIT	Round Valley Indian Tribes
RWMG	Regional Water Management Groups
RWQCB	Regional Water Quality Control Board
SB	Senate Bill
SCWA	Sonoma County Water Agency
SWAMP	Surface Water Ambient Monitoring Program
SWN	State Well Number System
SWRCB	State Water Resources Control Board
taf	Thousand Acre-Feet
THP	Timber Harvest Plan
TMDL	Total Maximum Daily Loads
TPZ	Timber Production Zone
TRD	Trinity River Diversion
TRFES	Trinity River Flow Evaluation Study
UKL	Upper Klamath Lake
USACE	U.S. Army Corps of Engineers
USBR	U.S. Bureau of Reclamation
USFS	U.S. Forest Service
USFWS	U.S. Fish and Wildlife Service
USGS	U.S. Geological Survey
VBF	Variable Base Flow
WAMP	Watershed Adaptive Management Plan
WDR	Waste Discharge Requirements
WRCC	Western Regional Climate Center
YT	Yurok Tribe

1 North Coast Hydrologic Region

2 North Coast Hydrologic Region Summary

3 The North Coast Hydrologic Region is a unique setting with an extreme diversity of land use and climate.
 4 Land use includes fisheries and larger urban areas near the coast to high mountain deserts with large cattle
 5 operations and low populations. This chapter begins with an overview of the region, describing the setting
 6 and conditions within the region. Topics include information on the watersheds and sub-watersheds in the
 7 region with emphasis on developed resources. The chapter continues with a review of activities
 8 concerning resource administration and laws affecting resource management. This chapter concludes with
 9 a discussion of suggested resource management strategies to help assist local water managers in planning
 10 for future water needs.

11 Current State of the Region

12 Setting

13 The North Coast Hydrologic Region encompasses coastal areas, redwood forests, inland mountain
 14 valleys, and the semi-desert-like Modoc Plateau. Most of the region is mountainous and rugged. The
 15 dominant topographic features in the region are the California Coast Ranges, the Klamath Mountains, and
 16 Modoc Plateau. The mountain crests, which form the eastern boundary of the region, are about 6,000 feet
 17 elevation with a few peaks higher than 8,000 feet. Much of the region is mountainous and rugged; only
 18 13 percent of the land is classified as valley or mesa, and more than half of that is in the higher-elevation
 19 northeastern part of the region in the upper Klamath River Basin.

20 **PLACEHOLDER Figure NC-1 North Coast Hydrologic Region**

21 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 22 the end of the report.]

23 The North Coast Hydrologic Region is one of 10 hydrologic regions defined by the California
 24 Department of Water Resources. It shares boundaries with the North Coast Region as defined in Section
 25 13200(a) of Porter-Cologne Water Quality Control Act, which divides the state into nine regional board
 26 boundaries:

27 “North Coast region, which comprises all basins including Lower Klamath Lake
 28 and Lost River Basins draining into the Pacific Ocean from the California-
 29 Oregon state line southerly to the southerly boundary of the watershed of the
 30 Estero de San Antonio and Stemple Creek in Marin and Sonoma Counties.”

31 The North Coast Region is divided into two natural drainage basins, the Klamath River Subbasin and the
 32 North Coastal Subbasin. The North Coast Hydrologic Region covers all of Del Norte, Humboldt, Trinity,
 33 and Mendocino counties, major portions of Modoc, Siskiyou, and Sonoma counties, and small portions of
 34 Glenn, Lake, and Marin counties.

1 The North Coast Region encompasses a total area of approximately 19,390 square miles, including
2 340 miles of scenic coastline and remote wilderness areas, as well as urbanized and agricultural areas.
3 The region is characterized by distinct temperature zones. Along the coast, the climate is moderate and
4 foggy, and the temperature variation is not great. For example, at Eureka, the seasonal variation in
5 temperature has not exceeded 63 °F for the period of record. Inland, however, seasonal temperature may
6 range into 100 °F or higher.

7 Precipitation over the North Coast region is greater than over any other part of California, and damaging
8 floods are a fairly frequent hazard. Particularly devastating floods occurred in the North Coast area in
9 December of 1955, in December of 1964, in February of 1986, and over New Year's of both 1997 and
10 2006.

11 **Watersheds**

12 The North Coast region includes many watersheds and basins within its boundaries. Each of the main
13 region's boundaries as defined by the State Water Resources Control Board (SWRCB), California
14 Department of Water Resources (DWR), and the North Coast Integrated Regional Water Management
15 Plan (NCIRWMP) coincide with each other. According to the NCIRWMP, several subbasins exist
16 including the Klamath River Subbasin, North Coastal Subbasin, North Coast River Basin, Russian River
17 and Bodega watersheds, each containing many subareas within their boundaries. (See Figure NC-2)

18 **PLACEHOLDER Figure NC-2 North Coast Watersheds**

19 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
20 the end of the report.]

21 *Klamath River Subbasin*

22 The Klamath River Subbasin contains Klamath River and all of its tributaries (including Trinity River),
23 the Smith River and its tributaries, Applegate, Illinois, and Winchuck rivers and includes the closed Lost
24 River and Butte Valley hydrologic drainage areas. The western portion of the subbasin is within the
25 Klamath Mountains and Coast Ranges provinces, characterized by steep, rugged peaks ranging to
26 elevations of 6,000 to 8,000 feet with relatively little valley area. The mountain soils are shallow and
27 often unstable. Precipitation ranges from 60 to 125 inches per year in the western portion. The 45-mile
28 coastline is dominated by a narrow coastal plain where heavy fog is common. The eastern portion of the
29 basin receives low to moderate rainfall and includes predominantly high, broad valleys such as the Butte,
30 Shasta, and Scott valleys. The Lost River and Butte Valley hydrologic areas are located in the Modoc-
31 Oregon Lava Plateau. This area is characterized by broad valleys ranging from 4,000 to 6,000 feet in
32 elevation. Typical annual precipitation is 15 to 25 inches. The Shasta Valley hydrologic area lies
33 principally within the Cascade Range. The valley floor elevation is about 2,500 to 3,000 feet, and
34 surrounding mountains range up to 14,162 feet (Mount Shasta). Annual precipitation ranges from below
35 15 inches in the valley to over 60 inches in the mountains. The Scott River hydrologic area is in the
36 Klamath Mountains to the west of Shasta Valley. The valley floor elevation is also about 2,500 to
37 3,000 feet, with surrounding mountain ranges up to approximately 8,500 feet. Annual precipitation ranges
38 from below 20 inches in the valley to over 70 inches in the western mountains.

Klamath River Watershed (Oregon and California)

The Klamath is the second largest river in California with an extensive watershed of almost 16,000 square miles including portions of California and Oregon. The Klamath River begins north of Klamath Falls, Oregon, and meets the Pacific Ocean near the town of Klamath, California. For the sake of this discussion, the Klamath Basin is divided into three areas; the upper, middle, and lower Klamath subbasins. Hydrologic subbasins within the Klamath Basin include Butte Valley, Lost River, Salmon River, Scott River, Shasta River, and Trinity River.

The Upper Klamath subbasin encompasses the area upstream of the Iron Gate Dam. Only a small part of this area is located in California. The primary subwatershed in California is the Lost River watershed, which covers approximately 1,689 square miles and includes Clear Lake Reservoir in Modoc County. The area around Clear Lake is characterized by high desert streams and is sparsely populated. Land uses in the California portion of the Klamath Basin are primarily cropped agriculture, grazing, and lands administered for the Klamath Basin National Wildlife Refuge. The basin is subject to many complex jurisdictional issues associated with water delivery and utilization of water infrastructure facilities including issues related to irrigation, hydropower, endangered species, tribal rights, and lake level management demands.

The Middle Klamath subbasin is contained wholly within California extending from Iron Gate Dam to the confluence of Scott River about 10 miles upstream from Seiad Valley, excluding the Shasta and Scott rivers. However, the Middle Klamath subbasin is influenced by adjacent Klamath River subbasins (the Upper Klamath, Lower Klamath, and Trinity River drainages) and by the direct effects of tributary rivers (the Shasta and Scott rivers) which flow into the Klamath within the area of the Middle Klamath subbasin. The lower, more western portion has a coastal influenced climate and is dominated by U.S. Forest Service lands while the upper, more eastern portion has a drier climate with mixed federal and private ownership.

The Lower Klamath subbasin begins below the confluence of the Klamath Scott River extending to the Klamath River delta at the Pacific Ocean. Trinity River watershed, although tributary to the Klamath in this subbasin, is considered its own watershed and is not in the Lower Klamath subbasin. The major industry in the watershed is silviculture and some limited mining. Salmon fishing has occurred in the basin since Native American occupation, although in 2006 the commercial fishery has been restricted due to record low populations.

Shasta River Watershed

The Shasta River watershed includes an 800 square-mile area of Siskiyou County. Mount Shasta to the south dominates the landscape, towering higher than 14,000 feet. However, melting snow from Mount Shasta does not contribute significantly to surface flows in the upper Shasta River because runoff sinks into the porous volcanic soils and reappears as springs on the Shasta Valley floor. The headwaters of the Shasta River are near Mount Eddy in the southwest portion of the basin. Mount Eddy is the tallest mountain in Trinity County and the Klamath Mountains at 9,025 feet. The upper river above Dwinell Reservoir is swift and falls in elevation rapidly. The river below Dwinell Reservoir is much slower and meanders along the Shasta Valley floor. Springs in this reach add to flows and provide much needed cool water for juvenile salmon and steelhead in summer. The Klamath Mountains to the west, strip most of the moisture from ocean air currents as they move eastward. The Shasta Valley itself receives only 11 to 17 inches of rain annually. Because so little rain falls in the Shasta Valley during the growing season,

1 ranchers rely heavily on streamflows and groundwater to irrigate crops and to water livestock. The
2 economy of Shasta Valley, like that of Siskiyou County generally, relies on ranching, farming, tourism,
3 and timber harvesting. Sport-fishing opportunities still draw visitors to Siskiyou County because of
4 numerous mountain lakes and productive streams. Yreka and Weed contain the largest populations in this
5 subwatershed.

6 **Scott River Watershed**

7 The Scott River watershed is a large area with substantial variation in geology and climate. The watershed
8 drains approximately 520,600 acres of land. Major tributaries to the 58-mile-long Scott River in Scott
9 Valley include Shackleford-Mill, Kidder, Etna, French, and Moffett creeks, including the South and East
10 Forks of the Scott River. Native vegetation consists of riparian vegetation along the streams, mixed-
11 conifer forest on the western mountain slopes, with scattered meadows and brush. The eastern mountains
12 are covered by extensive areas of brush, oak, western juniper, and both annual and perennial grasses. The
13 confluence of Scott and Klamath rivers is located approximately 10 miles upstream (along Klamath
14 River) from Seiad Valley. The Scott River drainage is bordered to the west and south by 7,000- to
15 8,000-foot elevation mountain ranges, including the Marble, Salmon, Trinity Alps and Scott mountains.
16 These ranges exert a strong orographic effect on incoming storms, which allows the higher elevation
17 mountains, along the west and south side of the Scott drainage, to receive 60 to 80 inches of precipitation
18 annually. In contrast, the rain-shadow effect that the westside mountains create reduces the amount of
19 annual precipitation to 12 to 15 inches on the east side of the watershed. Fort Jones, located at the
20 northern end of Scott Valley, averages 21 inches of precipitation, although rainfall has ranged from
21 10 inches in 1949 to 35 inches in 1970 showing the variability in the climate. Most of the precipitation in
22 the Scott River watershed falls on the west side, with snow prevailing during the winter above the
23 5,500 foot level. Snowfall is an important component of the water supply for the region.

24 **Salmon River Watershed**

25 The Salmon River flows from the Trinity Alps, Marble, Russian, and Salmon mountains joining the
26 Klamath River at Somes Bar, California, and is the second largest tributary to the Klamath next to Trinity
27 River. The watershed is almost entirely public land (Klamath National Forest) containing rugged
28 topography that is deeply incised by the river and its tributaries. Nearly the entire watershed is forested.
29 There are no dams, diversions, urban areas, or major industry in the watershed so the water is very high
30 quality. In addition, there are no dams between the Salmon River and the ocean, making it completely
31 accessible to anadromous fish. The cool, clean waters of the Salmon River are critical to the overall health
32 of the Klamath River fishery. The Salmon River provides genetic stock and quality habitat for fish and
33 other aquatic life making this watershed of great importance to the recovery of larger Klamath River
34 watershed. Elevations in the watershed range from 456 feet at its mouth to 8,560 feet at Caribou
35 Mountain in the Trinity Alps. The Salmon River remains culturally significant to the Shasta and Karuk
36 people, some of whom continue to reside on the river. Approximately 67 percent of the watershed is in
37 the Karuk Tribe's Ancestral Territory. Mean annual precipitation in the Salmon River watershed ranges
38 from about 35 inches in the South Fork Salmon River Canyon to about 85 inches in the headwaters of
39 North Fork/Little North Fork and Wooley Creek. The amount of precipitation generally decreases in an
40 easterly direction, and increases with elevation due to orographic effects. Seasonal precipitation patterns
41 include considerable snow, particularly at higher elevations. Approximately 90 percent of the
42 precipitation occurs from October to May. The remainder occurs during summer thunderstorms. Winter
43 precipitation occurs mainly as snow above 4,000 feet, with rain below 4,000 feet elevation. Fluctuation of

1 the snow level occasionally results in rain falling on snow, causing rapid snowmelt. Intense, localized
2 summer showers occur frequently and have been associated with soil erosion and debris torrents.

3 **Trinity River Watershed**

4 The Trinity River Basin drains an area of approximately 2,900 square miles of mountainous terrain. The
5 Trinity River is the largest tributary to the Klamath River. From its headwaters in the Klamath Mountains,
6 the river flows 172 miles south and west through Trinity County, then north through Humboldt County
7 and the Hoopa Valley and Yurok Indian reservations to its confluence with the Klamath River at
8 Weitchpec. Much of the watershed is prone to seismically induced landslides, especially during winter
9 months when soils are saturated. Additionally, inner valley gorges are considered highly unstable.

10 Groundwater resources are relatively plentiful throughout the watershed, but are not well defined. Annual
11 precipitation averages 57 inches a year with a low of 37 inches in Weaverville and Hayfork and a higher
12 rainfall of 75 inches in Trinity Center and 85 inches in the Hoopa Mountains. There are occasional
13 summer thunderstorms that produce extensive runoff and may start wildfires.

14 The Trinity River watershed is primarily rural with human populations centered near Trinity Center,
15 Weaverville, Lewiston, Hayfork, and Hyampom. Timber harvest has traditionally been a large factor in
16 the economy on both federal and private land. The U.S. Forest Service (USFS) and the Bureau of Land
17 Management (BLM) manage approximately 80 percent of the land in the Trinity watershed; of the
18 remaining 20 percent, about half are industrial timberlands.

19 In the early 1950s, two major water-development features were installed above river-mile 112 and the
20 community of Lewiston on the Trinity River. In 1955, the Trinity River Division Act authorized the
21 Trinity River Diversion (TRD). The TRD consists of Lewiston Dam and its reservoir and related facilities
22 and Trinity Dam and its reservoir (known as Trinity Lake). The TRD project diverts a majority of the
23 upper-basin's water yield at Lewiston for power generation and to support the U.S. Bureau of
24 Reclamation's (USBR) Central Valley Project (CVP). The hydrologic changes produced by the TRD
25 project have altered stream-channel conditions and instream habitat for many miles below Lewiston.
26 Trinity River downstream of the TRD provides habitat not only for anadromous salmonids and other
27 native species, but also the non-native brown trout. Operations of the TRD began in 1964 and were
28 integrated with operations of Shasta Dam.

29 Water quality in the Trinity River Basin ranges from the high quality, pristine waters that emerge from the
30 Trinity Alps wilderness to various degrees of impairment in the main stem and southern tributaries, which
31 are caused in part by human activity. Timber harvest, road construction, and associated activities are
32 recognized as sources of sedimentation and high summer water temperatures. Mining for gold, both
33 currently and historically, is also a source of impairment. Recreational instream suction dredging (mining)
34 causes sedimentation, especially in the main stem and canyon areas, and legacy effects from historic gold
35 mining include acid mine drainage and mercury pollution. Please see section on Regional Resource
36 Management Conditions for more information on instream mining (suction dredge mining).

37 **Smith River Watershed (Oregon and California)**

38 The Smith River is formed by the confluence of its Middle and North forks in Del Norte County, in the
39 extreme northwest corner of California, near the community of Gasquet. The Middle Fork originates in
40 Del Norte County, approximately 60 miles northeast of Crescent City, and flows west. The North Fork
41 Smith River originates in Oregon on the northeast slope of Chetco Peak in the Siskiyou Mountains. The

1 South Fork Smith River enters the Smith River near the community of Hiouchi, California. The South
2 Fork rises on the eastern edge of the Smith River National Recreation Area, approximately 30 miles east-
3 northeast of Crescent City, flowing southwest and then northwest. From the confluence with the South
4 Fork, the Smith River flows generally northwest, entering the Pacific Ocean near the community of Smith
5 River, approximately 10 miles north of Crescent City.

6 The Smith River estuary is located in Del Norte County near the community of Smith River. The
7 watershed is about 614 square miles. The Smith is the longest wild and scenic river in the United States;
8 as such, there are no impoundments. The Smith River system is the second largest free-flowing river in
9 California next to the South Fork Trinity River. It is considered one of the best fishing regions in the
10 United States with steelhead, Chinook, and other game fishes present. The region receives from 80 to
11 120 inches of rainfall annually.

12 In the Smith River Basin, no significant surface water development has occurred. Domestic, agricultural,
13 and industrial water needs are supplied through surface water diversions and groundwater pumping.
14 Further major developments on the Smith River and any of its tributaries are forbidden by the 1972
15 California Wild and Scenic Rivers Act. However, minor surface water supply projects for high value
16 crops in the Smith River area are possible. Because of both its geology and its limited development, the
17 Smith River is one of the healthiest river systems in California.

18 Federal land management dominates the Smith River Basin. Six Rivers National Forest manages the
19 Smith River Recreation Area, which includes 305,000 acres, or 476 square miles of the watershed.
20 Siskiyou National Forest manages 91 square miles of the basin within Oregon. Redwood National and
21 California Department of Parks and Recreation (California State Parks) have jurisdiction in 25 square
22 miles of the watershed. The total land managed by government agencies is about 83 percent of the
23 watershed, which leaves 126 square miles in private ownership, predominantly in the lower river basin.

24 *North Coastal Subbasin*

25 The North Coastal Subbasin consists of rugged, forested coastal mountains, including six major river
26 systems: the Eel, Russian, Mad, Navarro, Gualala, and Noyo rivers. In addition, among others, the North
27 Coastal Subbasin includes the Mattole and Garcia rivers and Redwood and Stemple creeks. Soils are
28 generally unstable and erodible, and rainfall is high. The area along the eastern boundary of the basin is
29 mostly National Forest land administered by the USFS. Major population areas are centered on Humboldt
30 Bay in the northern portion of the basin and around Santa Rosa in the southern portion. The Santa Rosa
31 area is on the northern fringe of the greater San Francisco Bay urban area and has experienced rapid
32 population growth in the period following the Second World War. The economy of the remainder of the
33 basin has developed more slowly than other areas in California.

34 **Humboldt Bay Watershed**

35 The Humboldt Bay watershed encompasses water bodies that drain to the Pacific Ocean from Humboldt
36 Bay north to Redwood Creek. The major river systems in the watershed are the Mad River and Redwood
37 Creek. Other water bodies within this watershed include Humboldt Bay and Mad River Slough, coastal
38 lagoons (Big, Stone, and Freshwater lagoons) and streams (Elk and Little rivers and Freshwater, Jacoby,
39 and Maple creeks). In the east, the terrain is elevated hill slope with coastal plain occurring in the west.
40 Precipitation ranges from 32 to 98 inches annually. The streams support production of anadromous
41 salmonids, including steelhead and cutthroat trout, coho and Chinook salmon.

The Mad River Watershed

The Mad River watershed has a long history of timber harvest on both USFS and private land. Gravel mining occurs in the lower portions of the watershed. Private landowners conduct grazing and limited agriculture in the flat areas around the bay. Humboldt Bay is an important commercial and recreational shellfish growing and harvesting area and provides the largest port between San Francisco and Coos Bay, Oregon. Urbanized areas include Trinidad, McKinleyville, Arcata, and Eureka; and rural residential areas are scattered throughout the watershed. The majority of the population lives in the Humboldt Bay area cities of Arcata and Eureka.

The Mad River is Clean Water Act (CWA) section 303(d) listed for sediment and temperature impacts. The primary issues for water quality are forestry related, with urbanization and associated industrial and public nonpoint sources. The drinking water for most of the Humboldt Bay area is supplied by Ranney Collectors in Mad River with other coastal streams providing drinking water for other communities. Mad River is continuously supplied with water via releases from Ruth Reservoir (with 48,030 acre-foot storage capacity), although these supplies are dependent on adequate precipitation and flows through the season. The Eureka waterfront was the site of several industrial operations that left the soil and groundwater contaminated with heavy metals, petroleum products, and pentachlorophenol's (PCPs). The waterfront is now undergoing redevelopment, and decontamination efforts continue.

Redwood Creek

Redwood Creek flows into the Pacific Ocean near the town of Orick located about 35 miles north of Eureka. Redwood Creek drains a 285 square-mile area and is about 67 miles long. The watershed is located entirely within Humboldt County.

Redwood Creek is a basin of mixed ownership and contains a rich blend of industrial and non-industrial timberlands, coastal and upland agricultural lands, State and federal national parks, other federal properties, and the unincorporated town of Orick. Redwood Creek supports three federally listed as threatened salmonids species as well as the non-listed coastal cutthroat trout and resident fish species. The watershed also provides domestic water supplies to rural communities and recreational opportunities.

Redwood Creek is a model watershed where government agencies, private landowners, nonprofit organizations and the local communities are cooperating to restore and protect water quality and the associated aquatic and riparian resources, providing economic opportunity to the Orick community. The watershed has a rich history of scientific studies that spans decades and well-established cooperation between groups with seemingly conflicting interests. The watershed is home to pioneering work in watershed restoration and erosion control.

The Redwood Creek watershed is a mixed ownership of private (56 percent) and public (44 percent) lands. More than 90 percent of the private lands are managed for timber production and ranching by eight private landowners. The upper two-thirds of the watershed contain vast expanses of timber and ranch lands managed primarily by seven landowners. Timberlands have been maintained in large unbroken tracts of lands, which have slowed rural residential development in upland areas. Located along the coast, the small town of Orick is the only municipality in the watershed and has a population of about 357 people (2010 U.S. Census). Orick is relatively isolated from other north coast communities and qualifies as a "disadvantaged community." (See "Demographics" section, subsection "Disadvantaged Communities.")

1 Redwood National Park and Prairie Creek Redwoods State Park are located in the lower part of the
2 Redwood Creek basin. This subbasin has been extensively researched and is considered a “reference
3 watershed” that displays nearly pristine conditions and is home to significant old growth stands of coast
4 redwood. The protection of streamside redwoods along Redwood Creek was a central issue for the
5 establishment and expansion of Redwood National Park and is linked to upstream watershed conditions.

6 Eel River

7 The Eel River and its tributaries are the third largest river system in California and the largest river system
8 draining to Humboldt County’s coast. The Eel River encompasses roughly 3,684 square miles. The main
9 tributaries to the Eel River are the Van Duzen River, the Bear River, and Yager, Larabee, Bull, and
10 Salmon creeks. Lake Pillsbury is located near the headwaters of the main stem Eel. The upper watershed
11 is mountainous, and soils are steep and highly erodible. In the west, the river meanders on a coastal plain
12 and is joined by the Salt River before entering the Pacific Ocean. Several dairies are located on the coastal
13 plain, as well as several small towns. Other communities in the watershed include Scotia, Garberville/
14 Redway, Laytonville, and Willits. In many of the alluvial valleys, surface water and groundwater are
15 closely connected, thus surface water withdrawals have a substantial effect on local groundwater supplies.
16 A Northwestern railroad line following along the Eel River has fallen into disrepair due to numerous
17 landslides and accidents. Currently, there are no plans to revive the railroad due to the high cost of
18 highway realignment and construction. The Eel River watershed is a well-known recreation destination
19 with numerous State and private campgrounds along its length; beneficial uses include both water contact
20 and noncontact uses such as swimming and boating. The river also supports a large recreational fishing
21 industry being the third largest producer of salmon and steelhead in the state. Due to the erodible soils,
22 steep terrain, and land use history, there is significant concern for the viability of this anadromous fishery
23 resource.

24 A longstanding transfer of water occurs downstream from Lake Pillsbury at Cape Horn Dam (Van
25 Arsdale Reservoir) moving water from the Eel River to the Russian River watershed (Potter Valley
26 Project). This out-of-basin transfer from the Eel River to the Russian River began in 1908 with the Eel
27 River Power and Irrigation Company. The purpose of this project was to supply the nearby town of Ukiah
28 with electricity and improve streamflows in the Russian River for municipal, industrial, and agricultural
29 uses. The areas served by this water have become dependent on this source, creating pressure to continue
30 the diversion in opposition to full restoration of the Eel River to its pre-diverted state. For additional
31 information on the Potter Valley Project, see "Project Operations" section in this report.

32 *North Coast Rivers Basin*

33 The North Coast River Basin also contains other major watersheds not listed above. These include the
34 Bear River, Mattole River, Ten Mile River, Noyo River, Big River, Albion River, Navarro River,
35 Greenwood, Elk and Alder creeks, Garcia River, Gualala River, and the Russian and Bodega watersheds.

36 **Bear River**

37 Bear River is a coastal stream located to the north of the Mattole River watershed draining approximately
38 53,287 acres to the Pacific Ocean. The connection between the Bear River and the Pacific Ocean is
39 periodically blocked by a temporary sand bar during summer low flow. The lagoon-type estuary is
40 approximately one-quarter mile in length. The two major land uses in the basin consist of agricultural
41 grazing and timber harvest. Humboldt Redwood Company (formerly Pacific Lumber) owns 16,537 acres
42 of land in the upper portion of the watershed, all of which is covered by its 1999 Habitat Conservation

1 Plan (HCP). The majority of remaining acreage in the watershed is in private ownership (36,839 acres),
2 and 161 acres are managed by California State Parks.

3 **Mattole River**

4 The headwaters of the Mattole River begin in Mendocino County, and it flows north 62 river miles,
5 through steep, forested lands in Humboldt County and into the ocean 10 miles south of Cape Mendocino.
6 Tributaries to the Mattole River include Mill, Squaw, Bear, Thompson, Honeydew, and Bridge creeks.
7 The watershed encompasses approximately 304 square miles and is subject to varying rainfall. Near the
8 coast, the river receives about 50 inches per year while near the headwaters, about 115 inches of rainfall
9 per year. The largest communities are Petrolia, Honeydew, and Whitethorn, but the 2,000-person
10 population is scattered throughout the watershed. Small landowners (those with less than 450 acres) own
11 about 43 percent of the watershed. The BLM owns about 12 percent, and commercial timber companies
12 own most of the remaining land. Silviculture and ranching are the predominant businesses. Water quality
13 problems are those associated with timber harvest, road building, forest conversion, and overgrazing. Fish
14 species known to inhabit the Mattole River include coho, Chinook, steelhead, rainbow trout, and brook
15 lamprey; other species include the southern torrent salamander and tailed frog.

16 **Ten Mile River**

17 The Ten Mile River watershed covers approximately 120 square miles. It is about 8 miles north of the
18 City of Fort Bragg and shares ridges with Pudding Creek and the North Fork of the Noyo River to the
19 south and Wages Creek and the South Fork of the Eel River to the north. Elevations range between sea
20 level and 3,205 feet. Near the coast, the terrain comprises an estuary and a broad river floodplain with
21 more rugged mountainous topography in the eastern portion of the watershed. Most of the basin, except
22 the northeast grasslands, coastal plain, and estuary, is characterized by narrow drainages bordered by
23 steep to moderately steep slopes. The watershed has abundant rainfall and cool temperatures during the
24 winter with dry, warm summers interspersed with breezes and coastal fog. Precipitation in the western
25 part of the watershed is about 70 inches per year while about 40 inches per year occurs in the eastern part
26 of the watershed.

27 The watershed is entirely privately owned. Hawthorne Timber Company, LLC, which is managed by
28 Campbell Timberland Management, LLC, owns about 85 percent of the watershed. Three small non-
29 industrial timber owners and a few residences make up the remainder of the ownership. The watershed
30 has a long history of timber harvest.

31 The coldwater fishery that supports coho, Chinook, and steelhead is the primary and most sensitive
32 beneficial use in the watershed. Protection of these species is considered to protect any of the other
33 beneficial uses identified in the watershed that could be impaired due to water quality.

34 **Noyo River**

35 The Noyo River watershed encompasses the 113 square-mile coastal drainage system immediately west
36 of the City of Willits, flowing into the Pacific Ocean at the City of Fort Bragg. The climate consists of
37 moderate temperatures — an annual average of 53 °F — and an average annual rainfall of 40 to
38 65 inches.

39 Silviculture is the primary land use within the watershed. Approximately 50 percent of the watershed is
40 owned by two commercial silviculture operations: the Mendocino Redwood Company and Hawthorne

1 Timber Company (managed by Campbell Timberland Management). The Jackson Demonstration State
 2 Forest (administered by the California Department of Forestry and Fire Protection — CALFIRE)
 3 encompasses about 19 percent of the watershed. Critical Coastal Areas in the vicinity of the watershed
 4 include Pudding Creek, Noyo River, and the Pygmy Forest Ecological Staircase. Minor land uses in the
 5 basin include ranching and recreation. The mouth of the Noyo River contains a marina and fish
 6 processing facilities in support of the local commercial fishing industry. The Noyo is the primary drinking
 7 water source for the City of Fort Bragg and also provides habitat for steelhead, coho, and Chinook. It is
 8 listed as impaired by sediment, due in part to timber harvest, grazing, and related human activities.

9 **Big River**

10 The Big River watershed drains about 181 square miles. The watershed drains from east to west, and
 11 shares ridges with the Noyo River watershed to the north, the Eel River watershed to the east, and the
 12 Little, Albion, and Navarro River watersheds to the south. The Big River estuary is located immediately
 13 south of the town of Mendocino. The climate is characterized by a pattern of low-intensity rainfall in the
 14 winter and cool, dry summers with coastal fog. Mean annual precipitation is approximately 40 inches near
 15 the western part of the watershed on the coast and about 51 inches at Willits to the east. The eastern part
 16 of this watershed receives more rainfall due to the orographic effect.

17 The predominant current and historical land use is silviculture with less area used for ranching. The
 18 largest community is the town of Mendocino. Together, the five largest property owners — four private
 19 timber companies and Jackson State Demonstration Forest — own 83 percent of the watershed.
 20 Thirty-one property owners own another 14 percent of the land (parcels from 160 acres to 3,760 acres),
 21 and private residences make up the rest of the land use.

22 **Albion River**

23 The Albion River watershed drains approximately 43 square miles. It drains primarily from east to west,
 24 and shares ridges with the Big River watershed to the north and northeast and the Navarro River
 25 watershed to the south and southeast. The Albion River estuary is located near the town of Albion, about
 26 16 miles south of the City of Fort Bragg. Elevations range from sea level to 1,566 feet, and the watershed
 27 is dominated by relatively flat marine terraces that extend several miles inland and are incised by gorges
 28 carved by the major river channels and streams. The climate in the watershed is characterized by a pattern
 29 of low intensity rainfall in the winter and cool, dry summers with coastal fog. Mean annual precipitation
 30 is about 40 inches near the western margin of the watershed and about 51 inches to the east at Willits. The
 31 main tributaries of the Albion River include Railroad Gulch, Pleasant Valley Creek, Duck Pond Gulch,
 32 South Fork Albion River, Tom Bell Creek, North Fork Albion River, and Marsh Creek.

33 More than half of the watershed (54 percent) is owned by Mendocino Redwood Company. Smaller
 34 industrial timberland ownerships, some ranches, and numerous smaller parcels that are mostly residences
 35 comprise the other half. The predominant historical and current land use is silviculture, with some
 36 agricultural and recreational uses. The Albion River estuary, which remains open to the sea year round, is
 37 used as a commercial and sport-fishing harbor for small boats. The river and estuary have historically
 38 served as habitat for coho, Chinook, and steelhead. Beneficial uses associated with the coldwater fishery
 39 are the most sensitive of the beneficial uses in the watershed; protection of these beneficial uses is thought
 40 to protect other beneficial uses harmed by excessive sediment.

1 **Navarro River**

2 The Navarro River watershed encompasses approximately 315 square miles. The Navarro River flows
3 through the Coast Ranges, Anderson Valley, and into the Pacific Ocean. It is the largest coastal basin in
4 Mendocino County. Rainfall averages about 40 inches per year at Philo and mostly occurs between
5 December and March.

6 Land uses in the watershed include silviculture (70 percent), rangeland (25 percent), and agriculture
7 (5 percent) with a small percentage devoted to rural residential development. Timber production,
8 ranching, and other agricultural activities are historical activities that continue to the present day, but the
9 fishery has decreased. Anderson Valley today supports orchards and a growing viticulture industry.

10 **Greenwood Creek**

11 The Greenwood Creek watershed encompasses approximately 25 square miles and is located on the
12 southern Mendocino Coast with Greenwood Ridge as its northern border, Clift Ridge as its southern
13 border, and Signal Ridge as its eastern border. Greenwood Creek is a Class I coastal stream and provides
14 habitat for steelhead and coho salmon.

15 Land use in the watershed is primarily for timber production, viticulture, fruit orchards, residential, and
16 some cattle ranching. Most of the watershed is privately owned. Mendocino Redwood Company holds
17 about 60 percent as Timber Production Zone (TPZ) land, and approximately 50 smaller landowners own
18 the rest of the watershed. The only public land in or adjacent to Greenwood Creek is Greenwood State
19 Beach, which contains the Greenwood Creek estuary and a small parcel owned by the Elk County Water
20 District.

21 **Garcia River**

22 The Garcia River watershed encompasses approximately 114 square miles in southwestern Mendocino
23 County. The river forms an estuary that extends from the ocean to the confluence of Hathaway Creek.
24 The floodplains of the lower portion of the watershed are primarily cropland.

25 The primary historical land uses include silviculture, dairy ranching, and gravel mining; these have not
26 changed during the past two decades. Timber-harvesting remains the dominant land use activity, but land
27 conversion to hillside vineyards is becoming a concern for production of sediment. The watershed is
28 completely privately owned by multiple owners. The river and estuary provide habitat for salmonids, and
29 identified beneficial uses include commercial and sport-fishing. The Garcia River has been listed as
30 impaired due to sediment.

31 **Gualala River**

32 The Gualala River watershed encompasses about 300 square miles; the Gualala River flows from
33 Mendocino County to Sonoma County in a north-south direction, reaching the ocean at the town of
34 Gualala. The watershed contains mostly mountainous terrain where tributaries flow through steep valleys
35 with narrow floors that contain erodible soil. Most of the annual precipitation occurs between October and
36 April, with the greatest amounts in January. Rainfall averages about 38 inches per year at the coast and up
37 to 100 inches per year on the inland peaks.

38 The primary historical land uses are silviculture, orchards, and ranching with timber harvest still an
39 important industry. Timber companies own about one-third of the watershed; Gualala Redwoods Inc. is

1 the largest commercial owner, holding about 30,000 acres. Orchards and ranching are on the decline
2 while the watershed has seen an increase in hillside vineyard development, which can threaten to impair
3 water quality with respect to sediment delivery. However, the SWRCB, through regulation, has put into
4 place requirements for runoff protections from hillside vineyards so sediment loading in the rivers from
5 vineyard development should not occur. The Gualala River provides the primary source of drinking water
6 for Sea Ranch and Gualala. The watershed supports an anadromous fishery that includes coho salmon.

7 *Russian River Watershed*

8 The Russian River watershed encompasses 1,485 square miles in Mendocino and Sonoma counties. It is
9 bounded by the Coast Ranges on both the east and west. The main stem is about 110 miles long and flows
10 from north of Ukiah southward through Redwood Valley (Mendocino County) to its confluence with
11 Mark West Creek, where it turns west, passes through the Coast Ranges, and empties into the Pacific
12 Ocean. The summer climate is moist and cool near the coast with temperatures increasing in the valley
13 areas, which are isolated from the cooling coastal influence. During winter, average rainfall ranges from
14 30 to 80 inches, depending on locale.

15 The reservoirs that provide flood protection and water supply storage include Lake Sonoma (Warm
16 Springs Dam) located at the confluence of Warm Springs Creek and Dry Creek west of Healdsburg and
17 Lake Mendocino (Coyote Valley Dam) on the East Fork Russian River near Ukiah. A diversion from the
18 Eel River via the Potter Valley Project (Van Arsdale Reservoir, Cape Horn Dam) for the purpose of
19 power production provides considerable benefit to the overall water storage in Lake Mendocino. The
20 Russian River watershed supplies drinking water for over 570,000 people.

21 The Russian River watershed is primarily an agricultural area with the greatest emphasis on vineyard and
22 orchard crops. Major orchard crops include prunes, pears, and apples; other crops such as cherries and
23 walnuts are also produced. Besides agriculture, there is a growing trend toward light industry and
24 commercial development and a significant telecommunications industry within the region. The production
25 and processing of timber, agricultural and animal products, gravel removal and processing, energy
26 production and miscellaneous light manufacturing operations are additional industrial activities in the
27 watershed. The Russian River watershed also has developed an international reputation for the production
28 of premium wines, contributing to a strong tourism industry within the region.

29 *Bodega Watershed*

30 The Bodega watershed contains streams with headwaters in the Coast Ranges entering the Pacific Ocean
31 south of the Russian River. Salmon, Americano, and Stemple creeks and their associated estuaries are the
32 main water bodies within this watershed. The terrain is relatively steep and erodible and is sensitive to
33 disturbance. Cooler temperatures and relatively high winter rainfall due to coastal influences typify the
34 climate of the Bodega watershed. Because of the Mediterranean climate, summertime flows are often
35 nonexistent in Americano and Stemple creeks; Salmon Creek flow is low but sustained. Each of these
36 subwatersheds have estuary areas; however, the Estero Americano (Americano Creek) and the Estero de
37 San Antonio (Stemple Creek) are prized for their resemblance to fjords and the enhanced resource values
38 associated with isolated estuarine environments.

39 **Groundwater Aquifers**

40 Groundwater resources in the North Coast Hydrologic Region are supplied by both alluvial and fractured
41 rock aquifers. Alluvial aquifers are composed of sand and gravel or finer grained sediments, with

1 groundwater stored within the voids, or pore space, between the alluvial sediments. Fractured-rock
 2 aquifers consist of impermeable granitic, metamorphic, volcanic, and hard sedimentary rocks, with
 3 groundwater being stored within cracks, fractures, or other void spaces. The distribution and extent of
 4 alluvial and fractured-rock aquifers and water wells vary significantly within the region. A brief
 5 description of the aquifers for the region is provided below.

6 *Aquifer Description*

7 **Alluvial Aquifers**

8 The North Coast Hydrologic Region contains 63 DWR Bulletin 118-2003 recognized alluvial
 9 groundwater basins and subbasins which underlie approximately 1,600 square miles, or 8 percent of the
 10 hydrologic region. The majority of the groundwater in the region is stored in alluvial aquifers.
 11 Figure NC-3 shows the location of the alluvial groundwater basins and subbasins and Table NC-1 lists the
 12 associated names and numbers. The most heavily used groundwater basin in the region is the Klamath
 13 River Valley Groundwater Basin, which is located in the northeastern portion of the region along the
 14 Oregon border in Modoc and Siskiyou counties. Other significant groundwater basins in the region are
 15 Santa Rosa Valley, Wilson Grove Formation Highlands, Eel River Valley, Butte Valley, Shasta Valley,
 16 and Scott River Valley.

17 **PLACEHOLDER Figure NC-3 Alluvial Groundwater Basins and Subbasins within the North Coast** 18 **Hydrologic Region**

19 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 20 the end of the report.]

21 **PLACEHOLDER Table NC-1 Alluvial Groundwater Basins and Subbasins within the North Coast** 22 **Hydrologic Region**

23 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 24 the end of the report.]

25 The Klamath River Valley Groundwater Basin is the largest groundwater basin in the North Coast
 26 Hydrologic Region, encompassing approximately 159,000 acres. It is composed of two subbasins — the
 27 Tule Lake and Lower Klamath, by Sheepy Ridge. The primary aquifers in the Klamath River Valley
 28 Groundwater Gasin consist of sand, silt, and clay sediments. Although these deposits are wide spread and
 29 hundreds of feet thick, the permeability of the sediments and therefore, the associated well yields, are
 30 generally low.

31 Santa Rosa Valley Groundwater Basin in Sonoma County is the second largest groundwater basin in the
 32 region, encompassing approximately 101,000 acres. It is composed of three subbasins — the Santa Rosa
 33 Plain, Healdsburg Area, and Rincon Valley. The Santa Rosa Plain Subbasin covers an area of
 34 approximately 80,000 acres and is home to approximately half of the population of Sonoma County. The
 35 subbasin's best water-producing units are stream channels filled with alluvial sands and gravels, basin-fill
 36 alluvium, and alluvial fan deposits that connect the Santa Rosa Plain with its bordering hills, and massive
 37 sandstone units of the Wilson Grove Formation. The Sonoma Volcanics, a thick sequence of lava flows
 38 present along the eastern boundary of the basin, produce variable amounts of water. The Glen Ellen
 39 Formation consists of continental deposits of partially cemented gravel, sand, silt, and clay, and yields
 40 modest amounts of water to smaller groundwater wells. Groundwater within the Santa Rosa Plain

1 Subbasin is generally present under confined conditions, except locally in the vicinity of clay or silt
2 horizons where conditions may be semi-confined or confined (Sonoma County Water Agency,
3 Groundwater Level Monitoring Plan for CASGEM, December 2011).

4 The Wilson Grove Formation Highlands Groundwater Basin covers approximately 80,000 acres and is
5 located in southwestern Sonoma County and northwestern Marin County. The primary groundwater-
6 bearing formation is the marine sedimentary deposits of the Wilson Grove Formation. This formation
7 consists of fine-grained sandstone with lenses of conglomerate and shale. The formation underlies most of
8 the basin and ranges from 300 to 2,000 feet in thickness. It is moderately permeable due to its high
9 porosity and moderate transmissivity. Well production data for the area is very limited (U.S. Geological
10 Survey 2004).

11 The Eel River Valley Groundwater Basin is in Humboldt County and encompasses approximately
12 73,000 acres. The groundwater basin includes the lower 8 miles of the Van Duzen River Valley and the
13 Eel River Valley. It is bounded by the Little Salmon fault on the north, the Wildcat geologic formation
14 series on the east, and the Carlotta Formation on the south. The primary groundwater-bearing alluvium
15 consists of the gravel, sand, and clay that underlie the floodplain of the Eel River. In the delta, the
16 thickness of the alluvium is estimated to be approximately 75 feet; in the valley portions of the region, the
17 gravel layer can be as thick as 200 feet. Wells constructed in these gravels can yield up to 600 gallons per
18 minute (gpm) (California Department of Water Resources 1965).

19 The Butte Valley Groundwater Basin is located in northeastern Siskiyou County and encompasses
20 approximately 79,000 acres. The primary groundwater-bearing formations are alluvial fan and lake
21 deposits. The alluvial fan deposits with thickness up to 350 feet, found on the west side of the valley,
22 have low permeability and yield small quantities of water. Lake deposits composed of fine grained silts
23 and clays have very low permeability; coarser layers in the western and northwestern part of the basin
24 generally yield sufficient water for stock wells. Well yield data indicate that groundwater production in
25 the Butte Valley Groundwater Basin varies between 200 gpm and 5,000 gpm, with an average yield of
26 about 2,300 gpm.

27 The Shasta Valley Groundwater Basin is located in central Siskiyou County and encompasses
28 approximately 52,000 acres. The primary groundwater-bearing formation is alluvium consisting of beds
29 of gravel, sand, silt, and clay with thickness up to 140 feet. The productivity of the alluvial deposits varies
30 greatly with well yields ranging from 150 gpm to 1,000 gpm. Most wells producing groundwater from the
31 alluvium throughout the valley are used for domestic and stock water and the majority of the irrigation
32 wells are on the western margin of the valley.

33 The Scott River Valley Groundwater Basin is located in southwestern Siskiyou County and covers
34 approximately 63,000 acres. The primary groundwater-bearing formations in the Scott River Valley are
35 the stream channel, floodplain, and alluvial fan deposits. The stream channel and floodplain deposits
36 consist of layered gravel, sand, and clay deposited by the Scott River and its tributaries with thickness up
37 to 260 feet or more. Data indicate that the greatest production is most often found along the east side of
38 the valley between Etna and Fort Jones. Wells in other areas may only produce sufficient quantities for
39 domestic users, and larger irrigation wells often produce 1,200 gpm to 2,500 gpm. The alluvial fan
40 deposits consist of sandy clay with some intermixed boulders deposited by the Scott River tributaries on
41 the west side of the valley. Little production data are available for wells constructed in these deposits. The

1 highly recognized interconnection between surface water and groundwater resources in Scott River
2 Valley resulted in the 1980 adjudication for both surface water and groundwater. A pending 2010 lawsuit
3 by Environmental Law Foundation seeks to apply the Public Trust Doctrine to further regulate
4 interconnected surface water — groundwater areas outside the “interconnected” area defined in the
5 1980 Scott River Adjudication.

6 **Fractured-Rock Aquifers**

7 Fractured-rock aquifers in the foothill and mountain areas adjacent to the many alluvial groundwater
8 basins also provide groundwater supply in the region. Groundwater from fractured rock aquifers tends to
9 supply individual domestic and stock wells, or small community water systems. Fractured rock aquifers,
10 and the wells that they supply, tend to have less capacity and reliability than the wells in alluvial aquifers.
11 However, localized fractured rocks within the Klamath, Butte, and Shasta Valley groundwater basins tend
12 to form some of the most highly productive fractured-rock aquifers in California.

13 In Klamath River Valley Groundwater Basin, the two major fractured rock aquifers are the Pleistocene
14 Intermediate Basalt and the Miocene to Pliocene Lower Basalt. Fracturing of the Pleistocene Intermediate
15 Basalt appears to be extensive resulting in high permeability and high well-yield in most locations
16 proving yield between 2,000 gpm and 4,000 gpm. The Miocene to Pliocene Lower Basalt is a jointed and
17 fractured basalt that underlies the lake deposits in most locations within the groundwater basin,
18 encountered in wells as deep as 1,200 feet in the center of the Tule Lake Subbasin. Wells constructed as
19 part of the 2001 Emergency Well Drilling Program in the Tule Lake Subbasin produce groundwater
20 primarily from the lower basalt at rates from 6,000 gpm to 12,000 gpm. However, groundwater elevation
21 data collected from these wells indicate that recharge within the basin are slow.

22 In Butte Valley Groundwater Basin, the primary fractured rock aquifers are the Butte Valley Basalt, the
23 Holocene and Pleistocene Pyroclastic Rocks, and the High Cascade Volcanics. The Butte Valley Basalt is
24 highly permeable, fractured, and vesicular basalt located primarily in the southern and southeastern
25 portions of the groundwater basin. The basalt occurs at depths of generally less than 150 feet below
26 ground surface and averages about 40 feet thick. The Butte Valley Basalt yields large amounts of
27 groundwater to wells; however, a study completed in the early 1980s determined that this aquifer was
28 already developed to its maximum productivity. The Holocene and Pleistocene Pyroclastic Rocks consist
29 of well consolidated tuffs and tuff breccias with great variability in permeability and porosity. The
30 deposits are up to 400 feet thick. Well yields vary greatly, but the formation is developed extensively for
31 stock supply wells. The High Cascades Volcanics consist of successive layers of basalt, tuff, tuff breccia,
32 and cinder-cone deposits; and the units generally range in thickness between 10 and 50 feet. DWR
33 Bulletin 118-2003 identified the average well yield in the Pluto’s Cave Basalt aquifer at 1,300 gpm, with
34 yields up to 4,000 gpm being recorded. A more recent review of 142 well drillers logs indicated an
35 average well yield of about 350 gpm and a maximum yield of about 1,400 gpm (California Department of
36 Water Resources 2011).

37 In Shasta Valley Groundwater Basin, the predominant fractured aquifer the Holocene age Pluto's Cave
38 Basalt is a highly productive and locally valuable fractured-rock aquifer. Williams (1949) describes the
39 formation as a series of overlapping flow units ranging in thickness from about 10 to 30 feet.
40 Groundwater from fractures and local lava tubes provides a significant source of water to wells and
41 springs for irrigation and domestic uses. Groundwater discharge to the Shasta River from the Pluto’s
42 Cave Basalt aquifer is the primary source of cold water inflow to the river during summer months and

1 relatively warmer water in winter months; both are critically important to the fishery (California
 2 Department of Water Resources 2011). DWR Bulletin 118-03 identified the average well yield in the
 3 Pluto’s Cave Basalt aquifer at 1,300 gpm, with yields up to 4,000 gpm. A more recent review of 142 well
 4 drillers’ logs indicate an average well yield of about 350 gpm and a maximum yield of about 1,400 gpm
 5 (DWR, 2011).

6 *More detailed information regarding the aquifers in the North Coast Hydrologic Region is available*
 7 *online from Update 2013 Volume 4, Reference Guide — California’s Groundwater Update 2013 and*
 8 *DWR Bulletin 118-2003.*

9 *Well Infrastructure and Distribution*

10 Well logs submitted to DWR for water supply wells completed during 1977 through 2010 were used to
 11 evaluate the distribution of water wells and the uses of groundwater in the North Coast Hydrologic
 12 Region. DWR does not have well logs for all the wells drilled in the region; and for some well logs,
 13 information regarding well location or use is inaccurate, incomplete, ambiguous, or missing. Hence, some
 14 well logs could not be used in the current assessment. However, for a regional scale evaluation of well
 15 installation and distribution, the quality of the data is considered adequate and informative. The number
 16 and distribution of wells in the region are grouped according to their location by county and according to
 17 six most common well-use types: domestic, irrigation, public supply, industrial, monitoring, and other.
 18 Public supply wells include all wells identified in the well completion report as municipal or public.
 19 Wells identified as “other” include a combination of the less common well types, such as stock wells, test
 20 wells, or unidentified wells (no information listed on the well log).

21 Six counties were included in the analysis of well infrastructure for the North Coast Hydrologic Region.
 22 Del Norte, Humboldt, Mendocino, and Trinity counties are fully contained within the hydrologic region,
 23 while Siskiyou and Sonoma counties are partially within the region. Well log data for counties that fall
 24 within multiple hydrologic regions were assigned to the hydrologic region containing the majority of
 25 alluvial groundwater basins within the county. Well log information listed in Table NC-2 and illustrated
 26 in Figure NC-4 show that the distribution and number of wells vary widely by county and by use. The
 27 total number of wells installed in the region between 1977 and 2010 is approximately 35,000, and ranges
 28 from a high of about 15,800 in Sonoma County to less than 1,300 for Del Norte County. In most counties,
 29 domestic use wells make up the majority of well logs — 10,800 is in Sonoma County, followed by about
 30 5,800 in Mendocino County, and 5,100 in Siskiyou County. The one exception is Humboldt County
 31 where over 60 percent of the wells are monitoring wells. Communities with a high percentage of
 32 monitoring wells compared to other well types may indicate the presence of groundwater quality
 33 monitoring to help characterize groundwater quality issues.

34 **PLACEHOLDER Table NC-2 Number of Well Logs by County and Use for the North Coast** 35 **Hydrologic Region (1977-2010)**

36 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 37 the end of the report.]

PLACEHOLDER Figure NC-4 Number of Well Logs by County and Use for the North Coast Hydrologic Region (1977-2010)

[Any draft tables, figures, and boxes that accompany this text for the public review draft are included at the end of the report.]

Figure NC-5 shows that domestic wells make up the majority of well logs (71 percent) for the region, while irrigation wells account for only about 5 percent of well logs. A higher percentage of domestic wells and lower percentage of irrigation wells point to the more rural-domestic setting and low use of groundwater for irrigation in the region.

PLACEHOLDER Figure NC-5 Percentage of Well Logs by Use for the North Coast Hydrologic Region (1977-2010)

[Any draft tables, figures, and boxes that accompany this text for the public review draft are included at the end of the report.]

Figure NC-6 shows a cyclic pattern of well installation for the region, with new well construction ranging from about 500 to 1,500 wells per year, with an average of about 1,000 wells per year. The large fluctuation of domestic well drilling is likely associated with population booms and residential housing construction. The increase in domestic well drilling in the region during the late 1980s and early 1990s is likely due to increases in housing construction during this time. Similarly, the 2007 to 2010 decline in domestic well drilling is likely due to declining economic conditions and related drop in housing construction.

PLACEHOLDER Figure NC-6 Number of Well Logs Filed per Year by Use for the North Coast Hydrologic Region (1977-2010)

[Any draft tables, figures, and boxes that accompany this text for the public review draft are included at the end of the report.]

The onset of monitoring well installation in the mid- to late-1980s is likely associated with federal underground storage tank programs signed into law in the mid-1980s. The installation of monitoring wells in the region peaked in 1991 at about 470 wells, with an average of about 320 monitoring wells installed per year from 1988 through 1995. From 1998 through 2004, about 300 wells were installed per year. Since 2004, monitoring well installation in the region has averaged approximately 160 wells per year.

As Figure NC-6 shows, irrigation well installation is more closely related to climate conditions, cropping trends and surface water supply cutbacks. Installation of irrigation wells averaged about 50 wells per year up until the late 1990s when the Klamath River Valley Basin and the Tule Lake Subbasin experienced an extended period of drought. From 1998 through 2003, averages of about 90 wells per year were installed. In the years following this long period of drought, installation of wells dropped back down to a rate of about 50 wells per year on average.

More detailed information regarding assumptions and methods of reporting well log information is available online from Update 2013 Volume 4, Reference Guide — California’s Groundwater Update 2013.

1 *California Statewide Groundwater Elevation Monitoring (CASGEM) Basin Prioritization*
 2 The Legislature in 2009, as part of a larger package of water-related bills, passed Senate Bill 7x 6
 3 (SBx7 6; Part 2.11 to Division 6 of the California Water Code § 10920 et seq.), requiring that
 4 groundwater elevation data be collected in a systematic manner on a statewide basis and be made readily
 5 and widely available to the public. DWR was charged with administering the program, which was later
 6 named the “California Statewide Groundwater Elevation Monitoring” or “CASGEM” Program. The new
 7 legislation requires DWR to identify the current extent of groundwater elevation monitoring within each
 8 of the alluvial groundwater basins defined under Bulletin 118-2003. The legislation also requires DWR to
 9 prioritize groundwater basins to help identify, evaluate, and determine the need for additional
 10 groundwater level monitoring by considering available data. Box NC-1 provides a summary of these data
 11 considerations and resulting possible prioritization category of basins. *More detailed information on*
 12 *groundwater basin prioritization is available online from Update 2013 Volume 4, Reference Guide —*
 13 *California’s Groundwater Update 2013.*

14 **PLACEHOLDER Box NC-1 California Statewide Groundwater Elevation Monitoring (CASGEM) Basin**
 15 **Prioritization Data Considerations**

16 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 17 the end of the report.]

18 Figure NC-7 shows the groundwater basin prioritization for the North Coast Hydrologic Region. Of the
 19 63 basins within the region, eight basins were identified as medium priority, two as low priority, and the
 20 remaining 53 basins as very low priority; no basin was identified as either high or very high priority.
 21 Table NC-3 lists the medium CASGEM priority groundwater basins for the region. The eight medium
 22 priority basins account for about 60 percent of the population and about 80 percent of groundwater use for
 23 the region. The basin prioritization could be a valuable tool to help evaluate, focus, and align limited
 24 resources for effective groundwater management, and reliability and sustainability of groundwater
 25 resources.

26 **PLACEHOLDER Figure NC-7 CASGEM Groundwater Basin Prioritization for the North Coast**
 27 **Hydrologic Region**

28 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 29 the end of the report.]

30 **PLACEHOLDER Table NC-3 CASGEM Groundwater Basin Prioritization for the North Coast**
 31 **Hydrologic Region**

32 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 33 the end of the report.]

34 *North Coast Hydrologic Region Groundwater Monitoring Efforts*

35 Groundwater resource monitoring and evaluation is a key aspect to understanding groundwater
 36 conditions, identifying effective resource management strategies, and implementing sustainable resource
 37 management practices. California Water Code (§10753.7) requires local agencies seeking State funds
 38 administered by DWR to prepare and implement groundwater management plans that include monitoring
 39 of groundwater levels, groundwater quality degradation, inelastic land subsidence, and changes in surface
 40 water flow and quality that directly affect groundwater levels or quality. This section summarizes some of

1 the groundwater level, groundwater quality, and land subsidence monitoring efforts within the North
 2 Coast Hydrologic Region. Groundwater level monitoring well information includes only active
 3 monitoring wells — those wells that have been measured since January 1, 2010. *Additional information*
 4 *regarding the methods, assumptions, and data availability associated with the groundwater monitoring is*
 5 *available online from Update 2013 Volume 4, Reference Guide — California’s Groundwater Update*
 6 *2013.*

7 **Groundwater Level Monitoring**

8 A list of the number of monitoring wells in the North Coast Hydrologic Region by monitoring agencies,
 9 cooperators, and CASGEM monitoring entities is provided in Table NC-4. The locations of these
 10 monitoring wells by monitoring entity and monitoring well type are shown in Figure NC-7. Table NC-4
 11 shows that a total of 194 wells in the region have been actively monitored for groundwater levels since
 12 2010. DWR monitors a total of 123 wells — 90 wells in 15 basins and 33 wells outside Bulletin 118-2003
 13 alluvial basins; the U.S. Geological Survey (USGS) monitors 37 wells in 10 basins; and three CASGEM
 14 monitoring entities monitor 34 wells in 6 basins. A comparison of Figure NC-7 discussed previously and
 15 Figure NC-8 indicate that several basins identified as having a medium priority under the CASGEM
 16 groundwater basin prioritization have not been monitored for groundwater levels.

17 **PLACEHOLDER Table NC-4 Groundwater Level Monitoring Wells by Monitoring Entity in the North** 18 **Coast Hydrologic Region**

19 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 20 the end of the report.]

21 **PLACEHOLDER Figure NC-8 Monitoring Well Location by Agency, Monitoring Cooperator, and** 22 **CASGEM Monitoring Entity in the North Coast Hydrologic Region**

23 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 24 the end of the report.]

25 The groundwater level monitoring wells are categorized by the type of well use and include domestic,
 26 irrigation, observation, public supply, and other. Groundwater level monitoring wells identified as “other”
 27 include a combination of the less common well types, such as stock wells, test wells, industrial wells, or
 28 unidentified wells (no information listed on the well log). Wells listed as “observation” also include those
 29 wells described by drillers in the well logs as “monitoring” wells. Domestic wells are typically relatively
 30 shallow and are in the upper portion of the aquifer system, while irrigation wells tend to be deeper and are
 31 in the middle-to-deeper portion of the aquifer system. Some observation wells are constructed as a nested
 32 or clustered set of dedicated monitoring wells, designed to characterize groundwater conditions at specific
 33 and discrete production intervals throughout the aquifer system. Figure NC-9 shows that wells identified
 34 as irrigation and other account for 36 and 33 percent, respectively, of the monitoring wells in the region,
 35 while wells listed as domestic comprise 19 percent of the total; public supply wells comprise less than 3
 36 percent of the total.

37 **PLACEHOLDER Figure NC-9 Percentage of Monitoring Wells by Use in the North Coast Hydrologic** 38 **Region**

39 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 40 the end of the report.]

1 **Groundwater Quality Monitoring**

2 Groundwater quality monitoring is an important aspect to effective groundwater basin management and is
3 one of the components that are required to be included in groundwater management planning in order for
4 local agencies to be eligible for State funds. Numerous State, federal, and local agencies participate in
5 groundwater quality monitoring efforts throughout California. A number of the existing groundwater
6 quality monitoring efforts were initiated as part of the Groundwater Quality Monitoring Act of 2001,
7 which implemented goals to improve and increase the statewide availability of groundwater quality data.
8 A summary of the larger groundwater quality monitoring efforts and references for additional information
9 are provided below.

10 Regional and statewide groundwater quality monitoring information and data are available on the
11 SWRCB Groundwater Ambient Monitoring and Assessment (GAMA) Web site and the GeoTracker
12 GAMA groundwater information system developed as part of the Groundwater Quality Monitoring Act of
13 2001. The GAMA Web site describes GAMA program and provides links to all published GAMA and
14 related reports. The GeoTracker GAMA groundwater information system geographically displays
15 information and includes analytical tools and reporting features to assess groundwater quality. This
16 system currently includes groundwater data from the SWRCB, Regional Water Quality Control Boards
17 (RWQCBs), California Department of Public Health (CDPH), Department of Pesticide Regulation
18 (DPR), DWR, USGS, and Lawrence Livermore National Laboratory (LLNL). In addition to groundwater
19 quality data, GeoTracker GAMA has more than 2.5-million depth to groundwater measurements from the
20 Water Boards and DWR, and also has oil and gas hydraulically fractured well information from the
21 California Division of Oil, Gas, and Geothermal Resources. Table NC-5 provides agency-specific
22 groundwater quality information. Additional information regarding assessment and reporting of
23 groundwater quality information is furnished later in this report.

24 **PLACEHOLDER Table NC-5 Sources of Groundwater Quality Information**

25 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
26 the end of the report.]

27 **Land Subsidence Monitoring**

28 Land subsidence has been shown to occur in areas experiencing significant declines in groundwater
29 levels. Most groundwater basins along the coastal portion of the North Coast Hydrologic Region have
30 limited risk for land subsidence due to groundwater withdrawal. Consequently, no land subsidence
31 monitoring efforts are known to exist along the coastal portion of the region. However, recent increases in
32 groundwater withdrawals for some inland groundwater basins have resulted in the installation of land
33 subsidence monitoring instruments, for example, in the Tule Lake subbasin. The Tule Lake GPS land
34 subsidence monitoring network was put in place in 2001 and 2002 as part of the 2001 Klamath Basin
35 Drought Emergency and in response to concerns about the potential for land subsidence in the thick
36 lakebed deposits of the basin after constructing several deep, high-production agricultural supply wells.
37 The existing GPS land subsidence network consists of 23 stations, with 16 stations within the Tule Lake
38 subbasin and 7 stations within hard rock along the outside edge of the basin.

39 **Ecosystems**

40 Natural ecosystems are the result of the interactions of the abiotic and biotic (nonliving and living)
41 components that interact as a unit. The climate, location, soil, biota, and topography of the North Coast

1 region have contributed to the development of large ecosystems that have come to characterize the region.
 2 Major ecosystems of the region include forests, estuaries and coastal tidelands, riverine, and sagebrush
 3 steppe.

4 Conditions in the region are conducive to forest ecosystems. From an ecosystem perspective, a forest
 5 ecosystem comprises all its plants, animals, and other organisms as well as the natural woodland units.
 6 Forests store large amounts of water because of their large size and physiological characteristics. They are
 7 important regulators of hydrologic processes, especially those involving groundwater, evaporation, and
 8 precipitation patterns. Forests accumulate large amounts of biomass and have been referred to as the most
 9 effective land cover for maintaining water quality. Forest cover has been directly linked to drinking water
 10 treatment costs: The more forest in a source watershed, the lower the treatment costs.

11 An estuary is a coastal area where fresh water from rivers and streams meets and mixes with salt water
 12 from the ocean. Estuaries and littoral (near shore) ecosystems are very significant to the North Coast
 13 because they provide feeding and nesting habitat for many species of waterfowl and shore birds and are
 14 an important feature for migratory birds along the Pacific Flyway. Estuaries and coastal ecosystems are
 15 valuable to foraging sea birds and marine mammals. Estuaries function as feeding and sheltering habitats
 16 for salmonids. The North Coast Hydrologic Region includes 340 miles of coastline.

17 Tidelands and marshes too are extremely important to many species of waterfowl and shore birds, both
 18 for feeding and nesting. Cultivated land and pasture lands also provide supplemental food for many birds,
 19 including small pheasant populations. Tideland areas along the North Coast provide important habitat for
 20 marine invertebrates and nursery areas for forage fish, game fish, and crustaceans. Offshore coastal rocks
 21 are used by many species of seabirds as nesting areas.

22 Riverine ecosystems are those environments that relate to, are formed by, or situated on streams or rivers.
 23 These systems are complex and result from the physical, chemical, and biological processes acting upon
 24 that system. Many of the rivers of the North Coast retain functional habitats and geomorphic processes
 25 but are affected by land use practices and invasion of non-native plants. The life cycle of salmonids is
 26 closely interwoven with water quality and quantity and, therefore, is an excellent indicator of the “health”
 27 of streams and rivers.

28 The common perception of the North Coast ecosystems are related to the forests, rivers, and proximity to
 29 the ocean. However, in the northeastern portion of this region, Modoc and Siskiyou counties, sagebrush
 30 steppe ecosystems are predominant. A sagebrush steppe ecosystem is largely treeless and dry with
 31 dominant plant communities consisting of sagebrush shrubs and short bunchgrasses.

32 **Flood**

33 In the North Coast Hydrologic Region, forest management practices are the most significant issue
 34 impacting flood management (See "Flood Management" subsection under "Regional Resources
 35 Management Conditions in this report). Maintaining the natural attenuation and function of floodplains in
 36 this hydrologic region will help to protect more than 320 sensitive species that live in the floodplains.
 37 Another issue is coastal flooding, including tsunamis, which can impact more than \$4 billion in assets
 38 (crops, buildings, and public infrastructure). See Box NC-2 "Near Coastal Issues." In addition, illegal
 39 cultivation of marijuana in the forests — with over fertilization and pesticide use, land clearing and illegal

1 water diversions — sets the stage for increased runoff during rain events carrying toxics and sediment
2 into the streams and rivers, degrading the environment.

3 **PLACEHOLDER Box NC-2 Near Coastal Issues**

4 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
5 the end of the report.]

6 Communities in the North Coast Hydrologic Region have suffered frequent flood damage since the winter
7 of 1861 when devastating floods were recorded. Torrential rains caused flooding throughout the
8 hydrologic region in 1937. Winter floods between 1935 and 1945 in Sonoma County spurred the U.S.
9 Army Corps of Engineers (USACE) to develop a flood management plan and construct Coyote Valley
10 Dam, which impounded Lake Mendocino upon completion in 1957 (U.S. Army Corps of Engineers,
11 Coyote Valley Dam 2010).

12 For a complete record of floods, refer to the California Flood Future Report Attachment C: Flood History
13 of California Technical Memorandum.

14 *Recent Tsunamis on the California Coast*

15 This region was struck by a tsunami in March 1964 as a result of an earthquake in Prince William Sound,
16 off the south coast of Alaska. The earthquake generated a tsunami that towered more than 20 feet when it
17 made landfall on the North Coast. The huge wave smashed into Crescent City in the early morning of
18 March 28 and devastated the community. Parts of Citizens Dock, a major distribution hub for the city's
19 bustling natural resources industry, were completely wrecked; and several fishing vessels were capsized.
20 The massive wave damaged 289 homes and businesses; 11 people were killed; and 3 were never found.
21 Damages were estimated at \$16 million in 1964 dollars.

22 In March 2011, a tsunami generated off the coast of Japan, recorded throughout the California coast,
23 struck Crescent City Harbor with an 8.1-foot wave, destroying much of the harbor and resulting in one
24 death near Klamath. There was also major damage to docks and boats at Noyo Harbor. Estimated damage
25 in the region was \$24 million.

26 **Climate**

27 Weather conditions vary dramatically within the North Coast Hydrologic Region from the cooler coastal
28 areas to the arid inland valleys in Siskiyou and Modoc counties. In the western coastal portion of this
29 region, average temperatures are moderated by the influence of the Pacific Ocean and range from highs in
30 the mid-80s in the summer to lows in the mid-30s during the winter. In the inland regions of Siskiyou and
31 Modoc counties, temperatures are more variable, where summer high temperatures usually reach the
32 100-degree mark and winter low temperatures are often in the low-30-degree range. The heavy rainfall
33 over the mountainous portions of the region makes it the most water-abundant area of California. Mean
34 annual runoff is about 29 million acre-feet (maf), which constitutes about 41 percent of the state's total
35 natural runoff, which is the largest volume compared to all other hydrologic regions of California. The
36 major rivers in decreasing order of average annual runoff are: the Klamath with 11 maf; the Eel, 6 maf;
37 the Smith, 3 maf; the Russian, 1.6 maf; the Mad, 1 maf, and the Mattole, 1 maf. The principal reaches
38 (and tributaries) of the Klamath, Eel, and Smith rivers have been designated wild and scenic under federal
39 and State law. Annual average precipitation in the North Coast region is 53 inches, ranging from over

1 100 inches per year in eastern Del Norte County to less than 15 inches annually in the Lost River drainage
 2 area of Modoc County. A relatively small fraction of the precipitation is in the form of snow; only at
 3 elevations above 4,000 feet does snow remain on the ground for appreciable periods.

4 Precipitation, or rainfall, varies greatly within the North Coast region depending upon location and time
 5 of year. The combination of mountainous terrain with high peaks and steep narrow valleys compared to
 6 higher elevation plateaus present conditions favorable to variable rainfall patterns. In general,
 7 precipitation is higher in the northwest mountains and decreases toward the east and southeast.

8 In the coastal communities to the north near Crescent City in Del Norte County, average precipitation for
 9 the period from 1971 through 2000 is about 64 inches with the highest rainfall normally during
 10 December. At Eureka in Humboldt County, average precipitation for the same period is about 48 inches.
 11 At Fort Bragg in Mendocino County along the coast, it is about 43 inches; at Bodega Bay in Sonoma
 12 County, about 37 inches.

13 In the mountains within the coastal counties, precipitation increases (compared to the coastal
 14 communities) due to the orographic effect causing moisture in the air to condense and fall as rain or snow.
 15 At Ship Mountain in Del Norte County with an elevation of approximately 5,320 feet, about 145 inches of
 16 rainfall occurs annually with the highest rainfall during the month of December. Moving south to Spike
 17 Buck Mountain in Humboldt County at approximately 5,480 feet, about 61 inches of rainfall occurs on
 18 average. In Mendocino County along Chamberlain Ridge at 2,020 feet elevation, about 48 inches of
 19 rainfall occurs with the highest precipitation during the month of January. At Sonoma Mountain in
 20 Sonoma County, at 2,460 feet elevation, precipitation averages about 29 inches with the heaviest amounts
 21 falling during January.

22 Moving inland toward northeast California, at Boulder Peak in Siskiyou County at 8,300 feet, about
 23 47 inches of rainfall normally occurs with the heaviest rainfall happening in January. Moving farther east
 24 to Mount Shasta in Siskiyou County at about 14,160 feet, average rainfall and snow amounts to near
 25 56 inches with the highest rainfall occurring during January. In contrast, at Weed in Siskiyou County at
 26 approximately 3,550 feet elevation and only 10 miles away from Mount Shasta (air miles), the average
 27 rainfall is about 31 inches. Moving to eastern Siskiyou County at Mount Hoffman near 7,910 feet
 28 elevation, about 47 inches of rainfall occurs.

29 In western Modoc County (the eastern portion of the North Coast region), representative precipitation in
 30 the Tule Lake agricultural area at the town of Newell, 4,042 feet elevation, amount to near 12 inches
 31 annually with November, December, and January having the highest amounts. At Blue Mountain near the
 32 eastern edge of the North Coast region at 5,750 feet elevation (about 27 air miles from the town of Newell
 33 and an increase of about 1,700 feet in elevation), precipitation amounts to an average of about 21 inches
 34 per year.

35 **Demographics**

36 The North Coast region includes all residents of Del Norte, Humboldt, Trinity, and Mendocino counties,
 37 the majority of Modoc, Siskiyou, and Sonoma counties, and a small percentage of the populations of
 38 Glenn, Lake and Marin counties.

1 *Population*

2 According to the California Department of Finance (DOF), the population of the entire North Coast
3 region was about 670,700 in year 2010, which is less than 2 percent of California's total population. More
4 than half of this region's population lives in its southern part, primarily in Santa Rosa and the surrounding
5 communities of Cotati, Healdsburg, Rohnert Park, Sebastopol, and Windsor along the Russian River
6 watershed. Urban growth in these cities, 261,485 people in year 2010 (DOF), is heavily influenced by the
7 overall urban expansion of the adjacent San Francisco Bay region.

8 The majority of the North Coast region's population (2010 U.S. Census, California Department of
9 Finance) is concentrated in the southern portion of the region, in Sonoma and Marin counties, with
10 370,025 and 316 residents respectively, or approximately 55 percent of all inhabitants in the region. Only
11 a portion of these two counties are in this hydrologic region. The remainder is part of the San Francisco
12 Bay Hydrologic Region. Marin County and part of Sonoma County are part of the nine-county Bay Area
13 Association of Bay Area Governments (ABAG). For additional information on ABAG, see:
14 <http://www.abag.ca.gov/>. Mendocino and Humboldt counties comprise 87,812 and 134,623 residents,
15 respectively.

16 The remainder of the region's population is distributed in its north/northeast and southeast sections. In the
17 north/northeast areas, Del Norte County had 28,610 residents, and Siskiyou County included 44,900
18 citizens. Other smaller communities in the northern portions of the region include Eureka, 27,191; Ukiah,
19 16,075; Arcata, 17,231; Crescent City, 7,643; and Yreka, 7,765 (California Department of Finance, U.S.
20 Census Data 2010). In 2007, the Eureka area had 114,362 people accounting for about 85 percent of the
21 population in Humboldt County. The Eureka area includes Blue Lake, Eureka, Ferndale, Fortuna,
22 McKinleyville, and adjacent towns. Three counties represent the southeast section's population: Glenn
23 with 0, Lake with 61 and Trinity with 13,881 residents (Association of Bay Area Governments 2008).
24 Trinity County is wholly contained within the region while Glenn and Lake counties are only partially
25 represented.

26 The North Coast region has experienced steady population growth over the past two decades and is
27 projected to continue positive growth through the year 2050 (California Department of Finance 2010).
28 Due to the rural nature of much of the region and the fact that there is a lower associated cost of living,
29 many communities within the region are seeing an influx of retirees from larger, more urbanized settings.
30 This has placed pressure on existing community services. Additionally, as population densities encroach
31 in the more urban settings, some of the more rural communities are becoming bedroom communities.
32 There is also a rise in migrant workers within the region. Modoc County has a county-operated migrant
33 camp. For both Modoc and Siskiyou counties, many of their migrant workers are becoming permanent
34 residents. Meanwhile, younger residents continue to leave the area seeking higher-paying jobs.

35 When compared with the year 2000 regional population of 636,000, the 670,300 in 2010 represents a
36 growth rate of 5.4 percent over the 10 years, which is a little more than half the statewide growth rate of
37 about 9.7 percent over the same period. Recent projections indicate that the regional population is
38 expected to grow to about 809,400 by year 2050, which represents approximately 21 percent increase
39 from year 2010 totals. Figure NC-10 provides a graphical depiction of the North Coast region's total
40 population from year 1960 through year 2010, with current projections to year 2050. More than half of
41 this projected growth is anticipated to occur in the Santa Rosa region, as urban populations from the San
42 Francisco Bay area continue to expand northward. Population increases in the rural communities in the

1 northern portion of this region are projected to grow more slowly due to the geographic location, few
2 transportation corridors, and a lack of adequate harbors.

3 **PLACEHOLDER Figure NC-10 Total Population 1960-2010 (2050), North Coast Hydrologic Region**

4 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
5 the end of the report.]

6 Despite the overall growth rates of the region, population growth rates are not as great as those of the rest
7 of the state, reflecting its rural character. In fact, some of the more remote counties of the region —
8 Modoc and Siskiyou — are projected to lose overall population in the coming decades. The most
9 populated area of the region, Sonoma County, experienced a higher growth rate than the state’s average in
10 1980 and 1990, and is estimated to continue this pattern with population increases of 15 percent and 14
11 percent by 2010 and 2020, respectively. Figure NC-11 describes the historic and projected population
12 growth trends for the North Coast Region.

13 **PLACEHOLDER Figure NC-11 Population Growth Trends, North Coast Hydrologic Region**

14 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
15 the end of the report.]

16 *Tribal Communities*

17 In the North Coast Hydrologic Region, 4 percent of the residents identify themselves as Native Americans
18 (Indigenous Peoples), significantly higher than in the 1.7 percent in the statewide population. Several
19 tribes live in the North Coast region, but the Yurok Tribe is the largest in both the North Coast and
20 California. Many of the tribes here are federally recognized, but some are not (Tables NC-6 and NC-7). In
21 addition, many tribes exist within the region, with both large and small numbers of registered individuals.
22 However, many tribes existing within the region have not obtained Federal Reservation status as of the
23 writing of this report. The following subsections include information on the Yurok Tribe and Reservation,
24 the Hoopa Valley Reservation, and the Round Valley Reservation.

25 **PLACEHOLDER Table NC-6 Federally Recognized Tribes in North Coast Hydrologic Region**

26 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
27 the end of the report.]

28 **PLACEHOLDER Table NC-7 California Native American Tribes (Non-Recognized) in North Coast
29 Hydrologic Region**

30 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
31 the end of the report.]

32 **Yurok Tribe and Reservation**

33 The Yurok Tribe is the largest tribe in California with more than 5,000 enrolled members. The tribe
34 provides numerous services to the local community and membership with its more than 200 employees.
35 The tribe’s major initiatives include the Hoopa-Yurok Settlement Act, dam removal, natural resources
36 protection, sustainable economic development enterprises, and land acquisition. The Yurok Tribe’s
37 territory consists of all Ancestral Lands, specifically including but not limited to the Yurok Reservation
38 lands, which extend from one mile on each side from the mouth of the Klamath River and upriver for a

1 distance of 44 miles. The Yurok Tribe's people are also known historically as the Pohlik-la, Ner-er-er,
 2 Petch-ik-lah, and Klamath River Indians. The Yurok Reservation (Yurok lands) includes 63,035 acres.
 3 Only a small portion of the Yurok Reservation has been developed for residential housing, and much of
 4 that lacks basic services such as electricity and telephone (Yurok Tribe 2006a).

5 The Yurok Tribe is in the process of establishing a hotel-casino in Klamath, California, with proceeds to
 6 go toward improving conditions for tribal members. Improvements will include electricity, potable water,
 7 and telephone services within the reservation. In addition, a per capita distribution plan will include a one-
 8 time dispersion of funds to all tribal members regardless of age concerning past timber harvesting on
 9 Yurok lands (SPOA Settlement 2013).

10 **Hoopa Valley Reservation**

11 The People of Hoopa Valley are one of California's first cultures. In 1864, a Peace and Friendship Treaty
 12 was negotiated with the United States. In 1896, the Department of the Interior began preparing a land
 13 allotment list, and in 1909 a proclamation was handed down by President Theodore Roosevelt. This list
 14 was not completed and approved until 1923. The "Hupa People" successfully avoided the physical
 15 destruction of their valley homeland, and in modern times created one of the first successful self-
 16 governance tribal structures in the nation. According to the 2010 U.S. Census, the Hoopa Valley
 17 Reservation includes 3,041 people with 82.4 percent of Native American heritage.

18 The Hupa people traditionally occupied lands in the far northwestern corner of California. The boundaries
 19 of the reservation were established by Executive Order on June 23, 1876, pursuant to the Congressional
 20 Act of April 3, 1864. The boundaries were expanded by executive order in 1891 to connect the old
 21 Klamath River (Yurok) Reservation to the Hoopa Valley Reservation. Further confirmation of the
 22 ownership by the Hupa Tribe of the Hoopa Valley Reservation came on October 31, 1988, with President
 23 Ronald Reagan's signature on Public Law 100-580, the Hoopa/Yurok Settlement Act.

24 The Hupa People have occupied their lands since time immemorial, and the past century has really been
 25 the shortest in their history. However, up until the late 1800s, there is little or no written record on the rich
 26 history and culture that is now the Hoopa Valley Tribe. Much of the tradition and lore that exists today
 27 has been passed along between generations via an extensive oral tradition. The ceremonies and traditions
 28 continue in the similar manners as they have since the beginning, and will continue into the future (Hoopa
 29 Valley Indian Tribe 2003).

30 **Round Valley Reservation (A Sovereign Nation of Confederated Tribes)**

31 The Round Valley Indian Reservation is federally recognized, lying primarily in northern Mendocino
 32 County with a small part of it extending northward into southern Trinity County. The total land area,
 33 including off-reservation trust land, is 36.27 square miles. More than two-thirds of this area is off-
 34 reservation trust land, including about 405 acres in the community of Covelo. Population estimates for
 35 2010 show just over 3,000 people are tribal members with about half living on the reservation (Center for
 36 Applied Research 2010).

37 The Round Valley Indians consist of the Covelo Indian Community. This community is an accumulation
 38 of small tribes; the Yuki (who were the original inhabitants of Round Valley), Concow Maidu, Little
 39 Lake and other Pomo, Nomlaki, Cahto, Wailaki, and Pit River peoples. These tribes were forced onto the
 40 land formerly occupied by the Yuki tribe. From years of intermarriage, a common lifestyle and a shared

1 land base, a unified community emerged. The descendants of these peoples formed a new tribe on the
 2 reservation, the Covelo Indian Community, later to be called the Round Valley Indian Tribes. Their
 3 heritage is a rich combination of different cultures with a common reservation experience and history.
 4 (Round Valley Indian Tribes 2010, 2011).

5 *Disadvantaged Communities*

6 Disadvantaged Community status is determined based on the DAC definition provided in Proposition 84
 7 and 1E IRWM Grant Guidelines (see:
 8 http://www.water.ca.gov/irwm/docs/Guidelines/GL_Final_07_20_10.pdf), dated August 2010. A Median
 9 Household Income (MHI) of less than \$48,706 is the DAC threshold in California (80 percent of the
 10 statewide MHI of \$60,882). In 2010, households in California included an average of 2.89 people.
 11 According to the 2010 U.S. Census, 47.64 percent of all census blocks in this region are within a DAC.
 12 This amounts to 46.25 percent of the North Coast region's population considered as disadvantaged.
 13 (See Figure NC-12.)

14 **PLACEHOLDER Figure NC-12 Disadvantaged Communities in the North Coast Hydrologic Region**

15 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 16 the end of the report.]

17 The NCIRWMP places a strong emphasis on ensuring the inclusion of DACs in the planning and
 18 implementation process. DACs have been involved in all aspects of the NCIRWMP planning effort from
 19 its inception, including plan review and input, attendance, and participation at meetings; DACs comprise
 20 a substantial portion of the priority project proponents who are currently implementing projects. See
 21 http://www.northcoastirwmp.net/Content/10344/North_Coast_IRWMP_Implementation_Projects.html
 22 for more information (North Coast Regional Integrated Water Management Plan, Phase III c2012).

23 The NCIRWMP identifies six primary objective (see "Water in the Environment" under "Regional
 24 Resource Management Conditions). One of the objectives is to address environmental justice issues as
 25 they relate to DACs — drinking water quality, and public health. Public discussions indicate that some
 26 believe that DACs in small rural areas and small family businesses carry a disproportionate burden when
 27 communities address environmental issues to protect endangered species, water quality, and natural
 28 resources. These requirements may impact the ability of local businesses (including agriculture,
 29 silviculture and mining), communities, economies and counties to provide services (affects tax base). One
 30 case in point is the removal of dams on the Klamath River. The decision is based on environmental
 31 improvement for fisheries, but it could have adverse effects on Siskiyou and Modoc counties (DAC areas)
 32 e.g., impacts to homeowners by dewatering of lakes with homes built along the shores, loss of
 33 recreational income for the counties, and decrease in flood control capability.

34 **Land Use Patterns**

35 Forest and rangeland represent about 98 percent of this region's land area. Much of the region is
 36 identified as national forests, State and national parks, under the jurisdiction of the federal BLM, and
 37 Native Indian lands such as the Hoopa Valley and Round Mountain reservations. The major land uses in
 38 the North Coast region consist of timber production, agriculture, fish and wildlife management, parks,
 39 recreational areas, and open space. In recent years, the timber industry has declined as a result of

1 economic issues and the expansion of environmental regulations (Timber Harvest Levels on the Major
2 National Forests in Siskiyou County 1978-2009, National Forest Growth 2009).

3 Failure to manage national forests by thinning and harvesting has caused an unnatural massive buildup of
4 biomass which has reduced water available to streams by canopy interception of snow and
5 evapotranspiration.

6 Vacationers, boaters, anglers, and sightseers are attracted to the region's 340 miles of scenic ocean
7 shoreline, including nearby forests with more than half of California's redwoods. The inland regions are
8 mountainous and include 10 wilderness areas run by the USFS. More than 40 State parks, numerous
9 USFS campgrounds, the Smith River National Recreation Area, and the Redwood National Park are
10 within this hydrologic region.

11 Climate, soils, water supply, and remoteness from markets are factors that limit the types of agricultural
12 crops that can be grown in the North Coast region. In the inland valley areas, there is more irrigable land
13 than can be irrigated with existing developed water. The trend in land use has been one of land
14 consolidation to form larger holdings and the conversion of prime agricultural land to urban growth. This
15 trend is a result of low crop values, the lack of additional inexpensive surface water, and the ability to use
16 only the most economically developable groundwater. The cost of environmental regulation and
17 uncertainty of continued water supply for irrigation also contribute to decisions to convert land from
18 agricultural use.

19 Irrigated agriculture in the North Coast region uses most of the region's developed water supplies.
20 Irrigation today accounts for about 81 percent of the region's non-environmental water use, while
21 municipal and industrial use is about 19 percent. About 422,300 acres, or about 3.4 percent of the region,
22 is irrigated. Of that, 276,840 acres lie in the Middle and Upper Klamath River basins, above the
23 confluence of the Salmon and Klamath rivers, where the main irrigated crops are pasture and alfalfa,
24 grain, potatoes, garlic, and a few other assorted truck crops. Agricultural areas in these basins include
25 Scott, Shasta, and Butte valleys and Tule Lake region and account for approximately 65 percent of
26 irrigated agriculture within the North Coast region. Even though the predominant crops in the remainder
27 of the region are pasture and alfalfa, there are significant acreages of other crops including orchards,
28 vineyards, and various row and truck crops. The highest value crops in the region are the substantial acres
29 of grapes and orchards in the Russian River Basin and ornamental flowers and bulbs in Del Norte County.
30 California has designated agricultural water use as a primary beneficial use of water.

31 In the southern portion of the region, the total acreage of fruit and nut orchards decreased over a 15-year
32 period. For example, in Sonoma County, orchards declined from 48,800 acres in 1992 to approximately
33 3,600 acres in 2007. However, the amount of irrigation water used on orchards did not decrease in the
34 same proportion because many of the apple, prune, and walnut orchards taken out of production were not
35 irrigated. In addition, as the acreage of orchards declined, the acreage planted in vineyards increased. In
36 Sonoma County, grape acreage increased from 34,399 acres in 1992 to 57,568 acres in 2007, an increase
37 of 23,169 acres.

38 Most of the newer grape vineyards use drip irrigation systems for irrigation allowing plantings in areas
39 previously unavailable, i.e., sloping hillsides. However, in addition to irrigation for production, overhead
40 sprinklers are used in vineyards for frost protection in the spring and for post-harvest irrigation in the fall,

1 increasing the water demand for this crop over the direct water use by the crop. Land previously
 2 nonirrigated and subsequently placed in production increases the water demand of the region beyond
 3 historical levels. With the development of low pressure drip irrigation systems, farmers are able to move
 4 in to areas unavailable prior to the low pressure technologies. This places a greater demand on the
 5 available water resources requiring surface water infrastructure improvements or reliance on groundwater
 6 (National Agricultural Statistics Service 1994, 2008, 2011; Sonoma County Agricultural Commissioner
 7 2008).

8 According to the 2007 Census of Agriculture, the North Coast Hydrologic Region contained 249 dairy
 9 farms with 54,234 milk cows. This amounted to about 11.5 percent of the dairy farms in California and
 10 about 2.9 percent of the milk cows. The majority of the dairy farms in the North Coast Region in 2007
 11 were in Humboldt County with 82 farms and in Sonoma County with 93 farms. A comparison of 2007 to
 12 2002 data, shows a trend of fewer and larger dairy farms in the region over the 5-year period (National
 13 Agricultural Statistics Service 2002, 2007).

14 Dairies can have water quality impacts resulting from discharges of waste and/or whey to streams, and
 15 from the presence of animals in waterways. The North Coast Regional Water Quality Control Board
 16 (RWQCB) Dairy Regulation Program offers three permitting options: a National Pollutant Discharge
 17 Elimination System (NPDES) permit*, a Waste Discharge Requirements Order, and a Waiver of Waste
 18 Discharge Requirements, depending on the level of risk to water resources. Unlike most other regions, the
 19 dairies in the North Coast are mostly small and family-run, concentrated in southern Sonoma County and
 20 the Eel River delta in Humboldt County. Groundwater impacts (such as nitrates) from dairies have not
 21 been documented, but groundwater monitoring will be performed, pursuant to the Dairy Program
 22 requirements.

- 23 • NPDES; As authorized by the CWA, the NPDES Permit Program controls water pollution by
 24 regulating point sources that discharge pollutants into waters of the United States. Point sources
 25 are discrete conveyances such as pipes or human-made ditches. Since its introduction in 1972,
 26 the NPDES Program has been responsible for significant improvements to the nation's water
 27 quality.

28 Urban acreage in the North Coast region is located primarily in the Eureka area and Russian River basin.
 29 (See content under "Population" in "Demographics" section.)

30 Land use issues in the region include activities causing soil erosion such as road construction, logging,
 31 and hillside agriculture (vineyards), which affects native fish spawning. However, since the principal
 32 reaches of the Klamath, Eel, and Smith rivers have been designated wild and scenic under federal and
 33 State law, they are protected from additional large-scale water development. Many of the region's
 34 watersheds support threatened and endangered species of plants and animals, and many North Coast
 35 streams and rivers support runs of salmon and steelhead trout.

36 *Diversity of Agriculture in North Coast Region*

37 Agriculture in the North Coast region is as diverse as its climate and people. Aquaculture along the coast
 38 to potatoes and wild rice grown in the high desert areas of Modoc County depict the diversity of crop
 39 production in the region. Although grain, hay, pasture, livestock and lumber account for the majority of
 40 crops produced, many other crops provide for the rural lifestyle and setting in Northern California. Each

1 county in the North Coast region is unique in its mix of crops, especially when compared to the region as
2 a whole. Agricultural Commissioners' Crop Reports (available online
3 http://www.nass.usda.gov/Statistics_by_State/California/Publications/AgComm/Summary/index.asp)
4 provide details on each of the counties major crop mix as well as approximate value of each commodity
5 in 2010.

6 Regional Resource Management Conditions

7 Water in the Environment

8 The NCIRWMP identifies six primary objectives for the region. These objectives are consistent with
9 State water management elements, State priorities and objectives, and IRWM Program Preferences. For
10 more information on IRWM, see the following links: <http://www.water.ca.gov/irwm/grants/index.cfm> and
11 <http://www.water.ca.gov/irwm/stratplan/>. The primary objectives are (1) conserve and enhance native
12 salmonid populations by protecting and restoring required habitats, water quality, and watershed
13 processes; (2) protect and enhance drinking water quality to ensure public health; (3) ensure adequate
14 water supply while minimizing environmental impacts; (4) support implementation of Total Maximum
15 Daily Loads (TMDLs), the North Coast RWQCB Watershed Management Initiative, and the Non-Point
16 Source Program Plan; (5) address environmental justice issues as they relate to disadvantaged
17 communities, drinking water quality and public health; (6) provide an ongoing, inclusive framework for
18 efficient intra-regional cooperation, planning, and project implementation.

19 *Instream Fisheries Requirements*

20 SWRCB adopted the North Coast Instream Flow Policy on May 4, 2010. It applies to applications to
21 appropriate water, small domestic use and livestock stock pond registrations, and water right petitions.
22 This policy applies to water diversions from all streams and tributaries discharging to the Pacific Ocean
23 from the mouth of the Mattole River south to San Francisco and all streams and tributaries discharging to
24 northern San Pablo Bay. The policy area includes approximately 5,900 stream miles and encompasses
25 3.1 million watershed acres (4,900 square miles) in Marin, Sonoma, portions of Napa, Mendocino, and
26 Humboldt counties (State Water Resources Control Board 2012).

27 Water Supplies

28 Many of the smaller communities and rural areas in the North Coast region are supplied by small local
29 surface water and groundwater systems. Larger water supply projects in this region include USBR's
30 Klamath Project, the USACE Russian River Project (Potter Valley Project including Lake Mendocino and
31 Lake Sonoma), and the Humboldt Bay Municipal Water District's Ruth Reservoir, which serves coastal
32 communities from Eureka to McKinleyville. Because the Upper Klamath River watershed is in both
33 California and Oregon, the federal Klamath Project includes water supply facilities in both states.
34 Facilities within the California portion include Clear Lake Reservoir for water supply, Tule Lake and
35 Lower Klamath Lake as waterfowl refuges, and Iron Gate Reservoir as a hydroelectric facility of Pacific
36 Power and Light Company. The primary water supply facilities on the Oregon side are Gerber Reservoir
37 and Upper Klamath Lake. The Klamath Project is the largest agricultural irrigation project in the region
38 and supplies water to about 240,000 acres, of which 62 percent is in Oregon and 38 percent is in
39 California. To maintain adequate instream fishery flows for the lower Klamath River, water releases must
40 be coordinated among the various reservoirs operated by different agencies within both states.

1 Two of the largest water supply reservoirs in the North Coast region are USBR’s 2.437-maf Trinity Lake
2 on the Trinity River and the USACE 380,000 acre-foot Lake Sonoma in the Russian River watershed.
3 These facilities provide water for instream flows, recreation, hydropower, and water supply purposes.
4 Water from Trinity Lake is exported from the North Coast region to the Sacramento River region through
5 USBR’s Clear Creek Tunnel. Lake Sonoma is operated to provide flood control and instream flows in the
6 Lower Russian River in Sonoma County. An intrabasin water transfer system known as the Potter Valley
7 Project has been in existence since 1908 and diverts water from the upper reaches of the Eel River at
8 Cape Horn Dam through a tunnel to the East Fork Russian River upstream from Lake Mendocino (see
9 Potter Valley Project" under "Project Operations" section). The water stored behind Coyote Dam (Lake
10 Mendocino, built in 1958) is used to meet instream flow requirements and urban and agricultural needs in
11 the lower Russian River watershed and the Santa Rosa area.

12 Early gold mining activities in the Scott and Shasta River valleys established water rights as early as the
13 1850s and 1860s. These rivers have been declared “fully appropriated” and are adjudicated under decree
14 of the Superior Court of Siskiyou County.

15 *Surface Water*

16 According to DWR (2011)), surface water storage in the North Coast region in 2006, a wet year, was
17 2,060 thousand acre-feet (taf) at the end of November. In 2007, during the beginning period of the most
18 recent drought, surface water storage at the end of November was 1,621 taf. In November 2008, reservoir
19 storage was 1,257 taf; in 2009, it was 1,169 taf; in 2010, 1,892 taf; and in 2011, it was 2,308 taf, showing
20 how variable the water supply can be. For comparison, reservoir storage at the end of November 1977
21 (the driest period in recent years) was 304 taf whereas the wettest period in recent times was in 1983
22 when the North Coast had 2,264 taf of storage (although less than in 2011). This water is used for urban,
23 municipal, rural residential needs, agriculture, State and federal water supply projects, managed wetlands,
24 required Delta outflow, instream flow, and wild and scenic rivers flow. When water supplies fall short, as
25 they did in 2008 and 2009, the wild and scenic rivers and environmental uses receive the largest
26 reductions (California Department of Water Resources 2011).

27 The amount of surface water in the North Coast region is extremely dependent upon precipitation as
28 described above. In very wet years, there may be a surplus; but in drought years, quantity is limited and
29 can become a source of contention between water users. For example, the Klamath Basin has had water
30 shortage problems in the recent past that have led to confrontations between farmers and regulators and
31 farmers and environmentalists. As the population of the region grows, drinking water will continue to
32 experience increases in demand, making the identification of alternative sources for agricultural and
33 landscape irrigation a high priority. The North Coast Regional Water Management Group (NCRWMG)
34 provides the framework for regional cooperation and collaboration to determine the optimal strategies to
35 ensure that surface water supply is able to meet environmental and human-related beneficial uses during
36 both surplus and drought water years. Please refer to Figure NC-13 for the regions inflows and outflows
37 in water year 2010.

38 **PLACEHOLDER Figure NC-13 North Coast Regional Inflows and Outflows in 2010.**

39 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
40 the end of the report.]

1 *Groundwater*

2 The amount and timing of groundwater extraction, along with the location and type of its use, are
 3 fundamental components for building a groundwater basin budget and identifying effective options for
 4 groundwater management. Although some types of groundwater extractions are reported for some
 5 California basins, the majority of groundwater pumpers are not required to monitor, meter, or publicly
 6 record their annual groundwater extraction amounts. Groundwater supply estimates furnished herein are
 7 based on water supply and balance information derived from DWR land use surveys, and from
 8 groundwater supply information voluntarily provided to DWR by water purveyors or other State agencies.

9 Groundwater supply is reported by water year (October 1 through September 30) and categorized
 10 according to agriculture, urban, and managed wetland uses. The associated information is presented by
 11 planning area (PA), county, and by the type of use. Reference to total water supply represents the sum of
 12 surface water and groundwater supplies in the region, and does not take into account local reuse.

13 **2005-2010 Average Annual Groundwater Supply and Trend**

14 In the North Coast Hydrologic Region, there is limited large-scale groundwater development due to the
 15 small number of significant coastal aquifers. Most of the groundwater development has occurred from
 16 shallow wells installed adjacent to rivers. However, as indicated previously, there are significant
 17 groundwater basins underlying the Klamath River valley along the Oregon border and the southern tip of
 18 the region underlying Santa Rosa in Sonoma County. Many domestic and small irrigation wells draw
 19 water from permeable zones within these deposits. Despite the limits on large-scale infrastructure,
 20 groundwater is utilized widely throughout the region for individual domestic, agricultural, and industrial
 21 water uses. Many rural areas rely exclusively on private wells for residential water. There are also an
 22 unknown number of small dams, and water-related infrastructure, which may have a large cumulative
 23 impact on groundwater. Groundwater is a significant water source for some small rural communities that
 24 rely on residential wells for water, but as discussed below groundwater contributes to about one-third of
 25 the total water supply in the region.

26 Groundwater basins in the Redwood Creek watershed are the Redwood Creek Area and Prairie Creek
 27 Area groundwater basins. The Orick Community Services District provides domestic water through a
 28 centralized distribution system that includes two wells located adjacent to Redwood Creek in the northern
 29 part of town. In the Redwood Creek watershed, there are no water development projects such as dams and
 30 surface water diversions.

31 Siskiyou County has developed several codes regarding groundwater. A Groundwater Advisory
 32 Committee has been appointed and is active for Scott Valley (Siskiyou County Code of Ordinances
 33 2012). Adjudication for the Scott Valley includes a defined interrelated groundwater area.

34 The amount of groundwater supply in the North Coast Hydrologic Region varies yearly with
 35 precipitation, infiltration, and the amount of withdrawals from groundwater basins. Withdrawals, in turn,
 36 are in part dependent on the amount of surface water available for municipalities that use both surface and
 37 groundwater for supply needs.

38 Table NC-8 provides the 2005-2010 average annual groundwater supply by PA and by type of use, while
 39 Figure NC-14 depicts the PA locations and the associated 2005-2010 groundwater supply in the region.
 40 The estimated average annual 2005-2010 total water supply for the region is about one maf. Out of the 1-

1 maf total supply, groundwater supply is 364 taf and represents 35 percent of the region’s total water
 2 supply; 42 percent (60 taf) of the overall urban water use and 44 percent (301 taf) of the overall
 3 agricultural water use being met by groundwater. Groundwater contributes to only 1 percent
 4 (2.5 taf) for meeting managed wetland uses in the region. Although statewide, groundwater extraction in
 5 the region accounts for only about 2 percent of California’s 2005-2010 average annual groundwater
 6 supply (see Figure NC-14), it accounts for 100 percent of the domestic supply for many rural
 7 communities in the region and is also heavily relied upon to meet local agricultural uses.

8 **PLACEHOLDER Table NC-8 North Coast Hydrologic Region Average Annual Groundwater Supply**
 9 **by Planning Area and by Type of Use (2005-2010)**

10 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 11 the end of the report.]

12 **PLACEHOLDER Figure NC-14 Contribution of Groundwater to the North Coast Hydrologic Region**
 13 **Water Supply by Planning Area (2005-2010)**

14 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 15 the end of the report.]

16 Regional totals for groundwater based on county area will vary from the PA estimates shown in Table
 17 NC-9 because county boundaries do not necessarily align with PA or hydrologic region boundaries. Del
 18 Norte, Trinity, Humboldt, and Mendocino counties are fully contained within the North Coast Hydrologic
 19 Region, while Siskiyou and Sonoma counties are partially contained within the region. Groundwater
 20 supply for Modoc and Lake Counties are reported in the Sacramento River Hydrologic Region. For the
 21 North Coast Hydrologic Region, county groundwater supply is reported for Del Norte, Siskiyou, Trinity,
 22 Humboldt, Mendocino, and Sonoma counties (Table NC-9). Overall, groundwater contributes to
 23 approximately 40 percent of the total water supply for the six-county area; the range varies from about
 24 30 to 70 percent for individual counties. Groundwater supplies in the six-county area are used to meet
 25 about 50 percent of the agricultural water use and 40 percent of the urban water use.

26 **PLACEHOLDER Table NC-9 North Coast Hydrologic Region Average Annual Groundwater Supply**
 27 **by County and by Type of Use (2005-2010)**

28 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 29 the end of the report.]

30 As shown in Table NC-8 and Figure NC-14, Upper Klamath PA is the largest user of groundwater in the
 31 region with an average annual groundwater supply equal to 192 taf (55 percent of the total groundwater
 32 supply for the region). Although the Upper Klamath relies on groundwater supplies for only 26 percent
 33 for meeting their overall water uses, 66 percent of the urban water use in the Upper Klamath is met by
 34 groundwater. Coastal PA and Russian River PA provides an average annual groundwater supply equal to
 35 82 taf and 75 taf providing respectively, 63 percent and 52 percent of the overall water supply water uses
 36 and meeting 78 percent and 62 percent of the agricultural water uses in the two PAs.

37 *More detailed information regarding groundwater water supply and use analysis is available online from*
 38 *Update 2013 Volume 4, Reference Guide — California’s Groundwater Update 2013.*

1 Changes in annual groundwater supply and type of use may be related to a number of factors, such as
 2 changes in surface water availability, urban and agricultural growth, market fluctuations, and water use
 3 efficiency practices.

4 Figures NC-15 and 16 summarize the 2002 through 2010 groundwater supply trends for the North Coast
 5 Hydrologic Region. The right side of Figure NC-15 illustrates the annual amount of groundwater versus
 6 surface water supply, while the left side identifies the percent of the overall water supply provided by
 7 groundwater relative to surface water. The center column in the figure identifies the water year along with
 8 the corresponding amount of precipitation, as a percentage of the 30-year running average for the region.
 9 Figure NC-16 shows the annual amount and percentage of groundwater supply trends for meeting urban,
 10 agricultural, and managed wetland uses.

11 **PLACEHOLDER Figure NC-15 North Coast Hydrologic Region Annual Groundwater Water Supply**
 12 **Trend (2002-2010)**

13 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 14 the end of the report.]

15 **PLACEHOLDER Figure NC-16 North Coast Hydrologic Region Annual Groundwater Supply Trend**
 16 **by Type of Use (2002-2010)**

17 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 18 the end of the report.]

19 Figure NC-15 indicates that the annual water supply for the region has remained relatively stable between
 20 2002 and 2010, which is likely due to relatively stable climatic conditions and surface water supply for
 21 the region. The driest year for the region was 2009, with precipitation equal to 76 percent of the 30-year
 22 average. The wettest year was 2006, with precipitation equal to 135 percent of the 30-year average.
 23 Between 2002 and 2010, annual groundwater supply fluctuated from about 300 taf in 2005 to about
 24 400 taf in 2007 and 2008 and provided between 34 and 36 percent of the total water supply for the region.
 25 Figure NC-14 indicates that groundwater supply meeting agricultural use ranged from 78 to 85 percent of
 26 the annual groundwater extraction, with the remaining groundwater extraction meeting mostly relatively
 27 stable urban use. Groundwater remained a minor supply at 1 to 2 percent for meeting managed wetland
 28 use.

29 *Reclaimed Water*

30 The City of Santa Rosa, the City of Arcata, and the Town of Windsor are using reclaimed water for
 31 landscape irrigation and holding tanks for fire suppression. On a regional scale, the North Coast
 32 RWQCB's Basin Plan recommends recycling portions of urban and agricultural water to help meet water
 33 demands for quality and supply.

34 *Geysers Recharge Project*

35 The Santa Rosa Sub-regional Reclamation System reclaims water, treats it to a tertiary level, and
 36 distributes it to agricultural users, golf courses, public and private landscaping, and The Geysers
 37 steamfield. Santa Rosa's reclamation system is one of the largest reclaimed water agricultural irrigation
 38 systems in the country. For the Geysers Recharge Project, reclaimed water is piped through a 42-mile
 39 pipeline and injected into underground wells in The Geysers steamfield in Sonoma and Lake counties.

1 Once within the wells, the water is gradually heated by geothermal activity to produce a steam that is then
 2 utilized to produce electricity at nearby power plants. The Geysers Recharge Project was chosen as a
 3 means to dispose of treated wastewater during the winter months, when there is no demand for
 4 agricultural irrigation. The Sub-regional Reclamation System had previously been discharging the unused
 5 water to the Russian River, but stricter water quality regulations removed this option. The Sub-regional
 6 Reclamation System is currently exploring other means of reusing or disposing of current and future
 7 amounts of reclaimed water in order to best manage water resources.

8 In November 2003, the Geysers Recharge Project began pumping 11 mgd of highly treated wastewater
 9 from the Laguna Treatment Plant to The Geysers steamfields, high in the Mayacamas Mountains. In
 10 January 2008, the delivery was up to 12.62 mgd helping to generate enough electricity for 100,000
 11 households in Sonoma and other North Bay counties.

12 The Geysers Expansion Project builds on the Geysers Recharge Project and will increase recycled water
 13 deliveries to the Geysers steamfield up to 19.8 mgd or as much as an additional 3,209 million gallons per
 14 year. Santa Rosa has completed negotiations with Calpine, the steamfield operator, and has signed a
 15 contract to send more water to The Geysers.

16 *Imported / Exported Water*

17 The North Coast region does not import water, but water transfers do occur within the region. For
 18 example, Eel River water is diverted at the Van Arsdale Dam into the Russian River (Potter Valley
 19 Project). The North Coast generally exports more water to other regions than the volume of water
 20 consumed within the region for agricultural and urban uses. Two out-of-region transfers include the
 21 CVP's TRD and the north San Francisco Bay Area (Petaluma Aqueduct). See "Project Operations"
 22 section of this document for additional information.

23 **Water Uses**

24 The principal developed uses of environmental water occur in the Lower Klamath Lake, Tule Lake and
 25 Clear Lake National Wildlife refuges, and the Butte Valley and Shasta Valley wildlife areas. In Butte
 26 Valley, most of the water for wildlife comes from about 3,000 af of groundwater. As a result of the
 27 passage of both federal and State Wild and Scenic Rivers Acts in 1968 and 1972, many of the major
 28 rivers in the North Coast region have been preserved to maintain their free-flowing character and provide
 29 for environmental uses. Most of the Eel, Klamath, Trinity, and Smith rivers are designated as wild and
 30 scenic, which preserves these river resources and protects them from new water development. On the
 31 Trinity River, efforts to restore the fishery led to a federal Record of Decision (ROD) in year 2000 to
 32 increase the fishery flow releases from Trinity Lake. After several years of legal challenges, this decision
 33 was upheld by a July 2004 federal court decision. The water allocated to downstream fishery flows is now
 34 being increased from the previous 340,000 af /yr, to a new schedule that ranges between 368,600 af in a
 35 critically dry year to more than 700,000 af per year in a wet water year. Biologists and CVP operators are
 36 still working on the development of daily, weekly, and monthly water-release schedules that will make
 37 the best use of these new water allocations.

38 *Drinking Water*

39 The region has an estimated 262 community drinking water systems. The majority (over 85 percent) of
 40 these community drinking water systems are considered small (serving fewer than 3,300 people) with
 41 most small water systems serving fewer than 500 people (Table NC-10 Summary of Community Water

1 Systems). Small water systems face unique financial and operational challenges in providing safe
 2 drinking water. Given their small customer base, many small water systems cannot develop or access the
 3 technical, managerial, and financial resources needed to comply with new and existing regulations. These
 4 water systems may be geographically isolated, and their staff often lacks the time or expertise to make
 5 needed infrastructure repairs; install or operate treatment; or develop comprehensive source water
 6 protection plans, financial plans or asset management plans (U.S. Environmental Protection
 7 Agency 2012).

8 **PLACEHOLDER Table NC-10 Summary of Community Water Systems within the North Coast**
 9 **Hydrologic Region**

10 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 11 the end of the report.]

12 In contrast, medium and large water systems account for less than 15 percent of region’s drinking water
 13 systems; however, these systems deliver drinking water to more than 80 percent of the region’s
 14 population (see Table NC-10). These water systems generally have financial resources to hire staff to
 15 oversee daily operations and maintenance needs and to hire staff for planning future infrastructure
 16 replacement and capital improvements. This helps to ensure that existing and future drinking water
 17 standards can be met.

18 *Agricultural Water Use*

19 **Annual Reference ET rates (Spatial ETo) for Selected Locations**

20 Following are the annual reference evapotranspiration rates from California Irrigation Management
 21 Information System (CIMIS):

22	Smith River (Del Norte County)	42.36 inches
23	Fortuna (Humboldt County)	44.58 inches
24	Ukiah (Mendocino County)	43.64 inches
25	Santa Rosa (Sonoma County)	40.24 inches
26	Etna (Siskiyou County, Scott Valley)	44.62 inches
27	Montague (Siskiyou County, Shasta Valley)	44.19 inches
28	MacDoel (Siskiyou County, Butte Valley)	43.50 inches
29	Tule Lake (Modoc and Siskiyou counties)	42.99 inches

30 Values estimated by CIMIS and Spatial CIMIS do not account for rainfall, light rain (trace), fog, or dew
 31 formation. These values are site-specific and require direct observation by those applying the information.
 32 Rainfall entering the crop-soil profile (effective precipitation) can be subtracted from the water use
 33 demand on a daily basis. Effective precipitation (EP) is the amount of rainfall actually entering the soil
 34 and available to the plant, not running off as surface water or percolating through the soil beyond the root

1 zone. For additional information on EP or rainfall, see web page at
2 <http://www.fao.org/docrep/S2022E/s2022e03.htm>. Light rain, fog, and dew contribute to lowering the
3 crop water demand by lowering the temperature and increasing the humidity in the micro-environment of
4 the plant. When present, trace precipitation, fog, and dew only form for short time periods requiring
5 frequent observation and good record-keeping. This is most important along the coast as fog and dew in
6 these areas can contribute a great deal to meeting the water use demands of the crop. For more
7 information on light rain, fog, and dew accounting for crop water use demand, see Correcting soil water
8 balance calculations for dew, fog, and light rainfall by R. Moratiel, D. Spano, P. Nicolosi and R.L.
9 Snyder, Irrigation Science paper: DOI 10.1007/s00271-011-0320-2 (California Department of Water
10 Resources 2009).

11 **Scott and Shasta Valley Study on Alfalfa Water Use**

12 Blaine Hanson, Extension Specialist (Emeritus), Land, Air and Water Resources (University of
13 California, Davis), in cooperation with Steve Orloff, Siskiyou County Farm Advisor (University of
14 California Cooperative Extension), et.al., have been working on a study of alfalfa water use in California
15 (including Scott and Shasta Valleys) from 2007 through 2010. The intention of the study is to develop
16 new crop water-use values for alfalfa to be used by agriculture and planning and to compare these
17 findings to historical text book assumptions. As of the writing of this document, study results are not yet
18 published. However, preliminary results indicate that historical, seasonal ET rates for alfalfa in California
19 have been overestimated, with the amount dependent on where the crop is grown and the type of soil.
20 Observed seasonal alfalfa water use from this study for the Scott and Shasta valleys ranged from 32.8 to
21 39.6 inches whereas historical estimates ranged from 36.5 to 44.0 inches. The average seasonal difference
22 between these two methods yielded a 3.25-inch over-application when using the historical values
23 compared to the observed amounts. The median difference between the two methods is 2.25 inches for the
24 season. As an example, if one were to apply an additional 3.25 inches of water over a typical 160-acre
25 field for the season, the additional water necessary to apply would amount to 520 acre-inches or 43.3 af of
26 additional water. This would be additional water required to meet expected seasonal crop demand if using
27 the historical values. Furthermore, this additional water would need to be diverted or pumped during the
28 irrigation season in order to meet the expected demand, which would require additional expense. (This
29 does not count the water necessary to overcome the irrigation system efficiency and assumes a good
30 uniformity of application.)

31 **New Vineyard Irrigation Practices**

32 The new vineyard installations use the latest technologies ensuring the optimum use of resources.
33 However, non-point-source pollution from vineyards, including pesticides, is still a concern. Current
34 cultural practices recommended by UC Cooperative Extension include minimum tillage to prevent soil
35 transport and minimum applications of fertilizer and pesticide at an agronomically proper rate. The goal
36 of these recommendations is to minimize the impact agricultural (vineyard) management has on the
37 environment. Although most vineyards with microspray and drip irrigation systems do not have much
38 runoff, agricultural tail water return systems and settling basins for runoff help to conserve and protect
39 water supplies.

40 ***Water Conservation Act of 2009 (SB x7-7) Implementation Status***

41 Thirteen North Coast urban water suppliers have submitted 2010 urban water management plans to DWR.
42 The Water Conservation Law of 2009 (SBx7-7) required urban water suppliers to calculate baseline water
43 use and set 2015 and 2020 water use targets. The urban water management plans indicate the North Coast

1 Hydrologic Region had a population-weighted baseline average water use of 147 gallons per capita per
 2 day with an average population-weighted 2020 target of 127 gallons per capita per day. The [Baseline and](#)
 3 [Target Data](#) for the North Coast urban water suppliers is available on the DWR Urban Water Use
 4 Efficiency Web site.

5 The Water Conservation Law of 2009 (SBx7-7) required agricultural water suppliers who supply more
 6 than 25,000 irrigated acres to prepare and adopt agricultural water management plans by December 31,
 7 2012, and update those plans by December 31, 2015, and every 5 years thereafter. All of the North Coast
 8 agricultural water suppliers supply fewer than 25,000 irrigated acres; as of August 2013, no agricultural
 9 water management plan had been submitted from the North Coast region.

10 **Water Balance Summaries**

11 Water balance figure and the narrative discussion below provide a detailed summary of the actual
 12 regionwide water supplies and water uses from years 2001 through 2010 for the entire North Coast
 13 region. Figure NC-17 summarizes the dedicated and developed urban, agricultural, and environmental
 14 water uses in the region for 2001 thru 2010. The figure also provides a graphical presentation of all of the
 15 water supply sources that are used to meet the developed water uses within this hydrologic region for
 16 these years. As shown on the first graph, the volume of water dedicated to wild and scenic rivers, called
 17 “statutory required outflows,” is the largest component of dedicated water uses in the region. The
 18 information presented in Table NC-11 also indicates that the volume of water exported to other regions is
 19 generally greater than all the water consumptively used for urban, agriculture and wildlife refuges within
 20 the North Coast region.

21 **PLACEHOLDER Figure NC-17 North Coast Water Balance by Water Year, 2001–2010**

22 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 23 the end of the report.]

24 **PLACEHOLDER Table NC-11 North Coast Hydrologic Water Balance Summary, 2001-2010** 25 **(Thousand af)**

26 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 27 the end of the report.]

28 The water balance information for North Coast Hydrologic Region is summarized by planning area. In
 29 this region, the four PAs are Upper Klamath (PA 101), Lower Klamath (PA 102), Coastal (PA 103), and
 30 Russian River (PA 104).

31 Upper Klamath PA is primarily agricultural in nature, with water use ranging from about 615 to 680 taf
 32 /yr, in comparison to about 10 to 12 taf urban use. This PA also supports considerable managed wetlands,
 33 which consumed about 225 taf water in 2010. This value is less than was applied in previous years, in
 34 which as much as 400 taf was used. There are no instream or wild and scenic designated rivers in this PA.
 35 Surface water supplies are split more or less equally between local supplies and the local federal project,
 36 at about 250 taf each. This supply is augmented with groundwater and reuse. Groundwater extraction
 37 averages about 180 to 200 taf, with about 70 to 75 taf being recharged back into the aquifer. Between
 38 2006 and 2010, 125 to 156 taf of applied water was reused. Previously stored water (22 to 25 taf) was
 39 added to make up the difference between applied water and supplies.

1 In the Lower Klamath PA, the primary application of water is for instream environmental uses. Instream
 2 and Wild and Scenic Requirements in these rivers of account for about 10 maf to 22 maf of applied water.
 3 The urban use is about the same as in the Upper Klamath PA. Agricultural applied water equals 20 taf to
 4 40 taf, and there are no managed wetlands in this PA. As can be expected, most of the water comes from
 5 local sources, with about 4 taf to 7 taf from groundwater and another 2 taf from reuse.

6 In the Coastal PA, there are still a few wild and scenic rivers, which account for most of the applied
 7 water, ranging between 3 maf to 11 maf. Instream use is next in volume with about 100 taf. Urban use is
 8 greater than either of the more northern PAs, at about 50 taf; and agricultural applied water ranges from
 9 about 55 taf to 100 taf. There are a few managed wetlands in the PA, using about a thousand ace-feet
 10 total. Most of the water supply is from local sources, with an additional 100 taf of groundwater being
 11 pumped. 30 taf are recharged back into the aquifer. There has been an increase to about 30 taf /yr in reuse
 12 during 2008 to 2010.

13 The Russian River PA is the most urbanized area of the North Coast Hydrologic Region. Urban applied
 14 water ranges from 80 to 94 taf. There is about 100 to 125 taf for agricultural uses and 90 to 100 taf from
 15 instream. There are no wild and scenic rivers or managed wetlands in this PA. Local supplies account for
 16 about 100 to 133 taf. Local imports have been decreasing from about 40 taf in 2006 to none in 2010.
 17 About 75 taf of groundwater supplements this, with about 15 to 25 taf recharged back into the aquifer.
 18 The instream flows are reused as part of the local supply. There is also some recycled water in this
 19 planning area, which varies from 12 taf to zero, depending on the year and other conditions.

20 The water portfolios are estimates of present water balances of water uses and supplies for each region in
 21 California. The water portfolios are aggregated to spatial scales unique to the California Water Plan
 22 (CWP) including the detailed analysis unit, planning area, and hydrologic region. Technical
 23 enhancements will allow this information to be evaluated at boundaries used by water purveyors and
 24 regional water management groups. A significant part of this work is to transition from the existing
 25 spreadsheet-based data storage of the water portfolio information to an enterprise data management
 26 system that will facilitate sharing of information through the Internet. Additional enhancements are under
 27 way to describe the hydrologic cycle components more fully within the water portfolios — groundwater
 28 in particular.

29 **Project Operations**

30 *Potter Valley Project*

31 The northern edge of Potter Valley in Lake County separates the Russian River watershed from the Eel
 32 River watershed, and in the year 1900 it was an ideal place to build a hydroelectric power plant. The
 33 Potter Valley Project was first licensed as a hydroelectric power plant in 1922 by the Federal Power
 34 Commission. The current license expires on April 14, 2022. See "Potter Valley Project FERC License"
 35 under "Water Governance" in this report. Annual flows in the Eel River are quite variable. In the
 36 relatively dry year of 2009, the peak flow in the beginning of March — as measured passing Cape Horn
 37 Dam at gage E-11 (downstream of the diversion) — for one day was over 5,000 cubic feet per second,
 38 quickly dropping to approximately 1,000 cfs and then back to the winter steady state of around 150 cfs
 39 before the next major rain. Peak winter flows can occasionally exceed 100,000 cfs. These winter storm
 40 events are captured and stored behind Scott Dam (Lake Pillsbury) for later use. Per a 2006 bathymetric
 41 survey, the maximum storage in Lake Pillsbury is 74,993 acre feet. From spring until fall, on an average

1 rainfall year, approximately 125 cfs is diverted through the Potter Valley Project into the Russian River
2 watershed. (Potter Valley Irrigation District 2010).

3 *Coyote Valley Dam and Lake Mendocino*

4 Lake Mendocino is located on the East Fork of the Russian River (downstream of the Potter Valley
5 Project), about 5 miles northeast of Ukiah in Mendocino County. The Coyote Dam (also known as
6 Coyote Valley Dam) project was authorized by the Flood Control Act of 1944 and completed in 1958 for
7 purposes of flood control, water supply, recreation, and streamflow regulation. Lake Mendocino has a
8 flood storage capacity of 122,400 af and a total surface area of 1,822 acres. The lake has an un-gated
9 spillway, designed for a maximum release of 35,800 cfs. Major facilities include an anadromous
10 endangered/protected fish species egg collection and imprinting facility, visitor cultural center complex,
11 park headquarters, sponsor run electrical power plant (hydropower), developed campgrounds (300 sites),
12 18 primitive boat-in/hike-in campsites, a trail system, 2 boat launch ramps, swim beach, and picnic areas.
13 Of the park's 5,110 acres, 689 are devoted to wildlife management (U.S. Army Corps of Engineers,
14 Coyote Valley Dam 2010).

15 *Warm Springs Dam and Lake Sonoma*

16 Warm Springs Dam and Lake Sonoma is located on Dry Creek in Sonoma County, approximately
17 14 miles above the confluence with the Russian River. The project is located on 15,966 acres of land,
18 situated approximately 14 miles northwest of Healdsburg.

19 Warm Springs Dam forms Lake Sonoma, which has a design capacity of 381,000 af and drains an area of
20 approximately 130 square miles, or about 9 percent of the total Russian River basin. Construction started
21 in 1967 and was completed in 1982. The dam is operated and maintained by USACE. The storage space
22 for water conservation is owned by the Sonoma County Water Agency (SCWA), while the remaining part
23 of the project is owned by USACE, which directs flood control releases from Warm Springs Dam.

24 The Don Clause Fish Hatchery (Warm Springs Fish Hatchery) is located on Dry Creek at the base of
25 Warm Springs Dam. This facility is operated by California Fish and Wildlife (DFW, formerly Department
26 of Fish and Game) under a cooperative agreement with USACE. The hatchery was created as part of the
27 Warm Springs Dam Project to compensate for loss of spawning and rearing habitat that was impounded
28 and made inaccessible to anadromous fish by the dam.

29 SCWA owns and operates the Warm Springs Dam hydroelectric facility. The hydroelectric facility was
30 completed in December 1988. SCWA operates the facility under a 50-year license issued by the Federal
31 Energy Regulatory Commission (FERC) on December 18, 1984. The 3,000-kilowatt Francis turbine
32 generators have a power rating of 2.6 megawatt (U.S. Army Corps of Engineers, Warm Springs Dam
33 2010).

34 *Petaluma Aqueduct*

35 SCWA owns and maintains a series of underground pipes that run from water collectors at Wohler Bridge
36 near Forestville on the Russian River to northern Marin County: the Petaluma Aqueduct serves the greater
37 Santa Rosa area, the City of Petaluma, and North Marin Water District. (See "North Marin Aqueduct"
38 subsection below.) In 1960, Petaluma leaders signed an agreement to receive water from SCWA. The
39 Petaluma Aqueduct was completed in December 1961. As a contractor of SCWA, Petaluma agrees to
40 purchase water at guaranteed rates while SCWA handles the two-county distribution system.

1 The Petaluma Aqueduct carries more than 90 percent of the water used by the City of Petaluma, over
2 8 mgd. Unfortunately, the underground structure is 50 years old, has exceeded its predicted lifespan, and
3 could rupture during an earthquake. While the Petaluma Aqueduct itself warrants monitoring and study,
4 Petaluma is only one user on an 85-mile system of water transmission lines. With Petaluma at the south
5 end of the system, any breakdown along the conveyance affects everyone downstream. SCWA staff
6 recalled times over the years when the water agency had to shut down its system. By implementing
7 conservation measures and using water held in storage, Petaluma was able to manage the temporary loss
8 of its primary supply. During times of supply curtailment, SCWA has 2- or 3-day supply in storage along
9 the aqueduct, and the City of Petaluma has a couple of days of storage and groundwater wells.

10 Typically, Petaluma's own source of municipal water only comprises 2 percent of the city's water use, but
11 in recent drought years, local wells were run more often and made up 10 percent of the city's average
12 water use (Petaluma 360 2012). Like most Sonoma County cities, Petaluma drew its own water from
13 municipal wells for decades. According to DWR, the original water source for the community was the
14 headwaters of Adobe Creek.

15 *North Marin Aqueduct*

16 The North Marin Aqueduct is an extension of the Petaluma Aqueduct to supply water to North Marin
17 Water District and Marin Municipal Water District for the city of Novato and surrounding communities.

18 Russian River water, which provides about 80 percent of Novato's water demand, originates in
19 Mendocino County from both the Eel River and the Russian River watershed. Eel River water flows from
20 the Potter Valley Project diversion on the Eel River to the east fork of the Russian River. Then,
21 downstream at a point about 10 miles upstream of Guerneville, near Forestville, water is collected by five
22 Ranney water collectors. This water is then pumped directly into the Petaluma Aqueduct system to supply
23 treatable water for potable use to a two-county area.

24 Stafford Lake, which provides approximately 20 percent of Novato's water demand, lies 4 miles west of
25 downtown Novato and collects runoff from 8.3 square miles of watershed property located upstream at
26 the upper tributary reaches of Novato Creek.

27 Since 2007, the Deer Island Recycled Water Facility near Novato, located adjacent to Highway 37, has
28 produced treated recycled water supplies to offset Russian River water and help improve Novato's water
29 supply for large landscape and fire protection (North Marin Water District 2013a, 2013b).

30 *R.W. Matthews Dam, Ruth Lake and Mad River*

31 R.W. Matthews Dam forms Ruth Lake in southern Trinity County. It impounds runoff from the upper
32 quarter of the Mad River Basin, an area of approximately 121 square miles. The lake capacity is
33 48,030 af.

34 A portion of the water stored in Ruth Lake is released each summer and fall to satisfy the Humboldt Bay
35 Municipal Water District's (HBMWD) downstream diversion requirements, as well as maintain minimum
36 bypass flow requirements in the Mad River below Essex. Although the HBMWD impounds water at Ruth
37 Lake and diverts water at Essex, the operations do not significantly affect the natural flow regime in the
38 Mad River. (Essex is located on the Mad River 3.5 miles northeast of Arcata at an elevation of 75 feet.)

1 The total volume of water impounded and diverted by HBMWD represents a small percentage of the
 2 natural yield of the Mad River watershed. The Mad River’s average annual discharge into the Pacific
 3 Ocean is just over 1 maf. Ruth Lake, in its entirety, represents less than 5 percent of the total average
 4 annual runoff from the Mad River basin. The entire 48,030-acre-foot capacity of Ruth Lake is not drawn
 5 down each year so the amount of winter-season runoff captured in the reservoir is yet a smaller
 6 percentage of the total runoff. With respect to diversions, the current withdrawal rate at Essex is
 7 approximately 25 mgd to 30 mgd (28,000 to 34,000 af /yr), which is 3 percent of the total annual average
 8 runoff of the Mad River watershed. The full diversion capacity of 75 mgd (84,000 af /yr) is 8 percent of
 9 the total annual average runoff of the watershed.

10 Tributaries downstream of Matthews Dam contribute significantly to, and are a major influence on,
 11 resulting flow rates in the Mad River. A former USGS gage station near Forest Glen was located 9 miles
 12 below the dam prior to the confluence of any major tributaries. Annual mean flow at the Forest Glen gage
 13 station increased by an average of 22 percent compared to the mean flows just below Ruth Lake. The
 14 more significant tributaries on the Mad River are located downstream of this former gage station. These
 15 tributaries contribute significantly to Mad River discharge and also provide a “buffering effect” during the
 16 few times the HBMWD is releasing from Ruth Lake less than the natural flow (e.g. during the first winter
 17 storms).

18 There is no out-of-basin transfer in the upper watershed, as occurs on some river systems. The water
 19 released by HBMWD flows down the main stem Mad River channel and augments flows, which would
 20 not occur naturally during the summer and fall. Flow augmentation has many beneficial effects, including
 21 expanding river habitat for the benefit of aquatic species and improving water quality in the summer and
 22 fall (Humboldt Bay Municipal Water District 2012).

23 *Iron Gate Dam and Klamath River*

24 Iron Gate Dam (Iron Gate Reservoir) is operated within the constraints of the Klamath Basin Operations
 25 Plan. The plan for the USBR’s Klamath Project, which is located within the upper Klamath River Basin
 26 in southern Oregon and northern California, describes project operations on an annual basis from April 1
 27 of one year through March 31 of the next, based upon current and expected hydrologic conditions.

28 Reclamation develops this plan annually to serve as a planning aid for agricultural water users, Klamath
 29 Basin tribes, national wildlife refuges, and other interested parties. The plan provides an estimated project
 30 water supply to the following areas:

- 31 • West Side delivery area: This area includes lands in southern Oregon and northern California
 32 that receive project water primarily from Upper Klamath Lake (UKL) and/or the Klamath
 33 River. This area also includes the Tule Lake and Lower Klamath National Wildlife Refuges.
- 34 • East Side delivery area: This area includes lands within Langell Valley Irrigation District and
 35 Horsefly Irrigation District (both in Oregon) on the east side of the project area. This area
 36 receives water from Clear Lake Reservoir (California), Gerber Reservoir (Oregon), and the
 37 Lost River (California and Oregon).

38 In response to both the 2010 National Marine Fisheries Service (NMFS) biological opinion (BO), and the
 39 2008 U.S. Fish and Wildlife Service (USFWS) BO, the USBR developed a Variable Base Flow (VBF)
 40 procedure to be used for operations. The VBF procedure was developed based on the following

1 objectives: (1) provide certainty in compliance with the UKL minimum elevations, as outlined in
 2 Table 2-1 of the 2008 USFWS BO and (2) provide a procedure that tracks the flows outlined in
 3 Table 18 of the 2010 NMFS BO and Reasonable and Prudent Alternatives. These objectives were
 4 designed to help meet the needs of coho salmon during critical periods of the year. For more information
 5 on the Klamath Basin Operations Plan, refer to the reference material listed at the end of this chapter
 6 (National Marine Fisheries Service 2010; U.S. Bureau of Reclamation 2012; U.S Department of Interior
 7 2008).

8 **Iron Gate, Copco, and Dwinnell Reservoirs**

9 Iron Gate and Copco reservoirs are operated for hydropower, water supply, and recreation. The Copco
 10 Reservoirs are located in the northern portion of Shasta Valley, upstream of Iron Gate Reservoir, and
 11 Dwinnell Reservoir is located in the southern portion of the valley. All of the reservoirs are important to
 12 the residences surrounding the lakes. Dwinnell reservoir (Lake Shastina) is for municipal water for the
 13 city of Montague, irrigation supply for the Montague Irrigation District, and recreation.

14 *Trinity Dam and Exports from Trinity River to Central Valley*

15 Trinity Dam stores water from the Trinity River in the Trinity Reservoir (Trinity Lake, formerly Clair
 16 Engle Lake). Water that is released from Trinity Dam is regulated by Lewiston Dam (directly
 17 downstream), which provides a forebay for diversion flows to the Clear Creek Tunnel. From the Clear
 18 Creek Tunnel, water then enters Whiskeytown Lake through Judge Francis Carr Powerhouse. Some of the
 19 water diverted from Whiskeytown Lake flows into the Clear Creek Unit South Main Aqueduct to irrigate
 20 lands in the Clear Creek Unit. The rest flows through the Spring Creek Power Conduit and Power Plant
 21 into Keswick Reservoir. From there, water goes through Keswick Power Plant to the Sacramento River.
 22 Exports from the TRD contribute to meeting minimum flow requirements in the Trinity and Sacramento
 23 rivers, help to maintain reservoir storage levels, and facilitate other CVP operating requirements such as
 24 compliance with the winter-run BO, which requires that certain temperature requirements be met in the
 25 Sacramento River below Keswick Dam.

26 Prior to construction of the TRD, average annual discharge at Lewiston was approximately 1.2 maf. Peak
 27 flows in excess of 100,000 cfs were recorded at the town of Lewiston, and daily average flows greater
 28 than 70,000 cfs occurred three times between 1912 and 1963. Following construction, instream flow
 29 releases were set at 120,500 ace-feet /yr (10 percent of the average unimpaired inflow). From 1964 to
 30 1996, TRD exports accounted for 14 percent of Keswick releases. In the period of 1986 through 1996, the
 31 TRD exports accounted for 12 percent of Keswick releases.

32 An outcome of TRD operations and the reduced instream Trinity River flows was degraded fish habitat
 33 and drastic reductions in anadromous fish populations. By 1980, it was estimated that fish populations
 34 had been reduced by 60 to 80 percent due to inadequately regulated harvest, excessive streambed
 35 sedimentation, and insufficient streamflows. The loss of fishery habitat was estimated to be 80 to
 36 90 percent. To help address these problems, Congress passed the Trinity River Stream Rectification Act
 37 in 1980 (addressing sedimentation issues) and passed the Trinity River Basin Fish and Wildlife
 38 Management Act in 1984. The 1984 act directed efforts to restore fish and wildlife populations to levels
 39 that existed prior to TRD construction.

40 One of the provisions of the 1992 Central Valley Project Improvement Act was the establishment of a
 41 minimum flow volume of 340,000 af for the Trinity River. The CVPIA also directed the completion of

1 the 12-year study (Trinity River Flow Evaluation Study) to establish permanent instream fishery flow
 2 requirements, TRD operating criteria, and procedures for restoration and maintenance of the fishery. The
 3 TRFES report recommended specific annual flow released, sediment management, and channel
 4 rehabilitation to provide necessary habitat. The subsequent Trinity River Environmental Impact Statement
 5 (EIS)/Environmental Impact Report (EIR) and ROD in 2000 identified the annual water allocation for
 6 specific water year types (see the section below for more information about the Trinity River Restoration
 7 Program Restoration Flows).

8 *USBR Flow Releases to the Trinity and Sacramento Rivers*

9 USBR releases to the Trinity and Sacramento rivers include two types of releases, namely, Safety of
 10 Dams and Other releases.

- 11 • Safety of Dams: During the winter, USBR maintains lower levels in Trinity Lake to provide a
 12 buffer in the event of an extremely large winter storm. The quantity of that buffer is based on
 13 several factors and primarily references many years of hydrologic record for the basin.
 14 Maintaining storage space is a very important aspect of flood control operations and is
 15 fundamental in protecting areas downstream of Trinity Dam, as well as the dam itself. As
 16 winter storms fill Trinity Lake, USBR may need to increase releases to maintain the lower lake
 17 levels. Because these elevated winter releases help protect the dam, they are commonly called
 18 “Safety of Dams releases” and may or may not occur in conjunction with actual winter storms.
 19 These releases are made independently from the ROD releases for river restoration. Safety of
 20 Dams releases are scheduled by USBR in response to current conditions and typically have no
 21 more advance warning than a few days. USBR uses a combination of increased releases to the
 22 Trinity River through Lewiston Dam and trans-basin diversions to the Sacramento River
 23 through the Clear Creek Tunnel to lower the water level in Trinity Lake (see
 24 <http://www.trrp.net/background/ops/>). Consequently, releases from Trinity Dam to Lewiston
 25 Reservoir may be higher than releases from Lewiston Dam to the Trinity River. Safety of Dams
 26 releases from Lewiston Dam to the Trinity River are typically no greater than 6,000 cfs, but
 27 may go higher if conditions warrant.
- 28 • Other Releases: USBR occasionally makes flow releases from Lewiston Dam to the Trinity
 29 River for other purposes such as tribal releases or to mitigate late summer conditions in the
 30 Lower Klamath River for fish health purposes. USBR coordinates these releases with the
 31 Trinity River Restoration Program and usually provides several weeks’ public notice. Such
 32 releases are independent from the ROD releases for river restoration (Trinity River Restoration
 33 Program 2012; U.S. Department of Interior 2000).

34 **Trinity River Restoration Program — Restoration Flows**

35 The ROD (ROD, U.S. Department of Interior, 2000) directs USBR to provide annual in-stream flows
 36 below Lewiston Dam deemed necessary to restore and maintain the Trinity River’s fishery resources.
 37 These restoration flows link two essential purposes: (1) flows to provide physical fish habitat (i.e.
 38 appropriate depths and velocities and suitable temperature regimes for anadromous salmonids); and (2)
 39 flows to restore the riverine processes that create and maintain the structural integrity and spatial
 40 complexity of the fish habitats. The ROD provides recommended daily release schedules for each of the
 41 five water year types (critical dry to extremely wet — Table NC-12). The ROD stipulates that “the daily
 42 schedule for releasing water for a given water year may be adjusted based on monitoring and studies
 43 guided by the Trinity Management Council but the associated annual water volume allocation may not be
 44

1 changed.” A water year is the 12-month period from any October 1 through September 30 of the
 2 following calendar year. The 2012 water year extends from October 1, 2011, to September 30, 2012.

3 **PLACEHOLDER Table NC-12 Trinity River ROD Water Year Types**

4 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 5 the end of the report.]

6 The predicted water year type is based on the April 1 forecast for the annual river runoff of the Trinity
 7 River at Lewiston, California. The annual runoff forecast is jointly developed by the National Weather
 8 Service and DWR for the entire state of California, including the Trinity River. Identical forecasts are
 9 published in the “Water Supply Outlook for California and Northern Nevada”
 10 (http://www.cnrfc.noaa.gov/water_supply.php) produced by the National Weather Service and in
 11 “Bulletin 120 Water Conditions in California” produced by DWR.

12 The finalized water-year forecast determines the water year type (e.g. wet, dry), and the ROD describes
 13 the volume of water available to the Trinity River Restoration Program for restoration releases for the
 14 different water year types. The Trinity River Restoration Program develops annual flow release
 15 recommendations through a collaborative process to meet ROD objectives for specific water year types.
 16 Trinity River Release and Diversion Summary since 2000 is listed in Table NC-13.

17 **PLACEHOLDER Table NC-13 Trinity River Release and Diversion Summary Since 2000**

18 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 19 the end of the report.]

20 The Trinity Management Council makes the final flow recommendation that is then forwarded to the U.S.
 21 Department of Interior for consideration (Trinity River Restoration Program 2012; U.S. Department of
 22 Interior 2000).

23 **Gravel Mining**

24 Historical gravel mining along many of the North Coast rivers and streams has presented a particular
 25 problem concerning sediment transport. Many (if not all) of the waterways have been affected by silt and
 26 clay deposition causing a negative impact on local and regional fish spawning areas. Several major gravel
 27 mining operations along the Russian River have been curtailed in recent years. Improvements such as
 28 settling basins have been implemented to control the amount of sediment outflow from these mining areas
 29 to help improve downstream water quality. The issuance of 401 water quality certifications is the primary
 30 mechanism for regulating water quality impacts from instream gravel mining. Some of the counties in this
 31 region (Humboldt and Sonoma) have gravel regulation programs in place that also play a significant role.

32 *Statewide Instream Mining (Suction Dredge Mining)*

33 Instream mining (specifically, suction dredge mining) has been curtailed in California as of 2008 with no
 34 set ending date on the moratorium. The Legislature and Governor have enacted Senate Bill 1018 (2012).
 35 A part of this legislation applies to suction dredge mining. Suction dredging, including the method known
 36 as “booming,” is prohibited within 100 yards of any California river, stream or lake (Fish & Game Code,
 37 § 5653 subd. (d)).

1 The current moratorium originally established by SB 670 and extended by Assembly Bill 120 and SB
2 1018 does not prohibit or restrict nonmotorized recreational mining activities, including panning for gold.
3 It also does not prohibit or restrict some other forms of mining, including, for example, practices known
4 as high banking, power sluicing, sniping, or using a gravity dredge so long as gravel and earthen materials
5 are not vacuumed with a motorized system from the river or stream. It is important to know that other
6 environmental laws may apply to some of these mining practices. In addition, these activities may be
7 subject to the authority of the appropriate RWQCB.

8 Small-scale suction dredge mining activity in California began in the 1960s and peaked during high gold
9 prices in the late 1970s and early 1980s. The existing regulatory framework governing the activity as
10 administered by DFW is rooted in statutory amendments to the Fish and Game Code that took affect
11 originally in the late 1980s. Under the statute and regulations, any California resident or nonresident
12 could (i.e., before the current moratorium) obtain a suction dredge mining permit from the DFW upon
13 payment of a fee required by statute. On average, DFW issued approximately 3,200 suction dredge
14 mining permits a year to California residents, and another 450 a year to nonresidents, from
15 1995 through 2009.

16 DFW's recent effort to amend the regulations and comply with the California Environmental Quality Act
17 (CEQA) was required by a court order issued in a lawsuit brought against DFW by the Karuk Tribe of
18 California. The lawsuit focused on the Klamath, Scott, and Salmon River watersheds in Northern
19 California; included allegations regarding impacts to various fish species, including coho salmon; and
20 contended that DFW's administration of the suction dredging program violated CEQA and various
21 provisions of the Fish and Game Code.

22 **Irrigated Lands Program**

23 Staff of the North Coast RWQCB are developing an Agricultural Lands Discharge Program to address
24 water quality impacts associated with irrigated agricultural lands in the North Coast region. Agricultural
25 lands have the potential to contribute to water quality problems through the over-application of fertilizers
26 and pesticides, human-caused erosion of sediment, pollutants in tailwater return flows, and the removal
27 and suppression of riparian vegetation. The RWQCB staffs are developing the program to address these
28 water quality issues and to meet the requirements of the California Water Code, the State Nonpoint
29 Source Policy, and the Klamath River TMDLs.

30 While the scope of the program has not been finalized, it will include certain types of agricultural lands in
31 the North Coast region and address discharges of waste. Staff expects the program to address, at
32 minimum, waste discharges from agricultural lands such as row crops, vineyards, orchards, medicinal
33 marijuana farms, nurseries, forage crops, and irrigated pasture. Dairies and dryland grazing are not
34 included in the program as dairies are being addressed through a separate RWQCB program, and dryland
35 grazing is likely to be addressed through a statewide effort that is currently under development.
36 Additionally, this effort will be coordinated with existing RWQCB programs, such as the TMDL
37 programs in the Scott, Shasta, and Garcia watersheds and grazing on USFS allotments.

38 **Tribal Water Rights**

39 Water rights in California have a long and complicated history. The interplay between State water law
40 and tribal water rights is especially complex in California for several reasons. First, while other western

1 states operate under a prior appropriation system, California maintains a system of both property-based
 2 rights and prior appropriation rights. Second, over 100 federally recognized Native American tribes are
 3 located in California — by far, more tribes than in any other state. A tribe’s individual history plays an
 4 important role in defining their water rights, thus requiring a review of each tribe’s history in order to
 5 accurately quantify each tribe’s rights. No historical reviews have been completed for the majority of
 6 California's Native American tribes. Third, California contains over 300 individual Native American
 7 allotments, located both on reservations and in the public domain. Each of these requires its own
 8 historical review, but to date there have been very few reviews of individual allotments.

9 Federally reserved waters on Native American reservations are governed by the Winters doctrine, which
 10 has evolved over more than a century in federal courts, and since 1955 in state courts as well. Two
 11 landmark U.S. Supreme Court cases, *Winters v. U.S.*⁵ and *U.S. v. Rio Grande Dam & Irrigation Co.*,⁶
 12 established several key principles: (1) federally reserved lands have a right to use sufficient water to fulfill
 13 the “primary purpose” of the reservation, and (2) these water rights cannot be destroyed by state water
 14 law or by water users acting in accordance with state law. Evaluation of a tribe’s water rights requires a
 15 determination of two factors: (1) the date on which the land became federally reserved (the “priority
 16 date”), and (2) the amount of water needed to fulfill the “primary purpose” for which the land was
 17 federally reserved (California Tribal Water Rights 2009).

18 *Tribal Water Rights on the Klamath and Trinity Rivers*

19 Interconnection of the Trinity and Sacramento rivers adds federally reserved Native American water and
 20 fishing rights to California’s Central Valley water Issues. Historically, the fishery resources of the
 21 Klamath and Trinity rivers have been the mainstay of the life and culture of the Hoopa Valley Tribe. The
 22 fishery was “not much less necessary to the existence of the Indians than the atmosphere they breathed.”
 23 *Blake v. Arnett*, 663 F.2d 906, 909 (9th Cir. 1981). The salmon fishery is central to Hoopa culture and its
 24 economy. The lower 12 miles of the Trinity River and a stretch of the Klamath River flow through the
 25 Hoopa Valley Reservation, established in 1864.

26 The Trinity River Division of the CVP was authorized in 1955 and completed in 1963. The Trinity River
 27 Division Act authorized the TRD (Trinity River Diversion). The TRD is the only source of water
 28 imported by the CVP to the Central Valley from within the region. Congress included area-of-origin
 29 protections for the Trinity River, including one establishing flow release procedures for Trinity River fish
 30 and wildlife preservation and propagation. The USBR informed Congress that it would divert
 31 approximately 50 percent of Trinity River water into the Sacramento River. However, until the 1992
 32 enactment of the CVPIA, Pub. L. 102-575, the USBR consistently diverted 90 percent of the Trinity
 33 River water. That procedure not only created undue reliance on water resources in the Central Valley, but
 34 it also devastated the Trinity River fishery (Hoopa Valley Indian Tribe, California Tribal Water Summit
 35 2009). Please see the “Tribal Water Rights on the Klamath and Trinity Rivers” Box NC-3 for more
 36 information.

37 **PLACEHOLDER Box NC-3 Tribal Water Rights on the Klamath and Trinity Rivers**

38 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 39 the end of the report.]

40 In March of 2013, the state of Oregon backed the Klamath Tribes' claim to have the oldest water rights in
 41 the upper Klamath Basin. The findings filed with the Klamath County Circuit Court in Klamath Falls

1 gives the tribes a new dominant position in the long-standing battles over sharing scarce water between
2 fish and farms in the Upper Klamath Basin. Farmers and ranchers who draw irrigation water from rivers
3 where the tribes now have the oldest claim could be restricted in drought years. As of the writing of this
4 report, the impact of this for California water users is unclear (Oregon backs Klamath Tribes water rights
5 2013).

6 **Water Quality**

7 *Surface Water Quality*

8 Large portions of the North Coast are listed for TMDLs. Unfortunately, rural regions have difficulties
9 maintaining State and federal drinking water standards due to financial and technical issues.

10 The surface water quality issues of most concern in the North Coast region are excess sediment, elevated
11 water temperatures, and excess nutrients. These water quality conditions are the result of point and
12 nonpoint sources (NPS) of pollution and other controllable factors (e.g., landscape alteration, road
13 building, etc.) and are exacerbated by hydrologic modification, water withdrawal, and the loss of
14 competent riparian zones and floodplains to development, agriculture, and logging. Many North Coast
15 aquatic ecosystems are impacted by these constituents and controllable factors, resulting in a loss of
16 streamside property to erosion, destruction of water intakes, loss of aquatic habitat and risk to threatened
17 and endangered aquatic species, increased winter flood potential, and increased risk of summer nuisance
18 algal blooms (including *microcystis* and other cyanobacteria).

19 There are more localized issues, as well. For example, surface water monitoring indicates a problem with
20 pathogens in Bodega Bay Hydrologic Area, Hare Creek Beach, and Pudding Creek Beach on the
21 Mendocino Coast, several coastal beaches in the Trinidad Hydrologic Unit, and riverfront beaches on the
22 Russian River and its tributaries, as well as the Laguna de Santa Rosa and its tributaries. In addition,
23 several of the region's water bodies are impaired by mercury from gold mining in the past, including Lake
24 Pillsbury, the Laguna de Santa Rosa, Lake Sonoma, Trinity Lake, and the East Fork Trinity River. Exotic
25 species are listed as a water quality problem in Bodega Bay and dioxin and PCBs are listed as impairing
26 Humboldt Bay.

27 **Nonpoint Source Pollution (NPS)**

28 NPS pollution in the region includes contamination of surface water due to NPS pollution from storm
29 water runoff, erosion, and sedimentation (roads, agriculture, and timber harvest), failing septic tanks,
30 channel modification, gravel mining, dairies, MTBE and dioxin contamination (from lumber mills) and
31 urban runoff. In areas where people can come into contact with contaminated waters, the SWRCB, North
32 Coast RWQCB, and California Coastal Commission have the responsibility to protect the people. Among
33 other priorities, one of the highest priorities of the North Coast RWQCB Basin Plan is to develop a
34 freshwater beach program in cooperation with the Sonoma County Health Department for the Russian
35 River. Sediment, temperature, and nutrients are the items of primary focus in the RWQCB 303(d) list of
36 impaired water bodies. Along the coast, NPS pollution can cause microbial contamination of shellfish
37 (and in particular, oyster) growing areas. In rivers, lakes, and reservoirs in the Klamath Basin, extreme
38 growths of blue green algae and accompanying microcystin neurotoxins have been found in high
39 concentrations, leading to issuance of a health advisory by the State.

1 **Mercury**

2 Mercury in fish tissue is a water quality concern in Lake Pillsbury (Eel River), Lakes Mendocino and
3 Sonoma (Russian River), and Trinity Lake (Trinity River); health advisories for mercury have been issued
4 for Lake Pillsbury and Trinity Lake.

5 **Erosion and Sedimentation**

6 The RWQCB has prepared a Work Plan to Control Excess Sediment in Sediment-Impaired Watersheds
7 (04-08-2008). The plan describes actions and tasks that staff is doing or intends to do over the next
8 10 years (as resources allow) to control human-caused excess sediment transport in the sediment-impaired
9 water bodies of the region. Besides harming aquatic life, excess sediment can limit the use of water for
10 municipal and domestic consumption, agriculture, industry, wildlife, fishing, and recreation, and it can
11 cause or contribute to flooding. When sediment transport and increased runoff do occur, they cause
12 changes in the downstream channels. These changes include gravel and sand deposition creating gravel
13 bars, degrading spawning habitat and scouring of stream channels due to higher flows.

14 **Five Counties Salmonid Conservation Program (5C Program)**

15 In 1997, the northwestern California counties of Del Norte, Humboldt, Mendocino, Siskiyou, and Trinity
16 agreed to collaborate on a proactive, positive response to the federal listing of coho salmon as a
17 threatened species by forming the Five Counties Salmonid Conservation Program (5C Program). The
18 primary 5C goal is "to strive to protect the economic and social resources of northwestern California by
19 providing for the conservation and restoration of salmonid populations to healthy and sustainable levels
20 and to base decisions on watershed rather than county boundaries.

21 In February 2009, the 5C Program transferred from Trinity County administration to the Northwest
22 California Resource Conservation & Development Council whose mission is to enhance the ability of
23 area residents to develop diverse opportunities through the utilization of available resources. The program
24 maintains its relationship with all five counties and will continue to build on the watershed restoration and
25 planning work that has been integral to the program over the past years.

26 The 5C Program's specific objectives include:

- 27 1. Improve county policies and road maintenance practices with a strong emphasis on training.
- 28 2. Identify potential restoration opportunities through inventories of fish passage barriers and potential
29 sediment sources on county maintained roads.
- 30 3. Increase the amount of salmonid habitat by replacing stream crossings that are barriers to migration
31 with structures that provide for passage. Improve water quality by treating identified sources of
32 road related sediment.
- 33 4. Devise methods to streamline permitting procedures, specifically under the ESA, CA ESA, the
34 Clean Water Act, and California Fish and Game Code.
- 35 5. Collaborate with other organizations, agencies, and regional groups on restoration and conserva-
36 tion.
- 37 6. Develop model regulations only where other means cannot be utilized to address land use activities
38 regulated by the Counties.
- 39 7. Secure grant program and project funding from a variety of Federal, State, and Local sources.

40 The 5C Program is highly effective in promoting and sustaining collaborative efforts that capitalize on
41 technical assets of participants and in leveraging financial support from numerous sources. 5C recognizes

1 that taking on these challenges will lead to a healthier environment, sustainable fisheries, and better
2 County facilities, all of which contribute to a more robust economy (Five Counties Salmonid
3 Conservation Program 2013).

4 **Local Tribes Cooperation**

5 Another group recently formed, although informal, is a collaboration of tribes in the North Coast led by
6 the Cher-Ae-Heights Indian Community of the Trinidad Rancheria. This group was formed to assist local
7 tribes interested in collaborating to develop an environmental assessment and implementation plan for
8 improving ecosystems and water quality in order to meet or exceed federal and State regulations regard-
9 ing water quality. Tribes currently involved in this collaboration include the Trinidad Rancheria in Trini-
10 dad, Blue Lake Rancheria Tribe in Blue Lake, Bear River Tribe in Loleta, and Big Lagoon Rancheria in
11 Arcata. One main function of the cooperation is to assist the members in obtaining grant funding for local
12 water quality infrastructure improvements.

13 *Groundwater Quality*

14 In 2009, the USGS, in conjunction with the SWRCB, collected untreated groundwater data from 58 wells
15 selected from the California Department of Public Health database within 34 groundwater basins located
16 in the North Coast region. Randomly selected wells included locations in Lake, Mendocino, Glenn,
17 Humboldt, and Del Norte counties. The results of the study are published in Mathany et al. (2011). All
18 detected concentrations of organic constituents, nutrients, major and minor ions, and radioactive
19 constituents were less than health-based benchmarks for the 30 wells sampled in the northern Coast
20 Ranges. There were a few detections of arsenic, boron, and barium in the 28 wells of the interior basins,
21 which exceeded MCLs or notification levels; but, these are likely related to the area's geology. The
22 results of this study indicate that community drinking water systems drawing from primary aquifer
23 systems in the North Coast region generally provide safe drinking water, with the exceptions noted.

24 Because the North Coast region is predominantly rural, many people rely on shallow (sometimes hand
25 dug) wells for their drinking water. Shallow groundwater cleanup, therefore, remains a high priority in
26 the region.

27 There may be contributions of nutrients and pesticides to shallow groundwater resulting from the
28 continued conversion of land to vineyards in Sonoma and Mendocino counties and other widespread
29 farming activities in the Upper Klamath River Basin and the Smith River Plain, among other disperse
30 locations of the region. Aging wastewater treatment ponds and leaking septic tanks may play a part in
31 shallow groundwater contamination in these areas as well (Mathany TM et al. 2011).

32 *Drinking Water Quality*

33 In general, drinking water systems in the region deliver water to their customers that meet federal and
34 State drinking water standards. Recently, the Water Boards completed a draft statewide assessment of
35 community water systems that rely on contaminated groundwater. This draft report identified 15
36 community drinking water systems in the region that rely on at least one contaminated groundwater well
37 as a source of supply (Table NC-14). Arsenic is the most prevalent groundwater contaminant affecting
38 community drinking water wells in the region (Table NC-15). The majority of the affected systems are
39 small water systems which often need financial assistance to construct a water treatment plant or alternate
40 solution to meet drinking water standards.

1 **PLACEHOLDER Table NC-14 Summary of Community Drinking Water Systems**
 2 **in the North Coast Hydrologic Region that Rely on One or More**
 3 **Contaminated Groundwater Well that Exceeds a Primary Drinking Water Standard**

4 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 5 the end of the report.]

6
 7 **PLACEHOLDER Table NC-15 Summary of Contaminants Affecting Community**
 8 **Drinking Water Systems in the North Coast Hydrologic Region**

9 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 10 the end of the report.]

11 **Groundwater Conditions and Issues**

12 *Groundwater Occurrence and Movement*

13 Aquifer conditions and groundwater levels change in response to varying supply, demand, and climate
 14 conditions. During dry years or periods of increased groundwater use, seasonal groundwater levels tend to
 15 fluctuate more widely and, depending on annual recharge conditions, may result in a long-term decline in
 16 groundwater levels, both locally and regionally. Depending on the amount, timing, and duration of
 17 groundwater level decline, nearby well owners may need to deepen wells or lower pumps to regain access
 18 to groundwater.

19 Lowering of groundwater levels can also impact the surface water–groundwater interaction by inducing
 20 additional infiltration and recharge from surface water systems, thereby reducing the groundwater
 21 discharge to surface water base flow and wetlands areas. Extensive lowering of groundwater levels can
 22 also result in land subsidence due to the dewatering, compaction, and loss of storage within finer grained
 23 aquifer systems.

24 During years of normal or above normal precipitation, or during periods of low groundwater use, aquifer
 25 systems tend to recharge and respond with rising groundwater levels. As groundwater levels rise, they
 26 reconnect to surface water systems, contributing to surface water base flow or wetlands, seeps, and
 27 springs.

28 The movement of groundwater is from areas of higher hydraulic potential to areas of lower hydraulic
 29 potential, typically from higher elevations to lower elevations. The direction of groundwater movement
 30 can also be influenced by groundwater extractions. Where groundwater extractions are significant,
 31 groundwater may flow towards the extraction point. Rocks with low permeability can restrict
 32 groundwater flow through a basin. For example, a fault may contain low permeability materials and
 33 restrict groundwater flow.

34 *Depth to Groundwater*

35 The depth to groundwater has a direct bearing on the costs associated with well installation and
 36 groundwater extraction operations. Understanding the local depth to groundwater can also provide a
 37 better understanding of the local interaction between the groundwater table and the surface water systems,
 38 and the contribution of groundwater aquifers to the local ecosystem.

1 Groundwater levels in the North Coast Hydrologic Region are highly variable from basin to basin.
 2 Resource and time constraints, depth-to-groundwater contours for the region could not be developed as
 3 part of the groundwater content enhancement for Update 2013. However, depth-to-groundwater data for
 4 some of the groundwater basins in the region are available online via DWR's Water Data Library, DWR's
 5 CASGEM system, and the USGS National Water Information System. In addition, basin-specific
 6 information may be obtained from the following sources. Please note that although a reference for
 7 Sonoma Valley is provided below, groundwater basins encompassing the southern portion of Sonoma
 8 County are discussed in the Regional Report for the San Francisco Hydrologic Region.

- 9 • Ground-Water Hydrology of the Upper Klamath Basin: USGS, 2010 Scientific Investigations
 10 Report 2007-5050. (<http://pubs.usgs.gov/sir/2007/5050/>)
- 11 • Santa Rosa Valley: Sonoma County Water Agency (<http://www.scwa.ca.gov/srgroundwater/>).
- 12 • Scott Valley Groundwater: UC Davis: (<http://groundwater.ucdavis.edu/Research/ScottValley/>).
- 13 • Sonoma Valley: USGS, 2006. (<http://pubs.usgs.gov/sir/2006/5092/pdf/sir2006-5092.pdf>)

15 **Groundwater Elevations**

16 Groundwater elevation contours can help estimate the direction of groundwater movement and the
 17 gradient, or rate, of groundwater flow. Although DWR monitors the depth to groundwater in some
 18 groundwater basins within the region, because of resource and time constraints groundwater elevation
 19 contours for the North Coast Hydrologic Region could not be developed as part of the groundwater
 20 content enhancement for Update 2013. Some references and links to local agencies that independently or
 21 cooperatively monitor the groundwater levels in the basins and develop groundwater elevation maps have
 22 been provided in the previous section.

23 **Groundwater Level Trends**

24 Plots of depth-to-water measurements in wells over time (groundwater level hydrographs) allow analysis
 25 of seasonal and long-term groundwater level variability and trend over time. Because of the highly
 26 variable nature of the physical aquifer systems within each groundwater basin, and because of the variable
 27 nature of annual groundwater availability, recharge, and surrounding land use practices, the hydrographs
 28 presented herein do not attempt to illustrate or depict average aquifer conditions over a broader region.
 29 Rather, the selected hydrographs are intended to help tell a story about how the local aquifer systems
 30 respond to changing groundwater pumping quantity and to the implementation of resource management
 31 practices. The hydrographs are designated according to the State Well Number System (SWN), which
 32 identifies each well by its location using the public lands survey system of township, range, section, and
 33 tract.

34 Hydrograph 48N04E31N002M

35 Hydrograph 48N04E31N002M (Figure NC-18a) is from a deep irrigation well that draws water from a
 36 fractured basalt portion of the aquifer underlying the Tule Lake subbasin and is located along the western
 37 edge of the Tule Lake subbasin. The Tule Lake subbasin has been designated as a CASGEM medium
 38 priority basin. In 2001, in response to one of the driest years on record for the Klamath Basin watershed,
 39 the USBR cutoff surface water deliveries from the Klamath Project to the Tule Lake subbasin area. In
 40 response, a drought emergency was declared and a number of new high-capacity wells were installed in
 41 the fractured-basalt portion of the Tule Lake subbasin aquifer. In subsequent years, ongoing
 42 environmental water shortages for the Klamath Project resulted in additional surface water cutbacks and
 43 the implementation groundwater substitution water transfers in nine out of the subsequent ten years. Due

1 to Oregon regulations that limit groundwater pumping, the majority of groundwater substitution pumping
2 came from the California portion of the Klamath Basin (Note: almost two thirds of the 210,000 irrigated
3 acres in the Klamath Project service area is in Oregon). In 2000, prior to the groundwater substitution
4 pumping, groundwater supply required from the Tule Lake subbasin was estimated to be 8,500 af. Over
5 the next couple of years, transfer operations resulted in groundwater extraction of 70,000 af in 2001 and
6 about 22,000 af /yr for 2002 and 2003 (DWR 2004). Groundwater pumping increased to 32,000 af in
7 2004 and then declined to an average of about 14,000 af a year for 2005 and 2006. No pumping amounts
8 were recorded for 2007 and 2008. No groundwater substitution transfers took place in 2009; however,
9 non-transfer related pumping of 8,500 af was estimated. But in 2010, groundwater extraction volume
10 increased to 51,000 af.

11 Although there is considerable annual variation in groundwater levels between 2001 and 2010, the
12 hydrograph shows that the overall rate of basin recharge has not been able to keep pace with the post
13 2001 increases in groundwater extraction. After the initial drop of seven feet between 2001 and 2002, the
14 hydrograph shows a slow but steady trend of declining groundwater levels until 2006, a period of
15 relatively stable levels from 2006 through 2009, and then another drop from 2009 to 2012. The period of
16 somewhat stable groundwater levels from 2006 to 2009 indicates that the annual rate of aquifer recharge
17 was likely sufficient to offset the average annual groundwater pumping volume of about 14,000 af.
18 Conversely, the post 2009 decline in groundwater levels associated with the increase in groundwater
19 extraction to 51,000 in 2010 indicates that annual extraction rates of 50,000 af /yr are not sustainable for
20 this portion of the basin aquifer. The hydrograph also highlights the importance of implementing
21 appropriate data collection and adaptive management practices when implementing conjunctive
22 management via groundwater substitution — especially in areas where aquifer response to increased
23 pumping is largely unknown. At the local level, a decline of 17 feet over 12 years in response to
24 groundwater substitution have resulted in impacts to shallow wells, increased the risk for future
25 subsidence within the fine-grained lakebed deposits above the fractured-basalt aquifer, and are bringing
26 into question the sustainability of land use practices that require greater than about 40,000 af of
27 groundwater extraction.

28 Uncertainties associated with the operation of the Klamath Project Water have led to the development of
29 the Klamath Water and Power Agency (KWAPA) to help align water supply and use. An On-Project Plan
30 is being implemented by KWAPA to help align long-term water supply and demand for the local service
31 area. In addition, conservation and management practices are currently being implemented by the Tule
32 Lake Irrigation District to help increase water supply reliability (Tulelake Irrigation District 2011).

33 Hydrograph 44N06W10F001M

34 Hydrograph 44N06W10F001M (Figure NC-18b) is from a 113-foot-deep domestic well that draws water
35 from shallow aquifer that consists of sand, gravel, clay and volcanic deposits, and is located near Grenada
36 in Shasta Valley Groundwater Basin, about 50-feet down gradient from Montague Water District
37 conveyance ditch. Shasta Valley basin is designated as a CASGEM medium priority basin. The
38 hydrograph for this well highlights dramatic seasonal effects of conveyance ditch losses to the underlying
39 shallow aquifer and the wells that draw water from it. Throughout most of California, precipitation
40 associated with Mediterranean climate conditions typically result in seasonal groundwater levels being the
41 highest during late winter to early spring months, and the lowest during summer or early fall months.
42 However, groundwater levels for well 44N06W10F001M are consistently 5- to 10-feet higher in the fall
43 relative to that in the spring. This reversed groundwater level trend is likely due to summer recharge from

1 conveyance ditch losses and the percolation of applied surface water for nearby agricultural water use.
 2 Once the irrigation season is over, the conveyance system is dewatered; and nearby groundwater levels
 3 decline. Prior to 2007, there were two conveyance canals located parallel to each other. In 2007, one of
 4 the two canals was replaced with an underground pipe system to reduce conveyance losses. This resulted
 5 in the overall lowering of the groundwater level by more than 5 feet, as shown in the hydrograph. The
 6 reversed trend of seasonal fluctuation continued, but at a lower elevation — indicating that the leakage
 7 from the remaining conveyance ditch is still occurring.

8 Hydrograph 02N01W08B001H

9 Hydrograph 02N01W08B001H (Figure NC-18c) is for a very shallow irrigation well constructed in the
 10 aquifer consisting of unconfined sand and gravel deposits in the Eel River Valley Groundwater Basin. Eel
 11 River Valley is designated as a CASGEM medium priority basin. The hydrograph highlights the close
 12 interaction between surface water systems and the numerous shallow groundwater wells that draw water
 13 from thin alluvial river plain aquifers along the California coast. Land use surrounding the well is
 14 predominantly rural pasture and dairy cattle. The hydrograph shows seasonal fluctuations in groundwater
 15 levels of about 6 to 8 feet during normal and drought years, and approximately 12 to 13 feet during wet
 16 years. A long-term comparison of spring-to-spring groundwater levels in the well shows a very slight
 17 decline and recovery of groundwater levels associated with the 1976-77 and the 1987-92 droughts.
 18 Groundwater levels in wells that are closely connected to nearby perennial surface water systems are
 19 typically more affected by wet rather than drought years. Perennial surface water systems tend to provide
 20 a consistent source of recharge which helps to govern the maximum seasonal decline in groundwater
 21 levels. Spring-to-spring groundwater levels in during years of normal precipitation show a trend of
 22 slightly declining groundwater levels since the late 1960s.

23 Hydrograph 06N08W15J003M

24 Hydrograph 06N08W15J003M (Figure NC-18d) is from an inactive well constructed in upper
 25 160 feet within the alluvial deposits and the Glen Ellen Formation of the aquifer in the Santa Rosa Plain
 26 Subbasin located in southern Sonoma County. Santa Rosa Plain Subbasin is designated as a CASGEM
 27 medium priority basin. The hydrograph depicts changes in groundwater supply and conjunctive
 28 management practices and shows the relationship between groundwater elevations and increased surface
 29 water supplies. The area surrounding the well is a combination of suburban residential and commercial
 30 land use. From 1950 to 1986, the groundwater elevation in the well declined approximately 50 feet due to
 31 groundwater extraction. During this time, municipal groundwater pumping in the southern Santa Rosa
 32 Plain increased from less than 1,000 af in 1969 to more than 5,000 af in 1986, while surface water
 33 deliveries during this time averaged less than 500 af /yr, with some years having no surface water supply
 34 to the area. Sonoma County Water Agency began increasing its municipal surface water deliveries in
 35 1986 from approximately 1,000 af /yr to more than 4,000 af /yr in 2003, and then to 6,000 af /yr in 2005.
 36 Between 1986 and 2000, groundwater continued to be pumped at a volume between 5,000 and 6,000 af
 37 /yr. As shown on
 38 Figure NC-17d, groundwater elevations did not start to recover until 2003 when groundwater pumping
 39 was reduced to less than 2,000 af in 2003 and about 500 af in 2005. The 40-foot groundwater level
 40 recovery between 2003 and 2005 was also the result of increased surface water deliveries from 4,000 af to
 41 6,000 af /yr. The conjunctive management efforts in the Santa Rosa Plain Groundwater Subbasin not only
 42 reflect the connection between groundwater extraction and surface water availability, but also the positive
 43 effects of water conservation and the use of recycled water supplies for irrigation.

1 **PLACEHOLDER Figure NC-18 Groundwater Level Trends in Selected Wells in the North Coast**
2 **Hydrologic Region**

3 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
4 the end of the report.]

5 *Change in Groundwater Storage*

6 Change in groundwater storage is the difference in stored groundwater volume between two time periods.
7 Examining the annual change in groundwater storage over a series of years helps identify the aquifer
8 response to changes in climate, land use, or groundwater management over time. If the change in storage
9 is negligible over a period represented by average hydrologic and land use conditions, the basin is
10 considered to be in equilibrium under the existing water use scenario and current management practices.
11 However, declining storage over a period characterized by average hydrologic and land use conditions
12 does not necessarily mean that the basin is being managed unsustainably or subject to conditions of
13 overdraft. Utilization of groundwater in storage during years of diminishing surface water supply,
14 followed by active recharge of the aquifer when surface water or other alternative supplies become
15 available, is a recognized and acceptable approach to conjunctive water management. Additional
16 information regarding the risks and benefits of conjunctive management are presented in Update 2013,
17 Volume 3, Chapter 9, “Conjunctive Management and Groundwater Storage.”

18 Because of resource and time constraints, changes in groundwater storage estimates for basins within the
19 North Coast Hydrologic Region were not developed as part of the groundwater content enhancement for
20 Update 2013. It is unknown if any of the local groundwater management agencies within the region have
21 developed change in groundwater storage estimates.

22 **Flood Management**

23 Traditional flood management has been focused on flood control infrastructure projects. These
24 infrastructures alter or confine natural watercourses — hydromodification — which are intended to
25 reduce the chance of flooding thereby minimizing damage to lives and property. This traditional approach
26 is based on the flood control principle of conveying floodwaters rapidly to a discharge point. A more
27 current understanding of floods and flooding takes into account the role of watershed management,
28 floodplain and river functions, and providing multiple resource management and societal benefits.
29 Activities under traditional flood management include physical modification of stream channels, dam and
30 surface impoundments, levees, and other structures.

31 Today, water resources and flood planning involves additional demands and challenges, such as multiple
32 regulatory processes and permits, coordination with multiple agencies and stakeholders, and increased
33 environmental awareness. These additional complexities call for an integrated water management
34 approach, incorporating natural hydrologic, geomorphic, and ecological processes to reduce flood risk by
35 influencing the cause of the harm, including the probability, extent, or depth of flooding (flood hazard).
36 State policy directs State agencies to implement integrated water management and other federal, regional,
37 and local agencies are transitioning to this approach. Integrated water management changes the
38 implementation approach based on the understanding that water resources are an integral component for
39 sustainable ecosystems, economic growth, water supply reliability, public health and safety, and other
40 interrelated elements. Additionally, it acknowledges that a broad range of stakeholders might have
41 interests and perspectives that could positively influence planning outcomes.

1 Projects that combine flood and ecosystem restoration also can provide areas of active- and passive-use
 2 recreation, increase open space, and provide scenic value, all of which result in economic and societal
 3 benefits. For example, in Humboldt County, the Rohner Creek Flood Control and Riparian Habitat
 4 Improvement project is a watershed-based, channel corridor-scale project with multiple objectives. The
 5 proposed project is taking a channel corridor approach in identifying opportunities to integrate habitat
 6 enhancement elements with flood reduction improvements through the 1-mile project corridor within the
 7 City of Fortuna (California’s Flood Future: Recommendations for Managing the State’s Flood Risk,
 8 Public Draft April 2013).

9 *Flood Hazard Exposure*

10 Historically, in the North Coast Hydrologic Region, flooding originates principally from melting of the
 11 Coastal Ranges snowpack and from rainfall. Flooding from snowmelt typically occurs in the spring and
 12 has a lengthy runoff period. Flooding from rainfall occurs in the winter and early spring, particularly
 13 when storms arriving from the Gulf of Alaska draw moisture-laden air from the tropics. This pattern is
 14 known as an Atmospheric River. This pattern also creates coastal storms that drive waves resulting in
 15 coastal flooding and erosion. Offshore earthquakes have caused tsunamis along the coast in the
 16 hydrologic region.

17 Flood exposure in the North Coast Hydrologic Region occurs along the coastline, Eel River, Scott River,
 18 around Crescent City Harbor and, Humboldt Bay. (See also Box NC-2 Near-Coastal Issues.) Flood
 19 exposure identifies who and what is impacted by flooding. In the North Coast Hydrologic Region, more
 20 than 43,000 people and over \$4.2 billion in assets are exposed to the 500-year flood event. Figures NC-19
 21 and NC-20 provide a snapshot of people, structures, crops, infrastructure, and sensitive species exposed to
 22 flooding in the region. Throughout the region, 320 State and federal threatened, endangered, listed, or rare
 23 plant and animal species are exposed to flood hazards.

24 **PLACEHOLDER Figure NC-19 Flood Exposure to the 100-Year Floodplain, North Coast Hydrologic** 25 **Region**

26 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 27 the end of the report.]

28 **PLACEHOLDER Figure NC-20 Flood Exposure to the 500-Year Floodplain, North Coast Hydrologic** 29 **Region**

30 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 31 the end of the report.]

32 *Sea Level Rise*

33 During the coming decades, sea level will continue to rise, bringing with it progressive flooding and
 34 inundation of low-lying areas as well as increased cliff and bluff erosion. North Coastal areas will be
 35 challenged to adapt to this rise especially in urban and rural coastal developed areas and ports, harbors
 36 and marinas with commercial and recreational facilities. It is imperative to minimize damage and losses
 37 through adaptation. Coastal managers are relying on historical coastal hazard-vulnerability data and
 38 projecting the types of hazards and risk associated with sea level rise. While the types of hazards may not
 39 change, their frequencies and magnitudes are changing, which will increase community vulnerability and
 40 risk.

1 Sea level rise will affect and threaten coastal communities, facilities and infrastructure through more
 2 frequent flooding and gradual inundation, as well as increased erosion of coastal bluffs, and river surges
 3 affecting local flooding. This will affect roads, utilities, wastewater treatment plants, outfalls and storm
 4 water facilities and systems as well as large wetland areas in addition to towns and cities. Where land is
 5 rising — tectonic effects — the rate of sea level rise may be exceeded by the rate of coastal uplift. In the
 6 North Coastal areas the rate of tectonic uplift is greater than current rate of sea level rise. For example, at
 7 Humboldt Bay's North Spit, sea level is rising by 18.6 inches per century (4.73 millimeters per year), the
 8 highest rate in California. At Crescent City, 80 miles north, sea level is dropping relative to the coastline
 9 by 2.5 inches per century. The shoreline at Humboldt Bay is subsiding, whereas Crescent City's coastline
 10 is rising.

11 The risk assessment for flooding is incorporating the vulnerability of the North Coast region based on the
 12 rate and magnitude of sea level rise and its impacts. Those communities and facilities at risk are
 13 incorporating hazard mitigation measures into planning and management strategies. As the "California
 14 Flood Futures" report identifies, the first strategy is to identify and evaluate sea level rise risks and
 15 determine those areas most vulnerable to future flooding, inundation, erosion, and wave impacts and to
 16 develop hazard mitigation and adaptation plans.

17 Where coastal bluff erosion is high, coastal cliff retreat is dramatic with collapsed roadways, undermined
 18 foundations, dangling decks and stairways and structures. Coastal erosion tends to be episodic, with long-
 19 term cliff and bluff failure occurring during a few severe storm events. Scientists consider the probability
 20 that these events will increase in frequency and intensity. The California Coastal Commission database
 21 for coastal erosion is a valuable resource and available on CD (Dare 2005). A key component to coastal
 22 management is understanding the adaptive capacity of the affected areas. This capacity is the ability to
 23 prepare for, respond to, and recover from sea level rise impacts.

24 As described earlier, the general principles of integrated water management includes adaptation planning
 25 to embrace sustainability (i.e., meeting the needs of the present without compromising the needs of future
 26 generations) with consideration given to equitable distribution and apportionment of costs and benefits of
 27 adaptation measures; and adaptation strategies should account for the distinct vulnerability of potentially
 28 affected DACs.

29 *Levee and Channel System*

30 The North Coast Hydrologic Region has four major flood management reservoirs — Lake Mendocino on
 31 the East Fork Russian River, Lake Sonoma on Dry Creek, Spring Lake off Santa Rosa Creek, and
 32 Matanzas Creek Reservoir on Matanzas Creek; two smaller flood management reservoirs on Paulin Creek
 33 and Middle Fork Brush Creek; and seven other reservoirs providing nondedicated flood-retention space.
 34 Other flood management projects include levees in the Eel River delta, levees and channel modifications
 35 on East Weaver Creek, Redwood Creek, the Klamath River, and the Mad River, and channel
 36 modifications on Santa Rosa Creek. Measures to mitigate the effects of tsunamis were part of Humboldt
 37 Harbor improvements, the Crescent City project, and Crescent City Harbor improvements.

38 *Levee Performance and Risk Studies*

39 In the North Coast Hydrologic Region, 26 local flood management projects or planned improvements
 40 have been identified. Fourteen of these projects have costs totaling more than \$108 million while the
 41 remaining projects do not have costs associated with them at this time. Fifteen local planned projects use

1 an integrated water management approach with a flood component. Examples of these local projects
 2 include the Mattole Integrated Watershed Management Initiative and the Big River Main Haul Road
 3 Phase I Restoration Project. For a complete list of these projects refer to "California's Flood Future
 4 Report" Attachment F: Information Gathering Technical Memorandum.

5 *Recent Tsunamis on the California Coast*

6 In March 2011, a tsunami generated off the coast of Japan and recorded throughout the California coast,
 7 struck Crescent City Harbor with an 8.1-foot wave, destroying much of the harbor and resulting in one
 8 death near Klamath. There was also major damage to docks and boats at Noyo Harbor. Estimated damage
 9 in the region was \$24 million.

10 *Redwood Coast Tsunami Work Group*

11 The Redwood Coast Tsunami Work Group (RCTWG) is an organization of local, State, and federal
 12 agencies, tribes, relief and service groups, land managers, and businesses from Del Norte, Humboldt and
 13 Mendocino counties. The group was formed in July 1996 to define the needs of local jurisdictions to
 14 mitigate the North Coast earthquake and tsunami hazard and to promote a coordinated, consistent
 15 mitigation program for all coastal areas. (See "Recent Tsunamis on California Coast" section in this
 16 report.)

17 In 2006, Humboldt County participated in FEMA's first ever tsunami response training exercise. In 2007,
 18 RCTWG helped the community of Samoa prepare for and conduct the first full-scale tsunami evacuation
 19 drill in California. In 2008, RCTWG members working with the California Governor's Office of
 20 Emergency Services planned and coordinated the first test of the tsunami warning communications
 21 system using actual (live) codes outside of Alaska. Additional drills have been conducted in schools and
 22 other North Coast communities. For more information on RCTWG, see the University of Humboldt's
 23 Web page at, <http://humboldt.edu/rctwg/site/about/>.

24 **Water Governance**

25 The North Coast region contains water service providers of all types, from small, private facilities that
 26 provide water for just a few neighboring residences to large municipal suppliers and wastewater treatment
 27 facilities. Private water districts include those representing counties or portions of counties,
 28 municipalities, irrigation districts, or particular water bodies. The only federal water boundaries in the
 29 region are Redwood Valley District in Mendocino and in the Klamath Lake and Tule Lake area as part of
 30 the Klamath Project, which are administered by USBR. A large number of North Coast residences are in
 31 rural areas with no water service and rely on groundwater wells or personal surface-water treatment
 32 facilities and onsite wastewater disposal systems, usually septic systems (North Coast Integrated Regional
 33 Water Management Plan, Phase III c2012). For a list of North Coast region's water management agencies,
 34 see Table NC-16.

35 **PLACEHOLDER Table NC-16 North Coast Hydrologic Region Water Management Agencies**

36 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 37 the end of the report.]

1 In 2009, state lawmakers passed four policy bills, the Safe, Clean, and Reliable Drinking Water Supply
2 Act of 2010, as a comprehensive water package (Water Conservation Act of 2009, SB x7-x). *For more*
3 *information on SB x7-x, see Update 2013 Volume 4, Reference Guide.*

4 *AB 2409 (Nestande, 2010)*

5 AB 2409 amends section 10632 of the California Water Code (Urban Water Management Planning Act)
6 to require urban water suppliers to prepare and adopt water shortage contingency plans including the
7 identification and treatment of artificially supplied water features, i.e., ponds, lakes, waterfalls and
8 fountains; separately from swimming pools and spas (California Legislature 2009-2010).

9 *California Water Code Sec 1259.4 AB 2121*

10 Water Code Sec 1259.4 AB 2121 requires the SWRCB to adopt principles and guidelines for maintaining
11 instream flows in Northern California coastal streams for the purposes of water right administration. The
12 geographic scope of the policy includes all coastal streams from the Mattole River to San Francisco and
13 coastal streams entering San Pablo Bay, and extends to five counties: Marin and Sonoma and portions of
14 Napa, Mendocino, and Humboldt counties.

15 *Fish and Game Code Section 5653*

16 Because instream dredging is a popular activity in this region, it should be noted that there have been
17 changes to rules that affect these activities. On April 27, 2012, the Office of Administrative Law
18 approved updated regulations governing suction dredge mining under Fish and Game Code section 5653
19 et seq., CEQA, and the Administrative Procedures Act. DFW has closed suction dredging for the next
20 several years. However, the closures are moot, as a statewide moratorium has been in place since 2008
21 and is planned to expire in 2016 after a planned court decision on the issue. For more information on
22 Suction Dredging, see DFW web page located at: <http://www.dfg.ca.gov/suctiondredge/> and "Statewide
23 Instream Mining (Suction Dredging)" section in this document.

24 *Water Code Division 5, Sections 8,000 - 9,651*

25 Water Code Division 5, Sections 8,000 - 9,651, has special significance to flood management activities
26 and is summarized in California's Flood Future Report Pages Attachment E: Information Gathering
27 Technical Memorandum.

28 *AB 70 (2007) Flood Liability*

29 AB 70 (2007) provides that a city or county might be responsible for its reasonable share of property
30 damage caused by a flood, if the State liability for property damage has increased due to approval of new
31 development after January 1, 2008.

32 *AB 162 (2007) General Plans*

33 AB 162 (2007) requires annual review of the land use element of general plans for areas subject to
34 flooding, as identified by FEMA or DWR floodplain mapping. The bill also requires that the safety
35 element of general plans provide information on flood hazards. Additionally, AB 162 requires the
36 conservation element of general plans to identify rivers, creeks, streams, flood corridors, riparian habitat,
37 and land that might accommodate floodwater for purposes of groundwater recharge and stormwater
38 management.

1 *Fish and Game Code Section 1602 Streambed Alteration Permits*

2 California Fish and Game Code section 1602 requires that any water user that alters a streambed, stream
3 bank or undertakes any other stream alteration to file for a permit with the DFW prior to performing any
4 work. On December 24, 2012, the Siskiyou County Superior Court issued an opinion granting declaratory
5 relief for the Siskiyou County Farm Bureau in a case challenging the DFW attempt to require farmers to
6 obtain streambed alteration permits for all agricultural water diversions. The court found that Fish and
7 Game Code section 1602 (“Section 1602”) does not require notification to DFW for the act of diverting
8 water pursuant to a valid water right where there is no alteration to the bed, bank, or stream. Although a
9 Superior Court case, this opinion has important potential statewide implications. This became effective
10 January 1, 2013.

11 *Potter Valley Project FERC License*

12 The Potter Valley Project was first licensed as a hydroelectric power plant in 1922 by the Federal Power
13 Commission. The original 50 year license expired in 1972. From 1972 until 1982, the project was
14 operated with a license that was granted annually while discussions regarding the operation were
15 undertaken by PG&E, FERC, Fishery agencies, and stakeholders. In 1978 a final environmental impact
16 statement (EIS) was issued by FERC. Several years of discussion ensued until, in 1983, the project was
17 relicensed for 50 years (from the original expiration date of 1972). The 1983 settlement agreement was
18 signed by PG&E, DFW, and the counties of Humboldt, Mendocino and Sonoma. Part of the new license
19 was Article 39 which requires a 10-year study be undertaken to determine what the new project flows
20 impact was on salmon and steelhead and to adjust them accordingly.

21 A Fisheries Review Group (FRG) was formed which consisted of scientists from PG&E, USFWS, DFW
22 and the NMFS. In March of 1998, after 10 years of studies, the FRG completed their findings and a report
23 was filed with FERC recommending flow modifications. FERC began its EIS process. Over the next year,
24 two other entities, including the Round Valley Indian Tribes (RVIT) and SCWA, submitted proposals to
25 FERC for minimum flow releases. FERC held public scoping meetings and many organizations,
26 municipalities, water districts, environmental groups, and governmental agencies joined as interveners in
27 the process. A draft EIS was completed by FERC in February 1999. After further public meetings, many
28 comments, additional proposed alternatives, and new modeling inputs; FERC issued its final EIS in May
29 2000.

30 The FERC recommendation was based predominately on the FRG proposal prepared by the scientists
31 with the most history and knowledge of salmon and steelhead populations specifically in the section of
32 the main stem of the Eel River impacted by the project. The resulting complex flow regimes were
33 calculated in such a way as to make the project nearly invisible to the environment by releasing flows
34 below Cape Horn Dam to mimic natural flows as closely as possible.

35 After a lengthy Section 7 Consultation between NMFS, PG&E and FERC, under the Endangered Species
36 Act, NMFS produced a BO and Reasonable and Prudent Alternative (RPA) for the project flows and
37 submitted it to FERC in November 2002. The NMFS RPA generated extensive discussion between the
38 agencies and stakeholders that had been involved in the license amendment proceedings since 1983.
39 Ultimately, FERC issued a Final Order Amending the License for the Project January 28, 2004. The
40 project license expires April 14, 2022 (Potter Valley Irrigation District c2012).

1 *Hydropower, a Renewable Energy*

2 In 2013, the California Public Utilities Commission is considering accepting large hydropower facilities
 3 as qualified “renewable energy” resources. This would allow power generating utilities in California to
 4 include these large hydropower sources in their list of renewable energy resources helping them to meet
 5 the requirement of 33% by 2020 goal set by the CPUC, i.e., Renewables Portfolio Standard (RPS) (PUC
 6 399.11 et seq). Until 2013, large hydropower facilities (producing over 30 megawatts) were not allowed
 7 to be considered renewable energy due to environmental concerns over the use of dams and their effect on
 8 fisheries. However, this new legal development may have a short-term benefit to the region's counties
 9 (particularly utilities in Siskiyou County) because plans are to remove the hydroelectric dams on the
 10 Klamath River (Iron Gate, and both Copco dams) pursuant to the Klamath Basin Restoration Agreement
 11 (CPUC Renewables Portfolio Standard Program 2007, Klamath Facilities Removal Environmental Impact
 12 Statement/Environmental Impact Report 2011).

13 *State Funding Received*

14 DWR and SWRCB administer planning grants intended to foster development or completion of integrated
 15 regional water management plans or components thereof, to enhance regional planning efforts, and to
 16 assist more applicants to become eligible for implementation grant funding (Table NC-17).

17 **PLACEHOLDER Table NC-17 State Funding Received**

18 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 19 the end of the report.]

20 *Flood Governance*

21 California’s water resource development has resulted in a complex, fragmented, and intertwined physical
 22 and governmental infrastructure. Although primary responsibility might be assigned to a specific local
 23 entity, aggregate responsibilities are spread among more than 85 agencies in the North Coast Hydrologic
 24 Region with many different governance structures. Agency roles and responsibilities can be limited by
 25 how the agency was formed, which might include enabling legislation, a charter, a memorandum of
 26 understanding with other agencies, or facility ownership.

27 The North Coast Hydrologic Region is the site of many flood management infrastructure including
 28 floodwater storage facilities and channel improvements funded and/or built by the State and federal
 29 agencies. Flood management agencies are responsible for operating and maintaining approximately 1,200
 30 miles of levees, more than 110 dams and reservoirs, and other facilities within the North Coast
 31 Hydrologic Region.

32 For a list of the entities that have responsibilities or involvement in flood and water resources
 33 management, and a list of major infrastructure, refer to California’s Flood Future Report Attachment E:
 34 Information Gathering Technical Memorandum. See <http://www.water.ca.gov/sfmp/> for more information
 35 on this report.

36 *Groundwater Governance*

37 California does not have a statewide management program or statutory permitting system for
 38 groundwater. However, one of the primary vehicles for implementing local groundwater management in
 39 California is a Groundwater Management Plan (GWMP). Some agencies utilize their local police powers
 40 to manage groundwater through adoption of groundwater ordinances. Groundwater management also

1 occurs through other avenues such as basin adjudication, IRWMPs, Urban Water Management plans, and
2 Agriculture Water Management plans.

3 **Groundwater Management Assessment**

4 Figure NC-21 shows the location and distribution of the GWMPs within the North Coast Hydrologic
5 Region based on a GWMP inventory developed through a joint DWR/Association of California Water
6 Agencies (ACWA) online survey and follow-up communication by DWR in 2011-2012. Table NC-18
7 furnishes a list of the same. GWMPs prepared in accordance with the 1992 AB 3030 legislation, as well
8 as those prepared with the additional required components listed in the 2002 SB 1938 legislation are
9 shown. Information associated with the GWMP assessment is based on data that was readily available or
10 received through August 2012. Requirements associated with the 2011 AB 359 (Huffman) legislation,
11 related to groundwater recharge mapping and reporting, did not take effect until January 2013 and are not
12 included in the current GWMP assessment. Sonoma County is split between the North Coast and San
13 Francisco hydrologic regions. The GWMP for the Sonoma County Water Agency is presented in the
14 regional report of the San Francisco Hydrologic Region.

15 **PLACEHOLDER Figure NC-21 Location of Groundwater Management Plans in the North Coast** 16 **Hydrologic Region**

17 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
18 the end of the report.]

19 **PLACEHOLDER Table NC-18 Groundwater Management Plans in the North Coast Hydrologic** 20 **Region**

21 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
22 the end of the report.]

23 The GWMP inventory indicates that four groundwater management plans exists within the region. Two of
24 the GWMPs are fully contained within the North Coast Hydrologic Region, while the other two plans
25 include portions of the adjacent Sacramento River Hydrologic Region. Three of the four GWMPs cover
26 areas overlying Bulletin 118-2003 alluvial groundwater basins. However, two plans also include
27 management areas that extend beyond Bulletin 118-2003 alluvial basins. Collectively, the four GWMPs
28 cover 90 square miles. This includes about 6 square miles (one percent) of the Bulletin 118-2003 alluvial
29 groundwater basin area in the region. All four GWMPs have been developed or updated to include the SB
30 1938 requirements and are considered active for the purposes of the *California Water Plan Update 2013*
31 GWMP assessment. As of August 2012, none of the eight basins identified as medium priority under the
32 CASGEM Basin Prioritization (see Table NC-3) were covered by an active GWMP. These eight medium
33 priority basins account for about 60 percent of the population and about 80 percent of groundwater supply
34 in the region. Some efforts are under way to develop additional GWMPs in the region, but additional
35 efforts are needed to develop and implement California Water Code-compliant GWMPs.

36 Based on the information compiled through inventory of the GWMPs, an assessment was made to
37 understand and help identify groundwater management challenges and successes in the region, and
38 provide recommendations for improvement. Information associated with the GWMP assessment is based
39 on data that were readily available or received through August 2012 by DWR. The assessment process is
40 briefly summarized below.

1 The California Water Code §10753.7 requires that six components be included in a groundwater
 2 management plan for an agency to be eligible for State funding administered by DWR for groundwater
 3 projects, including projects that are part of an integrated regional water management program or plan
 4 (Table NC-19). Three of the components also contain required subcomponents. The requirement
 5 associated with the 2011 AB 359 (Huffman) legislation, applicable to groundwater recharge mapping and
 6 reporting, did not take effect until January 2013 and was not included in the current GWMP assessment.
 7 In addition, the requirement for local agencies outside of recognized groundwater basins was not
 8 applicable for any of the GWMPs in the region.

9 **PLACEHOLDER Table NC-19 Assessment for SB 1938 GWMP Required Components, SB 1938**
 10 **GWMP Voluntary Components, and Bulletin 118-03 Recommended Components**

11 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 12 the end of the report.]

13 In addition to the six required components, Water Code §10753.8 provides a list of twelve components
 14 that may be included in a groundwater management plan (Table NC-19). Bulletin 118-2003, Appendix C
 15 provides a list of seven recommended components related to management development, implementation,
 16 and evaluation of a GWMP, which should be considered to help ensure effective and sustainable
 17 groundwater management plan (Table NC-19).

18 As a result, the GWMP assessment was conducted using the following criteria:

- 19 • How many of the post SB 1938 GWMPs meet the six required components included in SB
- 20 1938 and incorporated into California Water Code §10753.7?
- 21 • How many of the post SB 1938 GWMPs include the twelve voluntary components included in
- 22 California Water Code §10753.8?
- 23 • How many of the implementing or signatory GWMP agencies are actively implementing the
- 24 seven recommended components listed in DWR Bulletin 118 - 2003?

25 In summary, assessment of the GWMPs in the North Coast Hydrologic Region indicates the following:

- 26 • None of the four GWMPs adequately address all of the required components listed under Water
- 27 Code §10753.7; one plan that fails to meet all the required components, does not address the
- 28 Basin Management Objective (BMO) and Monitoring Protocol subcomponents for surface
- 29 water-groundwater interaction. Analysis of the GWMPs for other regions also reveals that
- 30 when a plan lacks BMO details for surface water and groundwater interaction, it generally lacks
- 31 details for Monitoring Protocols as well.
- 32 • One of the four GWMPs incorporates the 12 voluntary components listed in Water Code
- 33 §10753.8; one plan incorporates 10 of the voluntary components;, and the remaining two plans
- 34 incorporate five or fewer of the voluntary components.
- 35 • One of the four GWMPs includes all seven components, and the other three plans include five
- 36 or fewer of the seven components recommended in Bulletin 118-03.

37 The DWR/ACWA survey asked respondents to identify key factors that contributed to the successful
 38 implementation of the agency's GWMP. Only one agency from the region participated in the survey. The
 39 single survey respondent identified data collection and sharing, developing an understanding of common
 40 interest, sharing of ideas and information, broad stakeholder participation, and having adequate surface

1 water supplies as key factors for successful GWMP implementation. Having adequate funding and the
2 time necessary to develop the GWMP were also identified as important factors.

3 Survey participants were also asked to identify factors that impeded implementation of the GWMP. The
4 single survey respondent pointed to a lack of adequate funding as an impediment to GWMP
5 implementation. Funding is a challenging factor for many agencies because the implementation and the
6 operation of groundwater management projects typically are expensive and because the sources of
7 funding for projects typically are limited to either locally raised monies or to grants from State and federal
8 agencies. Limited access to planning tools and unregulated groundwater pumping were also identified as
9 factors that impede successful implementation of GWMPs.

10 Finally, the survey asked if the respondents were confident in the long-term sustainability of their current
11 groundwater supply. The single respondent felt long-term sustainability of their groundwater supply was
12 possible.

13 The responses to the survey are furnished in Tables NC-20 and NC-21. *More detailed information on the*
14 *DWR/ACWA survey and assessment of the GWMPs are available online from Update 2013 Volume 4,*
15 *Reference Guide — California’s Groundwater Update 2013.*

16 **PLACEHOLDER Table NC-20 Factors Contributing to Successful Groundwater Management Plan**
17 **Implementation in the North Coast Hydrologic Region**

18 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
19 the end of the report.]

20 **PLACEHOLDER Table NC-21 Factors Limiting Successful Groundwater Management Plan**
21 **Implementation in the North Coast Hydrologic Region**

22 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
23 the end of the report.]

24 **Groundwater Ordinances**

25 Groundwater ordinances are laws adopted by local authorities, such as cities or counties, to manage
26 groundwater. In 1995, the California Supreme Court declined to review a lower court decision (*Baldwin*
27 *v. Tehama County*) that says that State law does not occupy the field of groundwater management and
28 does not prevent cities and counties from adopting ordinances to manage groundwater under their police
29 powers. Since 1995, the *Baldwin v. Tehama County* decision has remained untested; thus the precise
30 nature and extent of the police power of cities and counties to regulate groundwater is still uncertain.

31 There are a number of groundwater ordinances that have been adopted by counties in the region (Table
32 NC-22). The most common ordinances are associated with groundwater wells. These ordinances regulate
33 well construction, abandonment, and destruction; however, none of the ordinances provide for
34 comprehensive groundwater management.

35 **PLACEHOLDER Table NC-22 Groundwater Ordinances that Apply to Counties in the North Coast**
36 **Hydrologic Region**

37 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
38 the end of the report.]

1 **Special Act Districts**

2 Greater authority to manage groundwater has been granted to a few local agencies or districts created
 3 through a special act of the Legislature. The specific authority of each agency varies, but the agencies can
 4 be grouped into two general categories: (1) agencies having authority to limit export and extraction (upon
 5 evidence of overdraft or threat of overdraft) or (2) agencies lacking authority to limit extraction, but
 6 having authority to require reporting of extraction and to levy replenishment fees.

7 **Court Adjudication of Groundwater Rights**

8 Another form of groundwater management in California is through the courts. There are currently
 9 24 groundwater adjudications in California. The North Coast Hydrologic Region contains one of those
 10 adjudications (Table NC-23 and Figure NC-22). In Scott River watershed of the region, all surface water
 11 rights and much of the groundwater rights (excluding the tributaries below Scott Valley) have been
 12 adjudicated. In 1950, court decree was issued by Siskiyou County Superior Court for Shackelford Creek
 13 and then in 1958, for French Creek. The remainder of the valley's water claims (including some of the
 14 groundwater) was established in 1980 through the Scott River Stream System Decree. The California
 15 Water Code was amended in 1970 to allow the Scott River Stream System's surface water and supporting
 16 underflow and groundwater to be considered interconnected. As indicated in Table NC-23 and Figure
 17 NC-22, the Scott Valley groundwater basin in the region is included in this adjudication. To ensure that
 18 water rights distribution set forth in the adjudication is followed, watermaster service is provided by the
 19 Scott and Shasta Valley Watermaster District for distributing and monitoring groundwater pumping and
 20 surface water diversions according to the court decree. (State Water Resources Control Board 1980;
 21 Ellison, Schneider & Harris 2004).

22 **PLACEHOLDER Table NC-23 Groundwater Adjudications in the North Coast Hydrologic Region**

23 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 24 the end of the report.]

25 **PLACEHOLDER Figure NC-22 Groundwater Adjudications in the North Coast Hydrologic Region**

26 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 27 the end of the report.]

28 A landmark case focusing on water management and regulation in Scott River Valley is currently being
 29 reviewed by the courts and has the potential to significantly alter the way groundwater is managed in
 30 other parts of California. A 2010 lawsuit (*Environmental Law Foundation, et al. v. State Water Resources*
 31 *Control Board and Siskiyou County*) alleges that the State and the county are not exercising their
 32 authority under the public trust doctrine to manage and regulate groundwater extractions which contribute
 33 to important base flows in the Scott River. The lawsuit claims that years of approving well drilling
 34 permits have seriously depleted the local aquifer, creating severe water depletion in the Scott River,
 35 which was once an important salmon-bearing tributary to the Klamath River and is still home to federally
 36 and State-protected coho salmon. The lawsuit focuses on the groundwater aquifer areas outside the
 37 interconnected groundwater–surface water zone identified in the 1980 adjudication. The courts have not
 38 yet ruled if the public trust doctrine applies to groundwater depletion and the effect it has on nearby
 39 surface water systems — which is a critically important issue factor for many of California's
 40 groundwater basins.

1 The pending lawsuit has great potential significance because the public trust doctrine has not previously
 2 been applied toward regulation and management of groundwater use, and “percolating” groundwater has
 3 not previously been subject to regulation by SWRCB. If successful, the lawsuit could result in precedent
 4 setting changes in the way groundwater is managed in California. If the State is required to take the public
 5 trust doctrine into account for allocation and use of interconnected surface water–groundwater resources,
 6 then many of California’s groundwater users could expect to see an increase in State management and
 7 regulation of groundwater, and increased oversight of local groundwater management practices.

8 **Other Groundwater Management Planning Efforts**

9 Groundwater management also occurs through other avenues such as IRWMPs, Urban Water
 10 Management plans, and Agriculture Water Management plans. Box NC-4 summarizes these other
 11 planning efforts.

12 **PLACEHOLDER Box NC-4 Other Groundwater Management Planning Efforts in the North Coast** 13 **Hydrologic Region**

14 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 15 the end of the report.]

16 *Surface Water Ambient Monitoring Program (SWAMP)*

17 Surface Water Ambient Monitoring Program (SWAMP) is a program administered by the State Water
 18 Board. SWAMP is tasked with assessing water quality in all of California’s surface waters. The program
 19 conducts monitoring directly and through collaborative partnerships; and provides numerous information
 20 products, all designed to support water resource management in California. SWRCB works on this
 21 program in cooperation with several statewide and local work groups including the Klamath Basin Water
 22 Quality Monitoring Coordination Group. Recent programs in the North Coast Region (Regional Work
 23 plans), as of the writing of this document, include the Russian River- Freshwater Beaches Program
 24 (2012), Water Quality Status and Trends (2012), Garcia River Watershed Condition Monitoring (2012),
 25 Toxicity in California Waters- North Coast Region (2012), and the Regional Work plan for 2006 and
 26 2007 (2007).

27 The Russian River, Redwood Creek (Humboldt County), and Klamath basins have long-term water
 28 quality data sets, which is necessary to evaluate water quality changes over time. The current SWAMP
 29 sampling will contribute to these data sets (California Environmental Protection Agency 2012).

30 **Current Relationships with Other Regions and States**

31 **Klamath Basin**

32 As shown on the region map (see Figure NC-1) the Klamath River Basin straddles the border with
 33 Oregon, such that water from the upper basin flows into Oregon and eventually returns to California
 34 above Iron Gate Reservoir. On the Oregon side of this interstate basin, two surface water diversions
 35 export an average of 29,600 af /yr from Klamath River tributaries into the Rogue River system in Oregon.
 36 The Klamath River Basin also receives a small amount of imported water (about
 37 2,000 af /yr) from the upper reaches of the Sacramento River Hydrologic Region through a canal called
 38 the North Fork Ditch within Shasta Valley in Siskiyou County.

1 The Klamath Basin Restoration Agreement and the Klamath Hydroelectric Settlement Agreement
2 (KHSA) are companion agreements between Klamath Basin tribes, irrigators, fishermen, conservations,
3 counties, states of Oregon and California, federal agencies, and dam owners. The agreements aim to
4 restore Klamath Basin fisheries and sustain local economies. They include removal of four dams in the
5 upper Klamath River, increased flows for fish; greater reliability of irrigation water deliveries,
6 reintroduction of salmon above the dams and into and above Upper Klamath Lake, investment in
7 comprehensive and coordinated habitat restoration, an electrical power program for basin farmers and
8 ranchers, mitigation to counties for the effects of dam removal, and investment in tribal economic
9 revitalization. The first dam is scheduled to be removed in 2020, pending CEQA and NEPA (California
10 Department of Water Resources, Statewide Agreements 2010).

11 **Trinity River**

12 The North Coast region exports a large volume of water from the upper reaches of the Trinity River into
13 the Sacramento River region through the USBR's CVP at Lewiston Dam and the Clear Creek Tunnel. In
14 1998, a wet year, Trinity River exports (by water year) were 851,610 af; in 2000, an above normal water
15 year, 1.110 maf; and in 2001, a dry year, 670,530 af showing the variability of flows related to changing
16 hydrology. In contrast, when looking at flows for years since the ROD was implemented (see "Trinity
17 River Watershed" under "Settings" section of this document), in 2006, a wet year, exports were 1.353
18 maf, in 2008, a critical dry year, exports were 555,929 af and in 2010, a below normal water year,
19 275,202 af. These examples show how hydrology plays an important part in the decision of how much
20 water to export. However, current year hydrology is only part of the decision. Instream requirements for
21 fisheries downstream on the Trinity River, past year hydrology, current year estimated hydrology, water
22 quality concerns in the Delta and Trinity River, reservoir levels and operational needs are all considered
23 when setting export quantities (U.S. Geological Survey c2012).

24 The Trinity River Restoration Program was founded in 2000, based on three comprehensive foundational
25 documents: the landmark TRFESFR prepared by the USFWS and the Hoopa Valley Tribe in consultation
26 with the USGS, USBR, NMFS, and DFW (U.S. Fish and Wildlife Service and Hoopa Valley Tribe 1999);
27 the Trinity River Environmental Impact Statement (TREIS) prepared by USFWS et al. 2000; and the U.S.
28 Department of the Interior Record of Decision 2000.

29 The program is administered by the USBR and USFWS, both bureaus of the U.S. Department of the
30 Interior, as co-leads. Other partner agencies make up and share in the decision-making process of the
31 Trinity Management Council: the Hoopa Valley Tribe (HVT), the Yurok Tribe (YT), Trinity County, the
32 California Resources Agency (consisting of DWR and DFW), USFS, and the NMFS.

33 The river was dammed, and most of the flow was diverted to the Sacramento Valley beginning in 1963, as
34 part of the Trinity River Division of the CVP. (As a note: The Trinity River Division Act of 1955
35 authorized the Trinity River Diversion or TRD.) The diverted water enters the Sacramento River near
36 Redding, California, and provides for a variety of uses such as agriculture, industry, drinking water,
37 recreation, electrical power generation, and habitat. According to the Trinity River Restoration Program
38 Annual Report (2011), in 1970 it was believed that this diversion of water to the CVP was causing a
39 population decline in the Trinity River fishery. Federal legislation at that time and in subsequent years has
40 called for a variety of protections to the river, including protection of pre-dam levels of fisheries and of
41 Native American tribal rights for access to Trinity River fish. For more information on the Trinity River
42 Watershed and Trinity River Division, see section on Setting; sub-section, Trinity River Watershed, in

1 this document. For further information concerning the Trinity River Restoration Program, go to
2 www.trrp.net (Trinity River Restoration Program 2012).

3 **Potter Valley Project**

4 The Russian River Basin began receiving Eel River water through the Potter Valley Project in 1908
5 (<http://www.pottervalleywater.org/history.html>) and with several modifications was diverting 154 taf /yr
6 into the basin. Communities grew up based upon the available supply in the augmented river system.
7 However, with the FERC relicensing and some lawsuits, the diversion has been cut 15 percent to
8 130.9 taf /yr.

9 Communities like Redwood Valley County Water District (RVCWD), are in an almost annual
10 summertime water shortage condition. In addition to diversion changes for the Potter Valley Project,
11 2007 through 2010 were low water years. RVCWD gathered most of the attention, but several small
12 community service districts and county water districts began having severe water supply problems. The
13 loss of supply also affected the reliability of SCWA to meet its demands, which affected supplies into the
14 San Francisco Bay Region.

15 **Sonoma-Petaluma Aqueduct**

16 In the most southern part of the region, a smaller export of roughly 33,000 af /yr is transported from the
17 lower Russian River into the northern portion of the San Francisco Bay Region through the Sonoma-
18 Petaluma Aqueduct, to supply communities in northern Marin County and southern Sonoma County. For
19 more information on the Petaluma Aqueduct, see section on "Project Operations," "Petaluma Aqueduct"
20 within this document.

21 **Regional Water Planning and Management**

22 The focus of regional planning activities varies significantly from north to south across the North Coast
23 region because of the diversity of water issues and involved water agencies. In the far north interstate
24 Klamath River watershed, much of the planning is being done by federal agencies such as the USBR, the
25 Natural Resources Conservation Service, and the USFWS, among others. These federal agencies are
26 working to balance the needs of the federal Klamath Project with water for fish, tribal interests, and
27 interests of communities affected by the federal project. Planning and issue resolution for the Trinity
28 River also have a significant federal lead role because of the federal CVP at Trinity and Lewiston
29 reservoirs. In general, many of the Northern California counties lack funding at the level available to
30 federal agencies to conduct regional planning.

31 In the central portion of the region, the communities and water issues in Humboldt, Trinity, and
32 Mendocino counties tend to be organized at the local or county levels, partly because these areas are
33 geographically separated from other developed regions. Planning activities of Humboldt Bay Municipal
34 Water District and the Humboldt County general plan update are one of the primary forums for regional
35 planning for the Arcata and Eureka areas. The Mendocino Council of Governments and the Mendocino
36 Community Services District are among the lead water planning agencies for the county, which includes
37 Ukiah and portions of the upper Russian River wine country.

38 Sonoma County is the southernmost county in the North Coast Hydrologic Region, and water planning is
39 closely associated with those of the adjoining San Francisco Bay region. Water planning is strongly

1 focused on meeting the urban needs of Santa Rosa and the surrounding communities served by SCWA.
 2 The agency coordinates with and is a member of several San Francisco Bay area regional planning
 3 groups, such as the Bay Area Water Agencies Coalition that provides significant direction and guidance
 4 for regional planning. Much of Sonoma County regional planning also focuses on the competing uses of
 5 the Russian River, which is the largest river in this part of the North Coast region. The Russian River
 6 Action Plan has been updated by SCWA, as a coordinated effort among federal, State, and local agencies
 7 to protect and restore salmonid fishery populations and habitat.

8 **Integrated Regional Water Management Coordination and Planning**

9 In the North Coast region, NCRWMG was formed to coordinate planning within the region. The
 10 NCRWMG is a consortium of counties working together on water management planning and project
 11 prioritization and implementation for the North Coast region. Currently the member counties of the
 12 NCRWMG are responsible for implementation of the NCIRWMP, with individual project proponents
 13 responsible for project implementation. More information about the authorizing resolutions for the
 14 existing institutional structure can be found at:
 15 http://www.northcoastirwmp.net/docManager/1000006298/NCIRWMP_Phase_I_2007.pdf “Authorizing
 16 Documentation and Eligible Applicant Documentation”.

17 You can read more about how the counties participate in the Regional Water Management Group at the
 18 same website. Some counties have expressed reservations about joining any collaborative planning effort
 19 that might conflict with their local authority. Please refer to Figure NC-23 for integrated regional
 20 management planning areas in the North Coast Region.

21 **PLACEHOLDER Figure NC-23 Integrated Regional Management Planning areas in the North Coast** 22 **Hydrologic Region**

23 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 24 the end of the report.]

25 **Accomplishments**

26 **Recent Initiatives or Actions to Improve Water Quality**

27 Since 2009, the North Coast RWQCB has been engaged in the following activities. Many of these
 28 activities support one or more of the 10 resource strategies developed by DWR with the primary objective
 29 of improving water quality. These activities also represent the North Coast RWQCB’s accomplishments
 30 for the period of 2009-2013.

- 31 1. Cleaning up and closing groundwater contamination sites at an accelerated rate.
- 32 2. Updating NPDES permits and Waste Discharge Requirements (WDRs). New permits incorpo-
 33 rate toxics rules and low-impact fevelopment techniques, where appropriate. Permits are also
 34 written to accommodate an increase in water recycling and water reuse, where possible. Non-
 35 municipal waste discharges typically regulated by NPDES permits in the North Coast include
 36 canneries, fish hatcheries, wineries and other food processing plants, groundwater cleanup pro-
 37 jects, hardboard manufacturing plants, pulp mills, sawmills, and gravel operations.
- 38 3. Implementing the statewide stormwater prevention regulations. Efforts include enrolling cities
 39 and other entities under the general stormwater permits and adopting individual stormwater

1 permits. A very large effort was made in the adoption of an MS4 permit* for the City of Santa
 2 Rosa.

3 *MS4 permit; Small Municipal Separate Storm Sewer System, The
 4 regulatory definition of an MS4 (40 CFR 122.26(b)(8)) is "a conveyance
 5 or system of conveyances (including roads with drainage systems,
 6 municipal streets, catch basins, curbs, gutters, ditches, man-made
 7 channels, or storm drains): (i) Owned or operated by a state, city, town,
 8 borough, county, parish, district, association, or other public body
 9 (created to or pursuant to state law) including special districts under state
 10 law such as a sewer district, flood control district or drainage district, or
 11 similar entity, or an Indian tribe or an authorized Indian tribal
 12 organization, or a designated and approved management agency under
 13 section 208 of the Clean Water Act that discharges into waters of the
 14 United States. (ii) Designed or used for collecting or conveying storm
 15 water; (iii) Which is not a combined sewer; and (iv) Which is not part of
 16 a Publicly Owned Treatment Works (POTW) as defined at 40 CFR
 17 122.2."
 18

19 In practical terms, operators of MS4s can include municipalities and
 20 local sewer districts, state and federal departments of transportation,
 21 public universities, public hospitals, military bases, and correctional
 22 facilities. The Storm water Phase II Rule added federal systems, such as
 23 military bases and correctional facilities by including them in the
 24 definition of small MS4s.

- 25 4. Continued monitoring of surface water quality trends at select locations around the region, as
 26 well as intensive watershed monitoring on a rotating schedule.
- 27 5. Evaluating available surface water data to identify impaired waters and schedule the develop-
 28 ment of a TMDL assessments and/or establish implementation measures to control known
 29 sources. A very large and complex TMDL was adopted for the Klamath River in 2010. Action
 30 Plans have been adopted for the Klamath River and the Lost River. An MOU was signed with
 31 the USFS for TMDL implementation efforts in the Scott and Salmon River watersheds. TMDL
 32 development efforts are underway in the Elk River, Freshwater Creek, Laguna de Santa Rosa,
 33 and Russian River. A Region-wide Temperature Implementation Policy is under development
 34 and will address identified temperature impairments in the Eel River, Mattole River, and Na-
 35 varro River.
- 36 6. Developing a program for controlling waste discharge from timber operations and other opera-
 37 tions on forested lands. Ownership-wide WDRs have been adopted for Mendocino Redwood
 38 Company and Green Diamond Resource Company. A waiver of WDRs has been developed for
 39 various activities on USFS lands. A general WDR for timber operation was adopted in 2004
 40 and a categorical waiver in 2009. Timber operators not otherwise covered by an ownership-
 41 wide permit or waiver must apply for coverage under either of the general programs — or ap-
 42 ply for an individual permit or waiver. Prior to this period, the RWQCB primarily addressed
 43 timber harvest related discharges on a THP-by-THP (Timber Harvest Plan) basis; thus, the new
 44 permit structure serves to more efficiently and effectively identify those operations requiring
 45 more or less oversight to insure the protection of water quality.

- 1 7. Protection of water quality from waste discharges associated with roads. Efforts include:
 - 2 A. Coordination with CAL FIRE on road-related Timber Harvest Rules;
 - 3 B. Development of a waiver of WDR for county road activities consistent with the fish-
4 friendly guidelines established by the five counties of Del Norte, Humboldt, Mendocino,
5 Trinity, and Siskiyou;
 - 6 C. Support of rural road closure, maintenance, and/or upgrade through the Mendocino County
7 Permit Coordination Program. This program supports a wide range of best management
8 practices related to erosion control and restoration, beyond road-related activities.
 - 9 D. Settlement agreement with Caltrans over discharges associated with the Confusion Hill by-
10 pass project.
- 11 8. Developing a program for the control of waste discharge from agricultural activities.
12 A Dairy Program was recently adopted by the Board (2012) in which the discharge of waste to
13 surface water will be controlled and shallow groundwater will be monitored to protect against
14 impacts from land application of dairy waste. A compliance program for irrigated agricultural
15 lands is also under development.
- 16 9. Ongoing enforcement activities to control water quality violations.
17

18 **Ecosystem Restoration**

19 Nearly 49 percent of the North Coast region is permanently protected as open space and includes parks,
20 reserves, recreation areas, national monuments, national forests, State forests, and other protected areas.
21 Over a million acres in the region have been designated as National Wilderness Areas. The North Coast
22 region also includes 21 areas listed as Critical Coastal Areas, 12 Marine Protected Areas, and 8 areas of
23 Special Biological Significance.

24 DFW recommends that priority be given to the following actions be taken in relation to water supply in
25 the North Coast region:

- 26 • Restoration projects that facilitate the improvement of aquatic habitat, including deep and
27 shallow open water;
- 28 • Actions that will offset, mitigate-for, or accommodate climate change-related environmental
29 issues such as sea water rise, temperature shifts, potential regime changes, etc.;
- 30 • Acquisition of conservation easements on lands;
- 31 • Protect or restore fish habitat through the improvement of fish passage conditions, gravel
32 augmentation, hydrology, fish screens, min/max flow, etc.;
- 33 • Development, collection and publication of instream flow data, including recommended
34 instream flow levels and minimum instream flow requirements;
- 35 • Prevent or reduce negative impacts from invasive non-native species including those associated
36 with water supply and conveyance projects such as quagga and zebra mussels, *Egeria densa*
37 (Dense waterweed, Brazilian Waterweed, Elodea), water hyacinth, and others;
- 38 • Restoration projects that facilitate the increase of populations and improvement of habitat for
39 salmon, especially coho;
- 40 • Restoration projects that improve upon existing wetlands, or create new wetlands in appropriate
41 areas;
- 42 • Improvements in the transparency and availability of environmental data;
- 43 • Acquisition of water for wildlife areas to assure health of the area;

- 1 • Water quality improvements (sediment, oxygen saturation, pollution, temperature, etc.) to
- 2 support healthy ecosystems;
- 3 • Improvements in coordination, management and implementation of watersheds.

4 Restoration efforts that support or are undertaken in conjunction with projects related to water supply
5 contribute to the protection and sustainability of ecosystems in the region. Presently, there are many
6 efforts to restore ecosystems in the region; to list them all is beyond the scope of this regional report. This
7 section describes a few representative projects that are being implemented in the region. They are notable
8 in that they are collaborative undertakings, involving State, federal, local agencies, and communities in
9 the North Coast region.

10 *Sage Steppe Ecosystem Restoration*

11 Restoration efforts in the upper Klamath Basin include the eradication of juniper within the sage steppe
12 ecosystem and associated vegetative communities of northeastern California. The effort began with a
13 series of information discussions between the Modoc National Forest, the BLM and local resource
14 agencies in the region. In April of 2008, the final EIS was issued for the Sage Steppe Ecosystem
15 Restoration Strategy. The restoration strategy EIS affects Modoc, Lassen, Shasta, and Siskiyou counties
16 as well as a portion of Washoe County in Nevada.

17 The action was undertaken because of the loss of sage steppe ecosystem processes and vegetation
18 conditions where the density of western juniper has created a shift in dominant vegetation in the region.
19 The purpose of the restoration strategy is to improve watershed function and condition, restore
20 biodiversity and productivity, manage fire fuel loads, and to implement where appropriate national
21 renewable energy directives. Projects have been completed recently to implement this strategy. A similar
22 effort is under way in southern Oregon as well.

23 *Klamath Basin National Wildlife Refuge Complex*

24 The Klamath Basin National Wildlife Refuge Complex is a wildlife refuge operated by the USFWS
25 located in the Klamath Basin in southern Oregon and Northern California. The complex consists of Lower
26 Klamath NWR (National Wildlife Refuge), Clear Lake NWR, Upper Klamath NWR, Tule Lake NWR,
27 Klamath Marsh NWR, and the Bear Valley NWR. Klamath Basin habitats include freshwater marshes,
28 open water, grassy meadows, coniferous forests, sagebrush grasslands, agricultural lands, and rocky cliffs
29 and slopes. These habitats support large numbers of resident and migratory wildlife. The refuge also
30 serves as a major stopping point for fall concentrations of Pacific Flyway waterfowl. See the following
31 section, "River Restoration - Klamath River" for information relating to the effect of the Klamath Basin
32 Restoration Agreement on the refuges.

33 **PLACEHOLDER Photo Geese and Mount Shasta as seen from the Klamath Basin National Wildlife** 34 **Refuge**

35 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
36 the end of the report.]

1 **River Restoration**

2 *Klamath River*

3 The Klamath Basin Restoration Agreement (KBRA) when implemented contains as its name implies,
4 strategies for restoring the fisheries and associated habitats for the Klamath River watershed. The
5 agreement is the result of a collaborative effort of a large group of stakeholders who have worked
6 together to find solutions to water conflicts in the region. The plan was adopted in January 2010 and will
7 implement fisheries restoration with the removal of four dams that were constructed in the early 1900s as
8 part of USBR's Klamath Reclamation Project.

9 The KBRA is intended to result in effective and durable solutions which:

- 10 10. In concert with the removal of four dams, will restore and sustain natural production and pro-
11 vide for full participation in ocean and river harvest opportunities of fish species throughout the
12 Klamath Basin;
- 13 11. Establish reliable water and power supplies which sustain agricultural uses, communities, and
14 National Wildlife Refuges; and
- 15 12. Contribute to the public welfare and the sustainability of all Klamath Basin communities.

16 According to the agreement, the dam removal would begin in 2020. Although the agreement has been
17 adopted, there are controversial issues and highly charged reactions to this plan.

18 *Shasta and Scott Rivers*

19 During the past 20 years, extensive restoration has been completed by the Shasta Resource Conservation
20 District and Coordinated Resource Management Program in the Shasta Valley and by the Siskiyou RCD
21 and Watershed Council in the Scott Valley. There are also water trusts in both valleys with Scott River
22 Water Trust beginning in 2007 and Shasta River Water Trust beginning in 2012:

23 Every water diversion accessible by coho has a fish screen. Diversions have
24 headgates and most are managed by a watermaster. Ninety percent (plus) of the
25 main stem Scott River is fenced to keep cattle from entering the stream. There
26 have been numerous riparian plantings, bank stabilizations and hundreds of
27 projects on both rivers. As part of the Five County Salmonid Conservation
28 program, hundreds of miles of barriers to fish passage have been removed; road
29 culverts and conditions have been inventoried and treated to improve overall
30 habitat and migration (Scott River Water Trust 2013; Shasta River Water Trust
31 2013; Shasta River Water Association 2013; Five Counties Salmonid
32 Conservation Program 2013).

33 *Salt River*

34 The Salt River Ecosystem Restoration Project is a collaborative effort to restore fish habitat, improve
35 water quality, and provide for flood protection. The project affects restoration of the Salt River, Francis
36 Creek, and Williams Creek near the City of Ferndale in Humboldt County. Sediment monitoring is also
37 conducted to provide guidance on how much suspended sediment can be expected to enter the Salt River
38 from Francis Creek watershed. The data will be used to enhance sediment routing and provide planning
39 data for future dredging downstream. The project is considered to be of an ecosystem scale that includes
40 the restoration of a large tidal wetland that will create a succession of biologically rich and diverse tidal
41 wetland habitats, including transitional wetlands and adjacent uplands as part of a sustainable estuary

1 system. The mission of the project is to restore natural hydrologic function to the Salt River for the
2 improvement of water quality, wastewater treatment, flood control, wetlands and fisheries enhancement.

3 *Big River*

4 The Big River Program undertaken by Mendocino Land Trust and California State Parks seeks to provide
5 permanent protection of the estuarine, wetlands, wildlife, and associated seral-stage forest of the Big
6 River Units of the Mendocino Headlands State Park. Activities that contribute to these goals include
7 invasive plant control, greenhouse development for seed collection, trails and road monitoring, research
8 and resource monitoring, outreach and education.

9 In 2002, most of the Big River Estuary and some associated upland areas were added to the California
10 State Park System. The Big River Parcel consists of 7,334 acres, which when added to the surrounding
11 State Park system creates a 74,000-acre wildlife corridor linking coastal and inland habitats into the
12 largest piece of connected public land contained entirely within Mendocino County.

13 Coho, steelhead, and Chinook inhabit the Big River watershed, but population numbers are low compared
14 to historical levels. The estuary and lower river provide critical habitat for spawning, rearing, and staging
15 for adult, juvenile, and smolting salmonids.

16 *Salmon Creek*

17 Another collaborative effort to address the decline of salmonid runs on the North Coast includes
18 restoration projects on Salmon Creek in Sonoma County. This restoration project provides for the
19 instream placement of large woody debris at critical locations in the Salmon Creek estuary. Post-
20 construction monitoring on a similar project on the Mattole River indicated high utilization by juvenile
21 salmonids and lower water temperatures contributing to project success.

22 *Russian River*

23 The Russian River watershed encompasses 1,485 square miles (approx. 950,000 acres) within Sonoma
24 and Mendocino counties. The USACE Russian River Ecosystem Restoration study will look at
25 opportunities to prevent or reduce flood damages, to restore riverine ecosystem values and the wise use of
26 floodplains, to restore watershed functions through restorative land-use practices, and to conserve
27 remaining hydrologic and ecological resources. The result of Phase I was the formation of the Russian
28 River Watershed Council with the mission to protect, restore, and enhance the biological health of the
29 Russian River and its watershed through a community-based process by facilitating communication and
30 collaboration among all interested parties. The Plan of Action for Phase II (POA) articulates critical issues
31 and potential actions and can be found at Web site.

32 http://www.krisweb.com/biblio/russian_scwa_scwa_2002_actionplan.pdf. Phase II will include the
33 completion of a Russian River Watershed Adaptive Management Plan (WAMP). The WAMP Synthesis
34 Report was completed to provide the watershed community with a catalog of existing data and a ranking
35 of over 1,800 watershed areas in the Russian River watershed.

- 36 • 2009 accomplishments: Completion of the WAMP Synthesis Report, Task 1. The USACE
37 collaborated with Mendocino County RCD to incorporate Synthesis Report into Task 2, the
38 Draft Russian River Watershed Adaptive Management Plan.

- 2010 accomplishments: With additional funding, continue work on the Draft Russian River WAMP and begin work on the Implementation Plan, and the Monitoring Plan.

Laguna de Santa Rosa

The Laguna de Santa Rosa (a tributary to the Russian River and a subset of the Russian River watershed), in Sonoma County is a biologically rich freshwater wetland complex that has retained much of its wildland character even as its surrounding neighborhoods have been converted to agriculture, commerce, and housing. The "laguna" has remained relatively strong and resilient in the face of severe pressures from habitat fragmentation, water pollution, floodplain encroachment, and urban development. Meanwhile, the general public perception of the area as a "wetlands jewel" has resulted in a widespread outpouring of public sentiment in support of its protection and restoration.

But a deeper look at the wetlands reveals a long list of ecological imbalances that portend a darker future. The need for enhancing the laguna becomes clearer when the historical record is examined — most notably the record of the land's great fertility and its former abundance of wildlife and diversity of plant life. When compared to today's remaining, simpler, less-diverse, plant and animal communities, the contrast is sharp.

Enhancing the laguna by removing invasive plants, planting native plants, re-contouring human-made water channels, and reducing water pollutants is a fundamental goal of the area's citizens. Caring for the laguna includes monitoring for changes, stewarding the land, educating the generations, studying the ecological processes of the laguna, and enacting public policy. Restoring and managing the laguna are complementary sets of activities that together will strengthen its ability to reach a balanced state of flux and resiliency.

Mattole River

Restoration efforts on the Mattole River include the replacement of poorly designed and installed culverts to improve fish passage and stabilize sediment. The Mattole Integrated Water Management program is a watershed-wide effort to meet water supply, water quality, and fish habitat goals for the coastal Mattole River. Benefits of the project will include increased water supply in a drought-prone area, reduction in sediment load, invasive plant eradication, and riparian ecosystem restoration at 47 sites.

Trinity River Restoration Program

The Trinity River Restoration program is a collaborative effort of federal, State, tribal, and local stakeholders who are working together to restore the physical processes of the Trinity River as a foundation for the recovery of the fishery. Methods of restoration include the management of flows through releases from Lewiston Dam, construction of channel rehabilitation sites, spawning gravel augmentation, watershed projects to control fine sediments, infrastructure improvements, environmental compliance, and science based adaptive management. More information about the Trinity River can be found in the setting and watershed sections of this regional report.

Shasta River

Recent projects in the Shasta River area include projects that are designed to reduce agricultural tailwater runoff to the river. Other efforts are considering the feasibility of providing water users in the Shasta River watershed with an incentive-based approach that relieves certain regulatory pressures in exchange for leaving water instream to support the fishery.

1 Challenges

2 The region faces many water quality and water supply challenges. The North Coast RWQCB's water
3 quality priorities highlight the need for control of NPS runoff from logging, rural roads, agriculture, and
4 urban areas. In fact, sediment, temperature, and nutrients are the primary focus of the RWQCB's 303(d)
5 list of impaired water bodies. Along the coast, NPS pollution can cause microbial contamination of
6 shellfish growing areas, especially oysters. Much of the region is characterized by rugged, steep, forested
7 lands, with highly erodible, loosely consolidated soils; taken together with wildfires, extensive timber
8 harvesting, and heavy precipitation primarily in the form of rain, the watershed is highly susceptible to
9 erosion and landslides. Such heavy runoff in turn causes stream sedimentation that impacts habitat for
10 spawning and rearing of anadromous fish. Channel modifications and water diversions have radically
11 changed water-quality conditions in many water bodies in the region, reducing natural flows that dilute
12 contaminant concentrations and lessen their impacts. In the southern portion of the region, the
13 development of new hillside vineyards is an increasing source of erosion and pesticides.

14 Fisheries can be adversely affected by a number of factors related to both water quality and water
15 quantity. The Eel, Mad, Trinity, Klamath and Russian rivers, as well as many other streams, suffer from
16 sedimentation, which can smother salmonid spawning areas. The North Coast RWQCB Basin Plan sets
17 turbidity restrictions to control erosion impacts from logging and related activities, such as road building.
18 The basin plan also specifically establishes temperature objectives for the Trinity River, in which reduced
19 flows have disrupted temperature and physical cues for anadromous fish runs. Because of water
20 diversions, summer temperatures in the Trinity as well as the Klamath can be lethal to salmonids.
21 Fisheries can be further affected by the lack of woody debris for pool habitat and sediment metering.

22 The North Coast RWQCB Basin Plan requires tertiary treatment of wastewater discharges to the Russian
23 River, a major source of domestic water, and establishes limits on bacteriological contamination of
24 shellfish-growing areas along the coast. The plan also prohibits or strictly limits waste discharges to the
25 Klamath, Trinity, Smith, Mad, and Eel rivers, as well as estuaries and other coastal waters. NPS runoff,
26 especially after heavy precipitation, has resulted in contamination and closure of shellfish harvesting beds
27 in Humboldt Bay. In the lower Russian River watershed storm water runoff also might be contributing to
28 high ammonia and low dissolved oxygen levels in Laguna de Santa Rosa, which is threatening aquatic
29 life. Mercury in fish tissue is a water quality concern in Lakes Pillsbury, Mendocino, and Sonoma; a
30 health advisory for mercury has been issued for Lake Pillsbury.

31 Groundwater quality problems in the North Coast region include contamination from seawater intrusion
32 and nitrates in shallow coastal groundwater aquifers; high total dissolved solids and alkalinity in
33 groundwater associated with the lake sediments of the Modoc Plateau basins; and iron, boron, and
34 manganese in the inland groundwater basins of Mendocino and Sonoma counties. Septic tank failures in
35 western Sonoma County, at Monte Rio and Camp Meeker, and along the Trinity below Lewiston Dam,
36 are a concern because of potential impacts to groundwater wells and recreational water quality.

37 Other water quality concerns include the impacts of boating fuel constituents such as MTBE to
38 recreational water use at Trinity, Lewiston, and Ruth lakes. Abandoned mines, forest herbicide
39 application, and historical discharge of wood treatment chemicals at lumber mills, including Sierra Pacific
40 Industries near Arcata and Trinity River Lumber Company in Weaverville, are also regional issues of

1 concern. Of note, the Klamath basin, Redwood Creek watershed, and the Russian River basin all have
2 long-term water quality data sets, which are necessary to evaluate water quality changes over time.

3 Even though the North Coast region produces a substantial share of California’s surface water runoff,
4 only about 10 percent of this runoff occurs in the summer; and water supplies are limited throughout
5 much of the area. Small surface-water supply projects generally have limited carryover capacity that
6 cannot supply adequate water during extended months of low rainfall. The drinking water for many of the
7 communities on the North Coast, such as Klamath, Smith River, Crescent City, and most of the Humboldt
8 Bay area, is supplied by Ranney collectors (horizontal wells adjacent to or under the bed of a stream).

9 Erosion is undercutting some of these collectors, such as those in the Mad River supplying the Humboldt
10 Bay Municipal Water District (which serves Eureka, Arcata, and McKinleyville). As such, these “wells”
11 may actually be under the direct influence of surface water, which would then require filtration. The city
12 of Willits has had chronic problems with turbidity, taste, and odor with water from Morris Reservoir, and
13 high arsenic, iron, and manganese levels in its well supply. Organic chemical contamination has closed
14 municipal wells in the cities of Sebastopol and Santa Rosa. The town of Mendocino typifies the problems
15 related to groundwater development in the shallow marine terrace aquifers; surveys in the mid-1980s
16 indicate about 10 percent of wells go dry every year and up to 40 percent go dry during droughts.

17 The Klamath River Basin is an interstate watershed with surface storage facilities in both California and
18 Oregon, with competing water needs for agriculture, Indian tribal rights, waterfowl refuges, and
19 endangered fish. The primary water storage facilities belong to the federal Klamath Project, which is
20 operated by USBR, in conjunction with other dams and diversion structures operated by local irrigation
21 districts, wildlife management agencies, and electric power companies. In 2001, the lack of rainfall
22 generated a severe drought, which aggravated water disputes and caused harsh effects to agriculture,
23 waterfowl refuges and the downstream fisheries. The endangered fish populations include listed species
24 such as the Lost River and shortnose suckers, coho salmon, and steelhead trout. During 2001, USBR was
25 able to deliver only about 75,000 af of water to agriculture in California, which is about 25 percent of
26 normal. In the Tule Lake and Lower Klamath Lake subbasins, this translated to a drought disaster for both
27 agriculture and the wildlife refuges. In 2002, about 33,000 adult salmon died due to water quality and
28 quantity problems while trying to swim up the Klamath.

29 The Eel River and its tributaries are the largest river system draining to the coast of Humboldt County,
30 and it is characterized by significant water quality problems during winter storm events due to massive
31 sediment loads from unstable soils. The Eel River is also host to Humboldt County’s largest fisheries of
32 salmon and steelhead, which depend on access to upstream tributaries for spawning. The only major
33 water storage in the upper reaches of the Eel River is the Potter Valley Project, which consists of Lake
34 Pillsbury and a downstream diversion dam and tunnel to the Russian River (Mendocino County). The
35 project was originally built in 1908 by Snow Mountain Water and Power Company. Lake Pillsbury was
36 constructed in 1922 for hydropower production, and the project was acquired by Pacific Gas and Electric
37 Company in 1930.

38 In recent years, fishery interest groups have argued that the amount of water diverted to the Russian River
39 has adversely affected salmon and steelhead in the Eel River. The water needs of the Eel River fishery
40 have been evaluated and disputed during the recent FERC hydropower license amendment proceeding of
41 the Potter Valley Project. In June 2004, FERC approved PG&E’s relicense amendment of the Potter

1 Valley Project and its associated water diversions to the Russian River. However, fishery groups are
2 litigating the FERC decision, so the future distribution of project water between the Eel and Russian
3 rivers is not yet resolved.

4 **Flood Challenges**

5 Precipitation, coastline, terrain, and other area factors translate to frequent floods and flooding in the
6 North Coast region. Finding solutions to reduce residual flood risk in California is a complex task that
7 will require a mix of both old and new tools and approaches to flood management and funding, evolution
8 of existing planning processes and policies, sustained action, and commitment from agencies at all levels
9 to achieve the desired result of public safety, environmental stewardship, and financial stability in the
10 state. To accomplish these goals, the public, policymakers, and agencies at all levels must work together
11 to address the flood risk that exists statewide. Also, flood management practices must continue to evolve
12 toward integrated water management, and flood management agencies must be brought into the IRWM
13 process as full partners with other water management agencies. The hazards and risks of floods and
14 flooding are indiscriminate:

- 15 • People are exposed to flood risk. Flood hazard exposure is distributed throughout the state, with
16 all counties having some level of exposure. In the North Coast region, 30,000 people are
17 exposed to flood risk (5 percent of population) in a 100-year floodplain with 40,000 people (6
18 percent of population) exposed in a 500-year floodplain.
- 19 • Structures are at risk. Property and assets are exposed to flood hazards in all regions of
20 California. In the North Coast region, \$3 billion worth of structures (8 percent) are exposed in a
21 100-year floodplain with \$4 billion (10 percent) exposed in a 500-year floodplain.
- 22 • California's agricultural economy is at risk. A major flood event in California has the potential
23 to devastate regional agriculture based economies and cause serious impacts to the State
24 economy. In the North Coast region, \$80 million of crop value is exposed in a 100-year
25 floodplain (108,000 acres or 25 percent of crop acreage). Within a 500-year floodplain in the
26 North Coast region, \$90 million in crop value from 112,000 acres (26 percent of crop land) is
27 exposed.
- 28 • Native American tribal lands at risk. Within the North Coast Region, 5,748 acres of tribal lands
29 are at risk in the 500-year floodplain.
- 30 • State and federal sensitive species are exposed to flood hazard. Within the North Coast region,
31 203 species of plants and 117 species of animals are exposed to flood risk in both the 100-year
32 and 500-year floodplains.
- 33 • Climate change may impact flood hazard risk. Climate change could have a significant impact
34 on the timing and magnitude of runoff in California. In addition, increasing temperatures could
35 result in a rise in sea level, which likely would result in an increase in flood events. These
36 changes could result in expansions of the 100-year and 500-year floodplains, thereby causing
37 an increase in the people, property, and infrastructure exposed to flood hazards in the future.

38 **Conjunctive Management and Groundwater Storage**

39 Conjunctive management, or conjunctive use, refers to the coordinated and planned use and management
40 of both surface water and groundwater resources to maximize the availability and reliability of water
41 supplies in a region to meet various management objectives. Managing both resources together, rather
42 than in isolation, allows water managers to use the advantages of both resources for maximum benefit.
43 Conjunctive use of surface water and groundwater has been utilized for decades by numerous coastal and

1 inland basins throughout the North Coast Hydrologic Region. Some basin examples include Eureka Plain,
 2 Eel River Valley, Santa Rosa Valley, Smith River Plain, Wilson Grove, Big Valley, Tule Lake Valley,
 3 Scott Valley, and Shasta Valley. Many agencies have erected systems of barriers to allow more efficient
 4 percolation of ephemeral runoff from surrounding mountains.

5 A survey undertaken in 2011-2012 jointly by DWR and ACWA to inventory and assess conjunctive
 6 management projects in California is summarized in Box NC-5. *More detailed information about the*
 7 *survey results and a statewide map of the conjunctive management projects and operational information,*
 8 *as of July 2012, is available online Update 2013 Volume 4, Reference Guide — California’s*
 9 *Groundwater Update 2013.*

10 **PLACEHOLDER Box NC-5 Statewide Conjunctive Management Inventory Effort in California**

11 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 12 the end of the report.]

13 *Conjunctive Management Inventory Results*

14 Although 89 conjunctive management programs were identified in California as part of the DWR/ACWA
 15 survey and although incidental and planned conjunctive management is known to occur in many basins in
 16 the North Coast Hydrologic Region, no agencies in the region responded to the survey. The lack of survey
 17 response from agencies in the region could be due to confusion over what constitutes a conjunctive
 18 management program. Confusion of the terminology and meaning of conjunctive management is
 19 common. Additional information regarding conjunctive management in California as well as discussion
 20 on associated benefits, costs, and issues can be found online from Update 2013 Vol. 3 Chapter 9,
 21 “Conjunctive Management and Groundwater Storage Resource Management Strategy.”

22 **Drought Planning**

23 *Klamath Basin Restoration Agreement Drought Plan 2011*

24 In 2011, representatives from the State of California and Oregon, USBR, tribal organizations, and other
 25 stakeholders (Klamath Basin Coordinating Council) under Section 19.2 of the Klamath Basin Restoration
 26 Agreement developed a Drought Plan for the Upper Klamath Region. The Drought Plan identifies a
 27 number of strategies that would be used to counteract the effects of drought and extreme drought in the
 28 region. Measures that could be implemented include voluntary water conservations, additional
 29 stored water, the use of groundwater and the reduction of diversions (Klamath Basin Coordinating
 30 Council 2011).

31 **Looking to the Future**

32 *Future Conditions*

33 **Future Scenarios**

34 For Update 2013, the CWP evaluates different ways of managing water in California depending on
 35 alternative future conditions and different regions of the state. The ultimate goal is to evaluate how
 36 different regional response packages, or combinations of resource management strategies from Volume 3,
 37 perform under alternative possible future conditions. The alternative future conditions are described as
 38 future scenarios. Together, the response packages and future scenarios show what management options

1 could provide for sustainability of resources and ways to manage uncertainty and risk at a regional level.
2 The future scenarios are composed of factors related to future population growth and factors related to
3 future climate change. Growth factors for the North Coast are described below. Climate change factors
4 are described in general terms in Chapter 5, Volume 1.

5 *Water Conservation*

6 The CWP scenario narratives include two types of water use conservation. The first is conservation that
7 occurs without policy intervention (called background conservation). This includes upgrades in plumbing
8 codes and end user actions such as purchases of new appliances and shifts to more water efficient
9 landscape absent a specific government incentive. The second type of conservation expressed in the
10 scenarios is through efficiency measures under continued implementation of existing best management
11 practices in the Memorandum of Understanding (California Urban Water Conservation Council 2004).
12 These are specific measures that have been agreed upon by urban water users and are being implemented
13 over time. Any other water conservation measures that require additional action on the part of water
14 management agencies are not included in the scenarios, and would be represented as a water management
15 response.

16 *North Coast Growth Scenarios*

17 Future water demand in North Coast hydrologic region is affected by a number of growth and land use
18 factors, such as population growth, planting decisions by farmers, and size and type of urban landscapes.
19 See Table NC-24 for a conceptual description of the growth scenarios used in *California Water Plan*
20 *Update 2013*. The CWP quantifies several factors that together provide a description of future growth and
21 how growth could affect water demand for the urban, agricultural, and environmental sectors in North
22 Coast region. Growth factors are varied between the scenarios to describe some of the uncertainty faced
23 by water managers. For example, it is impossible to predict future population growth accurately, so the
24 CWP uses three different, but plausible population growth estimates when determining future urban water
25 demands. In addition, the CWP considers up to three different alternative views of future development
26 density. Population growth and development density will reflect how large the urban landscape will
27 become in 2050 and are used by the CWP to quantify encroachment into agricultural lands by 2050 in
28 North Coast region.

29 **PLACEHOLDER Table NC-24** Conceptual Growth Scenarios

30 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
31 the end of the report.]

32 For Update 2013, DWR worked with researchers at the University of California, Davis, to quantify how
33 much growth might occur in North Coast region through 2050. The UPlan model was used to estimate a
34 year 2050 urban footprint under the scenarios of alternative population growth and development density
35 (see <http://ice.ucdavis.edu/project/uplan> for information on the UPlan model). UPlan is a simple rule-
36 based urban growth model intended for regional or county-level modeling. The needed space for each
37 land use type is calculated from simple demographics and is assigned based on the net attractiveness of
38 locations to that land use (based on user input), locations unsuitable for any development, and a general
39 plan that determines where specific types of development are permitted. Table NC-25 describes the
40 amount of land devoted to urban use for 2006 and 2050, and the change in the urban footprint under each
41 scenario. As shown in the table, the urban footprint grew by about 20,000 acre under the low population

1 growth scenario (LOP) by 2050 relative to 2006 base-year footprint of about 190,000 acres. Urban
 2 footprint under the high population scenario (HIP), however, grew by about 90,000 acres. The effect of
 3 varying housing density on the urban footprint is also shown.

4 **PLACEHOLDER Table NC-25 Growth Scenarios (Urban) – North Coast**

5 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 6 the end of the report.]

7 Table NC-26 describes how future urban growth could affect the land devoted to agriculture in 2050.
 8 Irrigated land area is the total agricultural footprint. Irrigated crop area is the cumulative area of
 9 agriculture, including multicrop area, where more than one crop is planted and harvested each year. Each
 10 of the growth scenarios shows a decline in irrigated acreage over existing conditions, but to varying
 11 degrees. As shown in the table, irrigated crop acreage declines by about 40,000 acres by year 2050 as a
 12 result of low population growth and urbanization in North Coast region, while the decline under high
 13 population growth was slightly higher by about 50,000 acres.

14 **PLACEHOLDER Table NC-26 Growth Scenarios (Agricultural) – North Coast**

15 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 16 the end of the report.]

17 *North Coast 2050 Water Demands*

18 In this section, a description is provided for how future water demands might change under scenarios
 19 organized around themes of growth and climate change described earlier in this chapter. The change in
 20 water demand from 2006 to 2050 is estimated for the North Coast region for the agriculture and urban
 21 sectors under 9 growth scenarios and 13 scenarios of future climate change. The climate change scenarios
 22 included the 12 Climate Action Team scenarios described in Chapter 5, Volume 1 and a 13th scenario
 23 representing a repeat of the historical climate (1962-2006) to evaluate a “without climate change”
 24 condition.

25 Figure NC-24 shows the change in water demands for the urban and agricultural sectors under 9 growth
 26 scenarios, with variation shown across 13 climate scenarios. The nine growth scenarios include three
 27 alternative population growth projections and three alternative urban land development densities, as
 28 shown in Table NC-24. The change in water demand is the difference between the historical average for
 29 1998 to 2005 and future average for 2043 to 2050. Urban demand is the sum of indoor and outdoor water
 30 demand where indoor demand is assumed not to be affected by climate. Outdoor demand, however,
 31 depends on such climate factors as the amount of precipitation falling and the average air temperature.
 32 The solid blue dot in Figure NC-24 represents the change in water demand under a repeat of historical
 33 climate, while the open circles represent change in water demand under 12 scenarios of future climate
 34 change.

35 **PLACEHOLDER Figure NC-24 Change in North Coast Agricultural and Urban Water Demands for**
 36 **117 Scenarios from 2006-2050 (thousand af /yr)**

37 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 38 the end of the report.]

1 Urban demand increased under all nine growth scenarios tracking with population growth. On average, it
2 increased by about 20 taf under the three low population scenarios, 30 taf under the three current trend
3 population scenarios and about 100 taf under the three high population scenarios when compared to
4 historical average of about 150 taf. The results show change in future urban water demands are less
5 sensitive to housing density assumptions or climate change than to assumptions about future population
6 growth.

7 Agricultural water demand decreases under all but one of the future scenarios due to reduction in irrigated
8 lands as a result of urbanization and background water conservation when compared with historical
9 average water demand of about 750 taf. Under the three low population scenarios, the average reduction
10 in water demand was about 60 taf while it was about 85 taf for the three high population scenarios. For
11 the three current trend population scenarios, this change was about 65 taf. The results show that low
12 density housing would result in more reduction in agricultural demand since more lands are lost under
13 low-density housing than high density housing.

14 **Integrated Water Management Plan Summaries**

15 Inclusion of the information contained in IRWMP's into the CWP regional reports has been a common
16 suggestion by regional stakeholders at the regional outreach meetings since the inception of the IRWM
17 program. To this end, the CWP update has taken on the task of summarizing readily available IRWMPs in
18 a consistent format for each of the regional reports. This collection of information will not be used to
19 determine IRWM grant eligibility. This effort is ongoing and will be included in the final CWP updates
20 and will include up to four pages for each IRWMP in the regional reports.

21 In addition to these summaries being used in the regional reports, we intend to provide all of the summary
22 sheets in one IRWMP Summary "Atlas" as an article included in Volume 4. This atlas will, under one
23 cover, provide an "at-a-glance" understanding of each IRWM region and highlight each region's key
24 water management accomplishments and challenges. The atlas will showcase how the dedicated efforts of
25 individual regional water management groups (RWMGs) have individually and cumulatively transformed
26 water management in California.

27 All IRWMPs are different in how they are organized. Therefore, finding and summarizing the content in a
28 consistent way proved difficult. It became clear through these efforts that a process is needed to allow
29 those with the most knowledge of the IRWMPs — those who were involved in the preparation — to have
30 input on the summary. It is the intention that this process be initiated following release of *California*
31 *Water Plan Update 2013* and continue to be part of the process of the update process for *California Water*
32 *Plan Update 2018*. This process will also allow for continuous updating of the content of the atlas as new
33 IRWMPs are released or existing IRWMPs are updated.

34 As can be seen in Figure NC-23, there is one IRWM planning effort that is ongoing in the North Coast
35 Hydrologic Region.

36 **Placeholder Text:** At the time of the Public Review Draft the collection of information out of the
37 IRWMPs in the region has not been completed. Below are the basic types of information this effort will
38 summarize and present in the final regional report for each IRWMP available. An opportunity will be
39 provided to those with responsibility over the IRWMP to review these summaries before the reports are
40 final.

1 **Region Description:** This section will provide a basic description of the IRWM region. This would
2 include location, major watersheds within the region, status of planning activity, and the governance of
3 the IRWM. In addition, a IRWM grant funding summary will be provided.

4 **Key Challenges:** The top five challenges identified by the IRWM would be listed in this section.

5 **Principal Goals/Objective:** The top five goals and objectives identified in the IRWMP will be listed in
6 this section.

7 **Major IRWM Milestones and Achievements:** Major milestones (Top 5) and achievements identified in
8 the IRWMP would be listed in this section.

9 **Water Supply and Demand:** A description (one paragraph) of the mix of water supply relied upon in the
10 region along with the current and future water demands contained in the IRWMP will be provided in this
11 section.

12 **Flood Management:** A short (one paragraph) description of the challenges faced by the region and any
13 actions identified by the IRWMP will be provided in this section.

14 **Water Quality:** A general characterization of the water quality challenges (one paragraph) will be
15 provided in this section. Any identified actions in the IRWMP will also be listed.

16 **Groundwater Management:** The extent and management of groundwater (one paragraph) as described
17 in the IRWMP will be contained in this section.

18 **Environmental Stewardship:** Environmental stewardship efforts identified in the IRWMP will be
19 summarized (one paragraph) in this section.

20 **Climate Change:** Vulnerabilities to climate change identified in the IRWMP will be summarized (one
21 paragraph) in this section.

22 **Tribal Communities:** Involvement with tribal communities in the IRWM will be described (one
23 paragraph) in this section of each IRWMP summary.

24 **Disadvantaged Communities:** A summary (one paragraph) of the discussions on disadvantaged
25 communities contained in the IRWMP will be included in this section of each IRWMP summary.

26 **Governance:** This section will include a description (less than one paragraph) of the type of governance
27 the IRWM is organized under.

28 **Resource Management Strategies**

29 Volume 3 contains detailed information on the various resource management strategies that can be used
30 by water managers to meet their goals and objectives. A review of the resource management strategies
31 addressed in the available IRWMPs are summarized in Table NC-27.

1 **PLACEHOLDER Table NC-27 Resource Management Strategies addressed in IRWMPs in the North**
 2 **Coast Hydrologic Region**

3 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
 4 the end of the report.]

5 *Regional Resource Management Strategies*

6 The following are the resource management strategies identified by DWR with great potential to benefit
 7 water quality in the North Coast Hydrologic Region.

- 8 1. Agricultural Water Use Efficiency
- 9 2. Urban Water Use Efficiency
- 10 3. Conjunctive Management and Groundwater Storage, with the caveat that shallow groundwater
 11 use is of critical human and ecological importance in the North Coast region
- 12 4. Recycled Municipal Water
- 13 5. Groundwater and Aquifer Remediation, with the caveat that shallow groundwater use is of
 14 critical human and ecological importance in the North Coast region
- 15 6. Pollution Prevention
- 16 7. Urban Runoff Management
- 17 8. Agricultural Lands Stewardship
- 18 9. Ecosystem Restoration
- 19 10. Forest Management
- 20 11. Land Use Planning and Management
- 21 12. Recharge Areas Protection, with the caveat that shallow groundwater use is of crucial human
 22 and ecological importance in the North Coast region
- 23 13. Water-dependent Recreation
- 24 14. Watershed Management

25 The following are resource strategies identified by DWR that address issues of importance in the North
 26 Coast Hydrologic Region but may not accurately capture the issues as they express themselves on the
 27 North Coast.

- 28 1. Surface Storage — Regional/Local. Instream impoundments in the North Coast Hydrologic
 29 Region often alter the natural pattern and range of flows in a river, reduce a water body's
 30 assimilative capacity for other perturbations, and sometimes result in unintended water quality
 31 consequences (e.g., nuisance algal blooms, including the production of toxic algae; elevated
 32 temperatures; alteration of downstream sediment delivery and sorting, etc.). The RWQCB is
 33 supportive of efforts to provide off-channel storage for summer agricultural use as an alternative
 34 to summer instream withdrawals. But, the construction of instream impoundments is not viewed,
 35 in most cases, as supportive of water quality goals.
- 36 2. Flood Risk Management. The North Coast Hydrologic Region has experienced increased
 37 flooding as a result of several interacting factors. These include historical land uses that have
 38 resulted in massive deliveries of sediment to water bodies; alterations to channel form and
 39 hydrology via roads, dams, armoring, and loss of riparian and floodplain habitat; reduction in
 40 baseflows due to surface and groundwater withdrawals; and increase in runoff rate and volume
 41 from landscape alterations. The RWQCB is supportive of efforts to address these causes of
 42 increased flood potential. The further reduction in natural hydrologic functioning via the
 43 construction of hardened flood control channels is not viewed, in most cases, as supportive of
 44 water quality goals.

1 **Climate Change**

2 For over two decades, the State and federal governments have been preparing for climate change effects
3 on natural and built systems with a strong emphasis on water supply. Climate change is already impacting
4 many resource sectors in California, including water, transportation and energy infrastructure, public
5 health, biodiversity, and agriculture (U.S. Global Change Research Program 2009, California Natural
6 Resources Agency 2009). Climate model simulations based on the Intergovernmental Panel on Climate
7 Change's 21st century scenarios project increasing temperatures in California, with greater increases in
8 the summer. Projected changes in annual precipitation patterns in California will result in changes to
9 surface runoff timing, volume, and type (Cayan 2008). Recently developed computer downscaling
10 techniques indicate that California flood risks from warm-wet, atmospheric river type storms may
11 increase beyond those that we have known historically, mostly in the form of occasional more-extreme-
12 than-historical storm seasons (Dettinger 2011).

13 Currently, enough data exist to warrant the importance of contingency plans, mitigation (reduction) of
14 greenhouse gas (GHG) emissions, and incorporating adaptation strategies; methodologies and
15 infrastructure improvements that benefit the region at present and into the future. While the State is taking
16 aggressive action to mitigate climate change through GHG reduction and other measures (California Air
17 Resources Board 2008), global impacts from carbon dioxide and other GHGs that are already in the
18 atmosphere will continue to impact climate through the rest of the century (Intergovernmental Panel on
19 Climate Change 2007).

20 Resilience to an uncertain future can be achieved by implementing adaptation measures sooner rather than
21 later. Because of the economic, geographical, and biological diversity of California, vulnerabilities and
22 risks from current and future anticipated changes are best assessed on a regional basis. Many resources
23 are available to assist water managers and others in evaluating their region-specific vulnerabilities and
24 identifying appropriate adaptive actions. (U.S. Environmental Protection Agency/California Department
25 of Water Resources 2011; California Emergency Management Agency/California Natural Resources
26 Agency 2012).

27 *Observations*

28 The region's observed temperature and precipitation vary greatly due to complex topography and relation
29 to the Pacific Ocean. Regionally specific air temperature data was retrieved through the Western Regional
30 Climate Center (WRCC). The WRCC has temperature and precipitation data for the past century.
31 Through an analysis of National Weather Service Cooperative Station and PRISM Climate Group gridded
32 data, scientists from the WRCC have identified 11 distinct regions across the state for which stations
33 located within a region vary with one another in a similar fashion. These 11 climate regions are used
34 when describing climate trends within the state (Abatzoglou et al. 2009). DWR's hydrologic regions,
35 however, do not correspond directly to WRCC's climate regions. A particular hydrologic region may
36 overlap more than one climate region and, hence, have different climate trends in different areas. For the
37 purpose of this regional report, climate trends of the major overlapping climate regions are considered to
38 be relevant trends for respective portions of the overlapping hydrologic region.

39 Locally in the North Coast region within the WRCC Northern Coastal climate region, mean temperatures
40 have increased by about 0.4 to 1.3 °F (0.2 to 0.7 °C) in the past century, with minimum and maximum
41 temperatures increasing by about 0.3 to 1.3 °F (0.2 to 0.7 °C) and 0.4 to 1.4 °F (0.2 to 0.8 °C),
42 respectively. Within the WRCC North Central climate region, mean temperatures have increased by about

1 0.5 to 2.8 °F (0.3 to 1.6 °C) in the past century, with minimum and maximum temperatures increasing by
2 about 1.2 to 2.1 °F (0.6 to 1.2 °C) and by 0.1 to 1.4 °F (0.0 to 0.8 °C), respectively. Within the WRCC
3 North East climate region, mean temperatures have increased by about 0.8 to 2.0 °F (0.5 to 1.1 °C) in the
4 past century, with minimum and maximum temperatures increasing by about 0.9 to 2.2 °F (0.5 to 1.2 °C)
5 and by 0.4 to 2.1 °F (0.2 to 1.2 °C), respectively (Western Regional Climate Center 2012). Mean annual
6 precipitation in Northern California has increased slightly in the 20th century, and precipitation patterns
7 in the region have considerable geographic and annual variation (California Department of Water
8 Resources 2006).

9 The Klamath River Basin has been affected by these climate trends with a decline in spring snowpack,
10 less precipitation falling as snow, and earlier snowmelt runoff (Knowles et al. 2007). Water year runoff
11 trends over the past century have increased in the Klamath, Salmon, Eel, and Russian River basins; the
12 largest increase was in the Eel River Basin with an additional 12 taf /yr more on average (California
13 Department of Water Resources 2006).

14 Historical sea level trends in this region are conflicting. A tide gage at North Spit, California, operating
15 since 1977, shows mean sea level (MSL) to be increasing at a rate equivalent to 1.55 feet (0.47 meters)
16 over the past century. A different tide gage at Crescent City, California, operating since 1933 shows MSL
17 to be decreasing at a rate equivalent to 0.21 feet (0.06 meters) over the past century (National Oceanic and
18 Atmospheric Administration 2012). Although we expect MSL to rise with climate change, MSL at
19 Crescent City is trending lower due to the Cascadia Subduction Zone, where the buildup of interseismic
20 strain is causing coastal uplift north of Cape Mendocino. Most gages south of Cape Mendocino show
21 relative sea-level rise, consistent with land subsidence. When adjusted for vertical land motions and for
22 atmospheric pressure effects, the rates of relative sea-level rise along the U.S. West Coast are lower than
23 the rate of global mean sea-level rise (National Research Council of the National Academies 2012).

24 Shifts in coastal fog patterns have been making conditions less favorable for coastal ecosystems. The
25 North Coast redwoods are currently experiencing drought stress under changing climate conditions
26 (Johnstone et al. 2010).

27 *Projections and Impacts*

28 While historical data is a measured indicator of how the climate is changing, it cannot project what future
29 conditions may be like under different GHG emissions scenarios. Current climate science uses modeling
30 methods to simulate and develop future climate projections. A recent study by Scripps Institution of
31 Oceanography uses the most sophisticated methodology to date and indicates that, by 2060-2069,
32 temperatures will be 3.4-4.9 °F (1.9-2.7 °C) higher across the state than they were from 1985 to 1994
33 (Pierce et al. 2012). Annual mean temperature of the North Coast region by 2060-2069 is projected to
34 increase by 3.4 °F (1.9 °C) for the WRCC Northern Coastal climate region, with increases of 2.7 °F
35 (1.5 °C) during the winter months and 4.3 °F (2.4 °C) during summer. The WRCC North Central climate
36 region has similar projections with annual mean temperatures increasing by 4.0 °F (2.2 °C), winter
37 temperatures increasing by 3.1 °F (1.7 °C), and summer temperatures increasing by 5.2 °F (2.9 °C). The
38 WRCC North East climate region projections have annual mean temperatures increasing by 4.7 °F
39 (2.6 °C), winter temperatures increasing by 3.4 °F (1.9 °C), and summer temperatures increasing by 6.5°F
40 (3.6 °C) Climate projections for this region, from Cal-Adapt indicate that temperatures between 1990 and
41 2100 will increase by 5 °F (2.8 °C) in the winter and 6 °F (3.3 °C) in the summer (California Emergency
42 Management Agency and California Natural Resources Agency 2012).

1 Changes in annual precipitation across California, either in timing or total amount, will result in changes
2 in type of precipitation (rain or snow) in a given area, and in surface runoff timing and volume. Most
3 climate model precipitation projections for the state anticipate drier conditions in Southern California,
4 with heavier and warmer winter precipitation in Northern California. More intense wet and dry periods
5 are anticipated, which could lead to flooding in some years and drought in others. In addition, extreme
6 precipitation events are projected to increase with climate change (Pierce et al. 2012). Because there is
7 less scientific detail on localized precipitation changes, there is a need to adapt to this uncertainty at the
8 regional level (Qian et al. 2010).

9 Climate model precipitation projections for Northern California are not all in agreement; simulated future
10 monthly average precipitation was found to be higher in the high Sierra and lower in the northern
11 drainage basins (Georgakakos et al. 2012). Rainfall and snowmelt dominated watersheds in the region
12 will each have a unique climate response and corresponding runoff, depending on the amount of warming
13 that occurs. With warmer temperatures and changes in precipitation patterns, the Klamath River Basin
14 may experience December-March runoff increases in streamflow and decreased April-June streamflow by
15 2100 (Markstrom et al. 2011).

16 While future precipitation and runoff is somewhat uncertain, greater flood magnitudes are anticipated as
17 more frequent atmospheric river storm events encounter the region. Recent computer downscaling
18 techniques indicate that California flood risks from warm-wet, atmospheric river type storms may
19 increase beyond those that we have known historically, mostly in the form of occasional more-extreme-
20 than-historical storm seasons (Dettinger 2011). These are periods of increased water vapor transported
21 toward the poles across the mid-latitudes within narrow, intense filamentary bands of moist air. A higher
22 proportion of precipitation falling as rain instead of snow and increased storm frequency will impact the
23 system's ability to provide effective flood protection.

24 Additionally, sea level is projected to continue to rise along California's coast. For the California coast
25 south of Cape Mendocino, the National Research Council (2012) projected that sea level will rise 1.5 to
26 12 inches (3.8 to 30 cm) by 2030, 4.5 to 24 inches (11.4 to 61 cm) by 2050, and 16.5 to 66 inches (41.9 to
27 168 cm) by 2100. For the Washington, Oregon, and California coast north of Cape Mendocino, sea level
28 is projected to change between falling 1.5 inches (3.8 cm) to rising 9 inches (23 cm) by 2030, falling 1
29 inch (2.5 cm) to rising 19 inches (48 cm) by 2050, and rising between 4 to 56 (10 to 142 cm) inches by
30 2100.

31 Projected climate changes are likely to upset the ecosystem balance, impacting sensitive fish and wildlife
32 species (Janetos et al. 2008). Warmer water temperatures will result in stress to fisheries, reducing
33 coldwater habitat for native species such as coho salmon, while potentially benefitting invasive species
34 such as quagga and zebra mussels. Increased water temperatures and nutrient loading will potentially
35 exacerbate toxic algae problems in the Klamath River with increases in extent, duration, toxicity, and
36 concentration of blue-green algal blooms (U.S. Bureau of Reclamation 2011b).

37 A further shift in coastal fog patterns along with temperature and precipitation changes may lead to range
38 shifts in vegetation. While a shift in vegetation patterns along the coast may decrease wildfire risk
39 (Lenihien et al. 2006), the non-coastal areas in the region will be at higher risk of wildfire (California
40 Natural Resources Agency 2012).

1 *Adaptation*

2 Climate change has the potential to impact the region, which California depends upon for its economic
3 and environmental benefits. These changes will increase the vulnerability of natural and built systems in
4 the region. Impacts to natural systems will challenge aquatic and terrestrial species with diminished water
5 quantity and quality, and shifting eco-regions. Built systems will be impacted by changing hydrology and
6 runoff timing, loss of natural snowpack storage, making the region more dependent on surface storage in
7 reservoirs and groundwater sources. Increased future water demand for both natural and built systems
8 may be particularly challenging with less natural storage and less overall supply.

9 Water managers and local agencies must work together to determine the appropriate planning approach
10 for their operations and communities. While climate change adds another layer of uncertainty to water
11 planning, it does not fundamentally alter the way water managers already address uncertainty (U.S.
12 Environmental Protection Agency and California Department of Water Resources 2011). However,
13 stationarity (the idea that natural systems fluctuate within an unchanging envelope of variability) can no
14 longer be assumed, so new approaches will likely be required (Milly et al. 2008).

15 IRWM planning is a framework that allows water managers to address climate change on a smaller, more
16 regional scale. Climate change is now a required component of all IRWM plans (California Department
17 of Water Resources 2010). IRWM regions must identify and prioritize their specific vulnerabilities and
18 identify adaptation strategies that are most appropriate for their sub-regions. Planning strategies to address
19 vulnerabilities and adaptation to climate change should be both proactive and adaptive, starting with
20 strategies that benefit the region in the present-day while adding future flexibility and resilience under
21 uncertainty.

22 Local agencies, as well as federal and State agencies, face the challenge of interpreting climate change
23 data and information to determine which adaptation methods and approaches are appropriate for their
24 planning needs. The *Climate Change Handbook for Regional Water Planning* (U.S. Environmental
25 Protection Agency and California Department of Water Resources 2011) provides an analytical
26 framework for incorporating climate change impacts into the regional and watershed planning process for
27 consideration of climate change. This handbook provides guidance for assessing the vulnerabilities of
28 California's watersheds and hydrologic regions to climate change impacts, and prioritizing these
29 vulnerabilities.

30 The primary water supply in the region is the Klamath, Eel and Russian River systems. With diminished
31 spring snowpack storage and very few significant aquifers, the potential for water supply shortages
32 increase. Agricultural water use efficiency and urban water use efficiency are resource management
33 strategies outlined in the CWP (see Volume 3) to adapt to water scarcity. These strategies would benefit
34 the region that has already developed most of its potential surface and groundwater supplies. Urban water
35 use efficiency focuses on conservation to lower municipal demand, and agriculture water use efficiency
36 helps the grower use water in a way that is most effective to the crop, while minimizing yield losses.

37 Many of the resource management strategies from *California Water Plan Update 2009* (Volume 2)
38 provide benefits for adapting to climate change in addition to meeting water management objectives.
39 These include:

- 40 1. Regional and local Conveyance
- 41 2. Conjunctive Management and Groundwater storage

- 1 3. Precipitation Enhancement
- 2 4. Regional and Local Surface Storage; Pollution Prevention
- 3 5. Ag Land Stewardship
- 4 6. Ecosystem Restoration
- 5 7. Forest Management
- 6 8. Land Use Planning and Management
- 7 9. Recharge Area Protection
- 8 10. Watershed Management
- 9 11. Integrated Flood Management

10 The myriad of resources and choices available to managers can seem overwhelming, and the need to take
 11 action given uncertain future conditions is daunting. However, there are many actions that water
 12 managers can take to prepare for climate change, regardless of the magnitude of future warming. These
 13 actions often provide economic and public health co-benefits. Water and energy conservation are
 14 examples of strategies that make sense with or without the additional pressures of climate change.
 15 Conjunctive management projects that manage surface water and groundwater in a coordinated fashion
 16 could provide a buffer against variable annual water supplies. Forecast-coordinated operations would
 17 provide flexibility for water managers to respond to weather conditions as they unfold.

18 Water managers will need to consider both the natural and built environments as they plan for the future.
 19 Stewardship of natural areas and protection of biodiversity are critical for maintaining ecosystem services
 20 important for human society such as carbon sequestration, pollution remediation, and habitat for
 21 pollinators. Increased cross-sector collaboration between water managers, land use planners, and
 22 ecosystem managers provides opportunities for identifying common goals and actions needed to achieve
 23 resilience to climate change and other stressors.

24 *Mitigation*

25 California's water sector has a large energy footprint, consuming 7.7 percent of statewide electricity
 26 (California Public Utilities Commission 2010). Energy is used in the water sector to extract, convey, treat,
 27 distribute, use, condition, and dispose of water. Figure 3-26, "Water-Energy Connection" in Volume 1,
 28 Chapter 3, "California Water Today" shows all of the connections between water and energy in the water
 29 sector; both water use for energy generation and energy use for water supply activities. Regional reports
 30 in *California Water Plan Update 2013* are the first to provide detailed information on the water-energy
 31 connection, including energy intensity (EI) information at the regional level. This EI information is
 32 designed to help inform the public and water utility managers about the relative energy requirements of
 33 the major water supplies used to meet demand. Since energy usage is related to GHG emissions, this
 34 information can support measures to reduce GHG's, as mandated by the State.

35 Figure NC-25 shows the amount of energy associated with the extraction and conveyance of one acre-foot
 36 of water for each of the major sources in this region. The quantity used is also included as a percent. For
 37 reference, Figure 3-26, "Water-Energy Connection" in Volume 1, Chapter 3, "California Water Today,"
 38 highlights which water-energy connections are illustrated in Figure NC-25 — only extraction and
 39 conveyance of raw water. Energy required for water treatment, distribution, and end uses of the water are
 40 not included. Not all water types are available in this region. Some water types flow by gravity to the
 41 delivery location and therefore do not require any energy to extract or convey (represented by a white
 42 light bulb).

1 **PLACEHOLDER Figure NC-25 Energy Intensity of Raw Water Extraction and Conveyance in the**
2 **North Coast Hydrologic Region**

3 [Any draft tables, figures, and boxes that accompany this text for the public review draft are included at
4 the end of the report.]

5 Recycled water and water from desalination used within the region are not show in Figure NC-25 because
6 their EI differs in important ways from those water sources. The EI of both recycled and desalinated water
7 depend not on regional factors but rather on much more localized, site, and application specific factors.
8 Additionally, the water produced from recycling and desalination is typically of much higher quality than
9 the raw (untreated) water supplies evaluated in Figure NC-25. For these reasons, discussion of EI of
10 desalinated water and recycled water are included in Volume 3, *Resource Management Strategies*.

11 EI, sometimes also known as embedded energy, is the amount of energy needed to extract and convey an
12 acre-foot of water from its source (e.g. groundwater or a river) to a delivery location, such as a water
13 treatment plant or a SWP delivery turnout. (Extraction refers to the process of moving water from its
14 source to the ground surface. Many water sources are already at ground surface and require no energy for
15 extraction, while others like groundwater or seawater for desalination require energy to move the water to
16 the surface. Conveyance refers to the process of moving water from a location at the ground surface to a
17 different location, typically but not always a water treatment facility. Conveyance can include pumping of
18 water up hills and mountains or can occur by gravity.) EI should not be confused with total energy — that
19 is, the amount of energy (e.g. kWh) required to deliver all of the water from a water source to customers
20 within the region. EI focuses not on the total amount of energy used to deliver water, but rather the energy
21 required to deliver a single unit of water (in kWh/acre-foot). In this way, EI gives a normalized metric
22 that can be used to compare alternative water sources.

23 In most cases, this information will not be of sufficient detail for actual project-level analysis. However,
24 these generalized, region-specific metrics provide a range in which energy requirements fall. The
25 information can also be used in more detailed evaluations using tools such as WeSim
26 (<http://www.pacinst.org/publication/wesim/>), which allows modeling of water systems to simulate
27 outcomes for energy, emissions, and other aspects of water supply selection. It's important to note that
28 water supply planning must take into consideration a myriad of different factors in addition to energy
29 impacts; costs, water quality, opportunity costs, environmental impacts, reliability, and other many other
30 factors.

31 EI is closely related to GHG emissions, but not identical, depending on the type of energy used (see
32 Volume 1, Chapter 3, "California Water Today," Water-Energy,). In California, generation of 1
33 megawatt-hour (MWh) of electricity results in the emission of about one-third of a metric ton of GHG,
34 typically referred to as carbon dioxide equivalent or CO₂e (eGrid 2012). This estimate takes into account
35 the use of GHG-free hydroelectricity, wind, and solar and fossil fuel sources like natural gas and coal.
36 The GHG emissions from a specific electricity source may be higher or lower than this estimate.

37 Reducing GHG emissions is a State mandate. Water managers can support this effort by considering EI
38 factors, such as those presented here, in their decision-making process. Water use efficiency and related
39 best management practices can also reduce GHGs (See Volume 3, *Resource Management Strategies*).

1 **Accounting for Hydroelectric Energy**

2 Generation of hydroelectricity is an integral part of many of the state’s large water projects. In 2007,
 3 hydroelectric generation accounted for nearly 15 percent of all electricity generation in California. The
 4 SWP, CVP, Los Angeles Aqueduct, Mokelumne Aqueduct, and Hetch Hetchy Aqueducts all generate
 5 large amounts of hydroelectricity at large multi-purpose reservoirs at the heads of each system. In
 6 addition to hydroelectricity generation at head reservoirs, several of these systems also generate
 7 hydroelectric energy by capturing the power of water falling through pipelines at in-conduit generating
 8 facilities (In-conduit generating facilities refer to hydroelectric turbines that are placed along pipelines to
 9 capture energy as water runs down hill in a pipeline [conduit]). Hydroelectricity is also generated at
 10 hundreds of smaller reservoirs and run-of-the-river turbine facilities.

11 Hydroelectric generating facilities at reservoirs provide unique benefits. Reservoirs like the SWP’s
 12 Oroville Reservoir are operated to build up water storage at night when demand for electricity is low and
 13 release the water during the day time hours when demand for electricity is high. This operation, common
 14 to many of the state’s hydropower reservoirs, helps improve energy grid stabilization and reliability and
 15 reduces GHG emissions by displacing the least efficient electricity generating facilities. Hydroelectric
 16 facilities are also extremely effective for providing backup power supplies for intermittent renewable
 17 resources like solar and wind power. Because the sun can unexpectedly go behind a cloud or the wind can
 18 die down, intermittent renewables need backup power sources that can quickly ramp up or down
 19 depending on grid demands and generation at renewable power installations.

20 Despite these unique benefits and the fact that hydroelectric generation was a key component in the
 21 formulation and approval of many of California’s water systems, accounting for hydroelectric generation
 22 in EI calculations is complex. In some systems like the SWP and CVP, water generates electricity and
 23 then flows back into the natural river channel after passing through the turbines. In other systems like the
 24 Mokelumne aqueduct water can leave the reservoir by two distinct outflows, one that generates electricity
 25 and flows back into the natural river channel and one that does not generate electricity and flows into a
 26 pipeline flowing into the East Bay Municipal Utility District service area. In both these situations, experts
 27 have argued that hydroelectricity should be excluded from EI calculations because the energy generation
 28 system and the water delivery system are in essence separate (Wilkinson 2000).

29 DWR has adopted this convention for the EI for hydropower in the regional reports. All hydroelectric
 30 generation at head reservoirs has been excluded from Figure NC-25. Consistent with Wilkinson (2000)
 31 and others, DWR has included in-conduit and other hydroelectric generation that occurs as a consequence
 32 of water deliveries, such as the Los Angeles Aqueduct’s hydroelectric generation at San Francisquito, San
 33 Fernando, Foothill, and other power plants on the system (downstream of the Owen’s River Diversion
 34 Gates). DWR has made one modification to this methodology to simplify the display of results: EI has
 35 been calculated at each main delivery point in the systems; if the hydroelectric generation in the
 36 conveyance system exceeds the energy needed for extraction and conveyance, the EI is reported as zero
 37 (0). That is, no water system is reported as a net producer of electricity, even though several systems do
 38 produce more electricity in the conveyance system than is used (e.g., Los Angeles Aqueduct, Hetch
 39 Hetchy Aqueduct). (For detailed descriptions of the methodology used for the water types presented, see
 40 Volume 5, *Technical Guide*).

1 **References**

2 References Cited

- 3 Agricultural Lands Discharge Program, 2012, North Coast Regional Water Quality Control Board,
4 California Environmental Protection Agency. Viewed online at:
5 http://www.waterboards.ca.gov/northcoast/water_issues/programs/agricultural_lands/. Accessed
6 on: Mar 20, 2013. Last update: Feb 28, 2013.
- 7 American Rivers, Mattole River. 1101 14th Street NW, Suite 1400, Washington, DC 20005, CA. Viewed
8 online at: <http://www.americanrivers.org/initiatives/dams/projects/mattole-river-restoration.html>.
9 Accessed: Aug 6, 2012. Last update: 2012.
- 10 Association of Bay Area Governments. 101 Eighth Street, Oakland, CA 94607, (510)464-7900. Viewed
11 online at: <http://www.abag.ca.gov/>. Accessed on: Aug 23, 2012. Last update: 2008.
- 12 Barker v. Harvey, 1901, U.S. Supreme Court. Viewed online at:
13 <http://caselaw.lp.findlaw.com/scripts/getcase.pl?court=US&vol=181&invol=481>. Accessed on:
14 Mar 13, 2013. Last update: May 13, 1901.
- 15 Bureau of Land Management, Modoc National Forest, Alturas Field Office, Modoc County, Sage Steppe
16 Ecosystem Restoration Strategy. Viewed online at:
17 [http://www.blm.gov/pgdata/etc/medialib/blm/ca/pdf/alturas.Par.67107.File.dat/sage%20steppe%20](http://www.blm.gov/pgdata/etc/medialib/blm/ca/pdf/alturas.Par.67107.File.dat/sage%20steppe%200rod.pdf)
18 [0rod.pdf](http://www.blm.gov/pgdata/etc/medialib/blm/ca/pdf/alturas.Par.67107.File.dat/sage%20steppe%200rod.pdf). Accessed; Feb 5, 2013. Last update; Dec 2008.
- 19 California Air Resources Board, AB 32 Climate Change Scoping Plan Document. 1001 "I" Street,
20 Sacramento, CA 95814. Viewed online at:
21 <http://www.arb.ca.gov/cc/scopingplan/document/scopingplandocument.htm>. Accessed: Feb 11,
22 2013. Last update: May 16, 2012.
- 23 California Department of Fish and Game (Now Department of Fish and Wildlife), Recovery Strategy for
24 California Coho Salmon, Species Recovery Strategy 2004-1. Viewed online at:
25 [http://www.dfg.ca.gov/fish/documents/SAL_SH/SAL_Coho_Recovery/ReportToCommission_20](http://www.dfg.ca.gov/fish/documents/SAL_SH/SAL_Coho_Recovery/ReportToCommission_2004/CohoRecoveryStrategy.pdf)
26 [04/CohoRecoveryStrategy.pdf](http://www.dfg.ca.gov/fish/documents/SAL_SH/SAL_Coho_Recovery/ReportToCommission_2004/CohoRecoveryStrategy.pdf). Accessed on: Oct 1, 2012. Last Update: Feb 2004.
- 27 California Department of Fish and Game, Suction Dredge Permitting Program. Viewed online at:
28 <http://www.dfg.ca.gov/suctiondredge/>. Accessed on: Jul 18, 2012. Last update on: Jul 11, 2012.
- 29 California Department of Food and Agriculture in cooperation with U.S. Department of Agriculture's
30 National Agricultural Statistics Service, Grape Acreage Report, 2010 Crop. California Field
31 Office, 650 Capitol Mall, Suite 6-100, P.O. Box 942871, Sacramento, CA 94721-001. Viewed
32 online at:
33 http://www.nass.usda.gov/Statistics_by_State/California/Publications/Grape_Acreage/201004gab
34 [tb00.pdf](http://www.nass.usda.gov/Statistics_by_State/California/Publications/Grape_Acreage/201004gab). Accessed on: Aug 13, 2012. Last update: Apr 2011.

- 1 California Department of Water Resources, California Drought Contingency Plan. Viewed online at:
2 http://www.waterplan.water.ca.gov/docs/meeting_materials/drought/2010.08.10/CA_Drought_Co
3 [ntingency_Plan-Public_Review_Draft-081010.pdf](http://www.waterplan.water.ca.gov/docs/meeting_materials/drought/2010.08.10/CA_Drought_Co). Accessed: Sept 28, 2012. Last update: Jun
4 2010.
- 5 California Department of Water Resources, Progress on incorporating Climate Change into Management
6 of California's Water Resources, Technical Memorandum Report, DWR Climate Change.
7 Viewed online at: <http://www.water.ca.gov/climatechange/docs/>. Accessed: Feb 5, 2013. Last
8 update: Jul 2006.
- 9 California Department of Water Resources, Managing an Uncertain Future, Climate Change Adaption
10 Strategies for California's Water. Viewed online at:
11 <http://www.water.ca.gov/climatechange/docs/ClimateChangeWhitePaper.pdf> . Accessed: Feb 5,
12 2013. Last update: Oct 2008.
- 13 California Department of Water Resources, Bulletin 73-79, Evaporation from Water Surfaces in
14 California, 163pp. Nov 1979.
- 15 California Department of Water Resources, California Data Exchange Center, Reservoir Water Storage.
16 Viewed online at: <http://cdec.water.ca.gov/cgi-progs/reservoirs/STORAGEW.11>. Accessed on:
17 Sept 24, 2012. Last update: Dec 2012.
- 18 California Department of Water Resources, California Data Exchange Center, Judge F Carr Power House.
19 Viewed online at: http://cdec.water.ca.gov/cgi-progs/staMeta?station_id=JCR. Accessed on: Sept
20 10, 2012. Last update: 2010.
- 21 California Department of Water Resources, California Irrigation Management Information System
22 (CIMIS). Viewed online at: <http://wwwcimis.water.ca.gov/cimis/welcome.jsp>. Accessed on: Aug
23 1, 2012. Last update: 2009.
- 24 California Department of Water Resources, 2012a, Guidelines, Integrated Regional Water Management,
25 Proposition 84 and 1E. Viewed online at:
26 http://www.water.ca.gov/irwm/grants/docs/Guidelines/GL_2012_FINAL.pdf. Accessed on: Feb
27 5, 2013. Last update: Nov 2012.
- 28 California Department of Water Resources 2012b, California Statewide Groundwater Elevation
29 Monitoring (CASGEM). Viewed online at: <http://www.water.ca.gov/groundwater/casgem/>.
30 Accessed: Sept 28, 2012. Last update: 2012.
- 31 California Department of Water Resources 2009, Comprehensive Water Package, Special Session Policy
32 Bills and Bond Summary. Viewed online at:
33 <http://www.water.ca.gov/legislation/docs/01272010waterpackage.pdf>. Accessed: Oct 1, 2012.
34 Last update: Nov 2009.
- 35 California Department of Water Resources, Integrated Regional Water Management Grants,
36 Disadvantaged Communities Mapping Tool. Viewed online at:

- 1 <http://www.water.ca.gov/irwm/grants/resourceslinks.cfm>. Accessed on: Feb 5, 2013. Last update:
2 Nov 29, 2012.
- 3 California Department of Water Resources, Statewide Agreements, Accomplishments. Viewed online at:
4 http://www.water.ca.gov/news/newsreleases/2010/122210statewide_agreements.pdf. Accessed:
5 Sept 10, 2012. Last Update: 2010.
- 6 California Emergency Management Agency and California Natural Resources Agency (Cal-EMA,
7 CNRA). 2012. California Adaptation Planning Guide.
8 (http://resources.ca.gov/climate_adaptation/local_government/adaptation_planning_guide.html).
- 9 California Emergency Management Agency and California Natural Resources Agency, California
10 Climate Change Adaptation Policy Guide. Viewed online at:
11 http://resources.ca.gov/climate_adaptation/docs/APG_-_PUBLIC_DRAFT_4.9.12_small.pdf.
12 Accessed: Feb 6, 2013. Last update: Apr 9, 2012.
- 13 California Energy Commission (CEC). 2006. Refining Estimates of Water-Related Energy Use in
14 California. Public Interest Energy Research Program (PIER) Final Project Report, CEC-500-
15 2006-118. Prepared by Navigant Consulting, Inc.
- 16 California Environmental Protection Agency, Surface Water Ambient Monitoring Program (SWAMP).
17 State Water Resources Control Board, 1001 I Street, Sacramento, CA 95814. Viewed online at:
18 http://www.waterboards.ca.gov/water_issues/programs/swamp/. Accessed on: Feb 6, 2013. Last
19 update: Jan 17, 2013.
- 20 California's Flood Future: Recommendations for Managing the State's Flood Risk, Public Draft, April
21 2013, California Department of Water Resources and U.S. Army Corps of Engineers. Viewed
22 online at: <http://www.water.ca.gov/sfmp/flood-future-report.cfm>. Accessed on June 21, 2013.
23 Last update: May 23, 2013.
- 24 California Indians K-344, 1942, Accounting and value of land taken without compensation under 18
25 unratified treaties, U.S. Supreme Court. Viewed online at: [http://www.tachi-yokut-
26 nsn.gov/Portals/0/Documents/ca_tribes1.pdf](http://www.tachi-yokut-nsn.gov/Portals/0/Documents/ca_tribes1.pdf). Accessed on: Mar 14, 2013. Last update: Dec 4,
27 1944.
- 28 California Land Settlement Act of 1851, Christine A. Klein, The University of Dayton School of Law.
29 Viewed online at: <http://academic.udayton.edu/race/02rights/guadalu6.htm#California>. Accessed
30 on: Mar 13, 2013. Last update: Apr 4, 2012.
- 31 California Legislature, Assembly Bill 2409, Urban Water Management, 2009-2010 Regular Session.
32 Viewed online at: [http://www.leginfo.ca.gov/pub/09-10/bill/asm/ab_2401-
33 2450/ab_2409_bill_20100406_amended_asm_v98.pdf](http://www.leginfo.ca.gov/pub/09-10/bill/asm/ab_2401-2450/ab_2409_bill_20100406_amended_asm_v98.pdf). Accessed: Sept 26, 2012. Last update:
34 Feb 2010.

- 1 California Native American Heritage Commission, Strategic Plan. California Native American Heritage
2 Commission, 915 Capitol Mall, Room 364 Sacramento, CA 95814 (916) 653-4082. Viewed
3 online at: <http://www.nahc.ca.gov/sp.html>. Accessed: Aug 3, 2012. Last update: [Unknown].
- 4 California Natural Resources Agency, 2009 California Climate Adaptation Strategy: A report to the
5 Governor of the State of California in Response to Executive Order S-13-2008. Viewed online at:
6 [http://www.energy.ca.gov/2009publications/CNRA-1000-2009-027/CNRA-1000-2009-027-](http://www.energy.ca.gov/2009publications/CNRA-1000-2009-027/CNRA-1000-2009-027-F.PDF)
7 [F.PDF](http://www.energy.ca.gov/2009publications/CNRA-1000-2009-027/CNRA-1000-2009-027-F.PDF). Accessed on: Feb 6, 2013. Last update: 2009.
- 8 California Natural Resources Agency (CNRA). 2009. 2009 California Climate Adaptation Strategy.
9 http://resources.ca.gov/climate_adaptation/docs/Statewide_Adaptation_Strategy.pdf.
- 10 California Powder Works Co v. Davis et al, 1894, U.S. Supreme Court. Viewed online at:
11 http://caselaw.lp.findlaw.com/scripts/printer_friendly.pl?page=us/151/389.html. Accessed on:
12 Mar 13, 2013. Last update: 2013.
- 13 California Public Utilities Commission, Renewables Portfolio Standard Program Overview, 2007.
14 Viewed online at: <http://www.cpuc.ca.gov/PUC/energy/Renewables/overview>. Accessed: Jun 26,
15 2013. Last update: Mar 1, 2013.
- 16 California Tribal Water Rights 2009, Briefing paper from the 2009 California Tribal Water Summit.
17 Viewed online at:
18 http://www.waterplan.water.ca.gov/docs/tws/CTWS_BriefingPaper_Rights_Parr_v1.pdf.
19 Accessed on: Jul 18, 2013. Last update: 2009.
- 20 California Urban Water Agencies, Public Investment White Paper, October 18, 2011. Viewed online at:
21 http://www.cuwa.org/pubs/PI_White_Paper_10_%20final.pdf. Accessed on: Sept 20, 2012. Last
22 update: Oct 2011.
- 23 Camp Dresser & McKee, 2008, Evaluation and Determination of Potential Liability Associated with the
24 Decommissioning and Removal of four Hydroelectric Dams on the Klamath River by Any Agent.
25 Prepared by: Camp Dresser & McKee Incorporated, Prepared for: U.S. Department of the
26 Interior, Through: U.S. Bureau of Reclamation. Viewed online at:
27 <http://www.usbr.gov/mp/kbao/kbra/docs/Doc2.pdf>. Accessed on: Mar 12, 2013. Last update: Jul
28 18, 2008. Cayan, D 2008. Climate change scenarios for the California region, Climatic Change,
29 87(s1), 21-S42. Viewed online at: http://tenaya.ucsd.edu/~dettinge/cccc08_scenarios.pdf.
30 Accessed on: Feb 6, 2013. Last update: Jan 2008.
- 31 City of Santa Rosa, City of Santa Rosa Geysers Project. Viewed online at: <http://ci.santa->
32 [rosa.ca.us/departments/utilities/irwp/geysers/Pages/default.aspx](http://ci.santa-rosa.ca.us/departments/utilities/irwp/geysers/Pages/default.aspx). Accessed: Sept 20, 2012. Last
33 update: 2010.
- 34 Common Waters Fund, Background on Forests and Water Quality. Common Waters Fund, P.O. Box
35 1029, Milford, PA 18337, 202-797-6530. Viewed online at:
36 <http://www.commonwatersfund.org/background-forests-and-water-quality>. Accessed: Aug 23,
37 2012. Last Update: [Unknown].

- 1 County of Humboldt, Community Development Services, Integrated Regional Water Management
2 Region Acceptance Process (RAP), Integrated Regional Water Management Program Applicant:
3 Humboldt County, Authorization and Eligibility Requirements. Contact Kirk Girard, Community
4 Development Services Director, County of Humboldt, Community Development Services, Clark
5 Complex, 3015 H Street, Eureka, CA 95501. Viewed online at:
6 http://www.northcoastirwmp.net/docManager/1000007982/Att1_IG1_NorthCoastIRWMP_Eligible_1of1.pdf. Accessed: Feb 6, 2013. Last update: 2011.
7
- 8 Cayan, D. 2008. Climate Change Scenarios for the California Region. *Climatic Change*, 87(s1), 21-S42.
- 9 de Jager, JM and van Zyl, WH, Atmospheric Evaporative Demand and Evaporation Coefficient Concepts,
10 Department of Agrometeorology, University of the Orange Free State, Bloemfontein, 9301, South
11 Africa, May 1988. Viewed online at:
12 <http://www.wrc.org.za/Lists/Knowledge%20Hub%20Items/Attachments/4182/524%20abstract.pdf>. Accessed on: Jul 25, 2012. Last update: [Unknown].
13
- 14 Dettinger, Michael 2011, Climate change, atmospheric rivers, and floods in California - A multi-model
15 analysis of storm frequency and magnitude changes, *Journal of the American Water Resources*
16 *Association*, 47(3): 514-523.
- 17 Doctrine of Indian Right to Occupancy and Possession of Land, 1927, *Indian Affairs: Laws and Treaties*.
18 Viewed online at: http://digital.library.okstate.edu/kappler/Vol4/html_files/v4p1166.html.
19 Accessed on: Mar 13, 2013. Last Update: 1929.
- 20 EcoTrust, Preliminary Economic Assessment of Dam Removal: The Klamath River, EcoTrust, 721 NW
21 Ninth Avenue, Suite 200, Portland, OR 97209. Viewed online at:
22 http://www.ecotrust.org/workingpapers/WPS2_Klamath_Dam_Assess.pdf. Accessed: Mar 12,
23 2013. Last update: Jan 31, 2006. Environmental Protection Agency Region 9, Resources Legacy
24 Fund, U.S. Army Corps & Department of Water Resources, 2011, *Climate Change Handbook for*
25 *Regional Water Planning*. Viewed online at:
26 <http://www.water.ca.gov/climatechange/CCHandbook.cfm>. Accessed on: Feb 6, 2013. Last
27 update: Nov 2011.
- 28 Eighteen Unratified Treaties, 1852, Tachi Yokut Tribe, Santa Rosa Rancheria, 16835 Alkali Drive,
29 Lemoore, California 93245. Viewed online at: http://www.tachi-yokut.com/treaty_intro.html.
30 Accessed on: Mar 14, 2013. Last update: 2011.
- 31 Fifty years past the deadline, 2001, Stapleton, Nolan, MacGregor & Thompson, St. Paul, Minnesota.
32 Viewed online at:
33 http://www.citizensalliance.org/links/pages/articles%20and%20CERA%20news/Indian%20Claims%20Commission%20Article.htm#_edn23. Accessed on: Mar 14, 2013. Last update: 2001.
34
- 35 Five Counties Salmonid Conservation Program, 2013, 5 Counties Salmonid Conservation Program, 40
36 Horseshoe Lane/ PO Box 2571, Weaverville, CA 96093-2571. Viewed online at:
37 <http://www.5counties.org/>. Accessed on: Mar 13, 2013. Last update: Mar 6, 2013.

- 1 Food and Agriculture Organization of the United Nations, Sustainable dryland cropping in relation to soil
2 productivity - FAO soils bulletin 72. Viewed online at:
3 <http://www.fao.org/docrep/V9926E/v9926e00.htm#Contents>. Accessed on: Jul 25, 2012. Last
4 update: 1995.
- 5 Gannett M, 2013, USGS Oregon Water Science Center, Upper Klamath Basin Ground-Water Study.
6 Viewed online at: http://or.water.usgs.gov/projs_dir/or180/background.html, Accessed: Feb 6,
7 2013. Last update: Jan 2013.
- 8 Georgakakos K, Graham N, Kistenmacher M, et al. Value of adaptive water resources management in
9 northern California under climatic variability and change: Dynamic hydroclimatology. Journal Of
10 Hydrology. January 4, 2012;412-413:47-65.
- 11 Georgia State University, Department of Physics and Astronomy, Relative Humidity. Georgia State
12 University, Atlanta, Georgia 30302-4106, August 2000. Viewed online at:
13 <http://hyperphysics.phy-astr.gsu.edu/hbase/kinetic/relhum.html#c5>. Accessed on: Jul 26, 2012.
14 Last update: [Unknown].
- 15 Hayt Charles D v. U.S. et al, [date unknown], Justice Howry ruling, (case not appealed to the Supreme
16 Court). Viewed online at: <http://users.sisqtel.net/armstrng/excerptsIndIndposs..htm>. Accessed on:
17 Mar 13, 2013. Last update: [Unknown].
- 18 Hoopa Valley Indian Tribe, California Tribal Water Summit 2009, Tribal Water Authorities-Rivers,
19 Dams & Fish. Hoopa Valley Tribe, P.O. Box 1348 Hoopa, CA 95546, Thomas P. Schlosser,
20 October 21, 2009, PDF. Viewed online at:
21 http://www.waterplan.water.ca.gov/docs/tws/CTWS_BriefingPaper_Rivers2_Hoopa_v1.pdf.
22 Accessed on: Feb 6, 2013. Last update: Oct 2009.
- 23 Hoopa Valley Indian Tribe, Tribal History. Hoopa Valley Indian Tribe, PO Box 1348 Hoopa, California
24 95546 (530) 625.4211. Viewed online at: <http://www.hoopa-nsn.gov/culture/history.htm>.
25 Accessed on: Sept 6, 2012. Last update: 2003.
- 26 Hoopa-Yurok Settlement Act, 1988a. Viewed online at:
27 <http://www.yuroktribe.org/government/tribalattorney/documents/HoopaYurokSettlementAct.pdf>.
28 Accessed on: Jul 8, 2013. Last update: Oct 31, 1988.
- 29 Hoopa-Yurok Settlement Act, 1988b. Viewed online at:
30 <http://www.govtrack.us/congress/bills/100/s2723>. Accessed on Jul 8, 2013. Last update: Oct 31,
31 1988.
- 32 Humboldt Bay Municipal Water District, Water Supply. Humboldt Bay Municipal Water District, 828
33 Seventh Street, Eureka, California, (707)443-5018. Viewed online at:
34 http://www.hbmwd.com/water_supply. Accessed on: Sept 12, 2012. Last update: 2012.
- 35 Humboldt County Public Works, Natural Resources & Water Management. Humboldt County Public
36 Works, Environmental Services Division, 1106 2nd Street in Eureka, CA. (707) 445-7741.

- 1 Viewed online at: <http://co.humboldt.ca.us/pubworks/resources-water.asp>. Accessed on: Aug 7,
2 2012. Last update: 2005.
- 3 Humboldt County Resource Conservation District, Salt River Ecosystem Restoration Project. Humboldt
4 County Resource Conservation District, 5630 South Broadway, Eureka, CA 95503. Viewed online
5 at: http://humboldtred.org/index_files/salt_river_ecosystem_restoration_project.htm. Accessed:
6 Aug 7, 2012. Last update: [Unknown].
- 7 Humboldt State University, Department of Geology, Redwood Coast Tsunami Work Group c2010a,
8 About the RCTWG. Humboldt State University, Department of Geology, 1 Harpst Street, Arcata,
9 California, 95521. Viewed online at: <http://humboldt.edu/rctwg/site/about/>. Accessed on: Aug 22,
10 2012. Last update: [Unknown].
- 11 Humboldt State University, Department of Geology, Redwood Coast Tsunami Work Group c2010b,
12 Projects. Viewed online at: <http://humboldt.edu/rctwg/site/projects/>. Accessed on: Aug 22, 2012.
13 Last update: [Unknown].
- 14 Hynes v. Grimes Packing Company, 1949, U.S. Supreme Court. Viewed online at:
15 <http://caselaw.lp.findlaw.com/cgi-bin/getcase.pl?court=US&vol=337&invol=86&friend=nytimes>.
16 Accessed on: Mar 14, 2013. Last update: May 31, 1949.
- 17 Indian Claims Commission Act of 1946, U.S. Department of Justice. Viewed online at:
18 <http://www.justice.gov/enrd/3147.htm>. Accessed on: Mar 14, 2013. Last update: Nov 2010.
- 19 Intergovernmental Panel on Climate Change, Climate Change 2007 Synthesis Report: Contribution of
20 Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on
21 Climate Change, [Pachauri, R.K. and Reisinger, A. (eds.)]. Viewed online at:
22 [http://www.ipcc.ch/publications_and_data/publications_ipcc_fourth_assessment_report_synthesis](http://www.ipcc.ch/publications_and_data/publications_ipcc_fourth_assessment_report_synthesis_report.htm)
23 [_report.htm](http://www.ipcc.ch/publications_and_data/publications_ipcc_fourth_assessment_report_synthesis_report.htm). Accessed on: Feb 6, 2013. Last update: Jun 2012.
- 24 Intergovernmental Panel on Climate Change (IPCC). 2007. IPCC Fourth Assessment Report: Climate
25 Change 2007: Synthesis Report. http://www.ipcc.ch/publications_and_data/ar4/syr/en/main.html.
- 26 Irrigated Lands Discharge Program, 2006. Prepared by: Jones & Stokes Consultants. Prepared for:
27 Central Valley Water Board. Irrigated Lands Program, Existing Conditions Report, Feb 2006.
- 28 Janetos A, Hansen L, Inouye D, Kelly BP, Meyerson L, Peterson B, and Shaw R, "Biodiversity", The
29 effects of climate change on agriculture, land resources, water resources, and biodiversity in the
30 United States. A report by the U.S. Climate Change Science Program and the subcommittee on
31 Global Change Research, Washington, DC. 362 pp. 2008.
- 32 Johnstone JA, Dawson TE, 2010. Climatic context and ecological implications of summer fog decline in
33 the coast redwood region, Proceedings of the National Academy of Science 107: 4,533-4,538.
- 34 Jones HG, Plants and Microclimate: A Quantitative Approach to Environmental Plant Physiology, Jun 26,
35 1992, second edition; ISBN-10: 0521425247 | ISBN-13: 978-0521425247.

- 1 Jurisdictional Act of May 18, 1928, amended Apr 29, 1930 and Jun 30, 1948, enacted by the Senate and
2 House of Representatives of the United States. Viewed online at:
3 http://digital.library.okstate.edu/kappler/vol6/html_files/v6p0502b.html. Accessed on: Mar 14,
4 2013. Last update: May 24, 1950.
- 5 Klamath Basin Coordinating Council, 2011 Annual Report, Implementing the Klamath Basin Settlement
6 Agreements. Viewed online at: <http://www.klamathcouncil.org/>. Accessed on: Sept 28, 2012.
7 Last update: May 2011.
- 8 Klamath Basin Crisis, 2009, Klamath River Basin Compact, Klamath Basin Crisis, PO Box 314,
9 Tulelake, CA 96134. Viewed online at:
10 <http://www.klamathbasincrisis.org/history/klamathcompact101705.htm>. Accessed: Mar 12, 2013.
11 Last update: May 7, 2009.
- 12 Klamath Basin Settlement Agreement documentation, 2013, Klamath Basin Settlement Agreement,
13 County of Siskiyou. Viewed online at: <http://www.co.siskiyou.ca.us/BOS/kbra.aspx>. Accessed
14 on: Mar 12, 2013. Last update: Jul 24, 2012.
- 15 Klamath Basin Water Crisis, 2001, Klamath Basin Crisis, PO Box 314, Tulelake, CA 96134. Viewed
16 online at: <http://www.klamathbasincrisis.org/Photos/photos-list.htm>. Accessed on: Mar 15, 2013.
17 Last update: May 7, 2009.
- 18 Klamath Facilities Removal Environmental Impact Statement/Environmental Impact Report - Comments
19 of the County of Siskiyou, City of Yreka, City of Dorris, City of Etna, City of Montague, City of
20 Weed, and Town of Fort Jones, November 21, 2011. Viewed online at:
21 [http://www.co.siskiyou.ca.us/BOS/DOCS/KBRA/Comments_of_SiskiyouCountyandCities_to_K](http://www.co.siskiyou.ca.us/BOS/DOCS/KBRA/Comments_of_SiskiyouCountyandCities_to_KlamathFacilitiesRemoval_EIR_EIS%2011_21_11.pdf)
22 [lamathFacilitiesRemoval_EIR_EIS%2011_21_11.pdf](http://www.co.siskiyou.ca.us/BOS/DOCS/KBRA/Comments_of_SiskiyouCountyandCities_to_KlamathFacilitiesRemoval_EIR_EIS%2011_21_11.pdf). Accessed on: Mar 12, 2013. Last update:
23 Nov 21, 2011. Klamath Restoration Agreements, Klamath Basin Restoration Agreement. Viewed
24 online at: <http://klamathriverrestoration.org/kbra-summary.html>. Accessed: Aug 24, 2012. Last
25 update: 2010.
- 26 Klamath River Basin Compact, 1957a, California Water Code 5900-5901. Viewed online at:
27 [http://www.leginfo.ca.gov/cgi-bin/displaycode?section=wat&group=05001-06000&file=5900-](http://www.leginfo.ca.gov/cgi-bin/displaycode?section=wat&group=05001-06000&file=5900-5901)
28 [5901](http://www.leginfo.ca.gov/cgi-bin/displaycode?section=wat&group=05001-06000&file=5900-5901). Accessed on: Mar 15, 2013. Last update: 1957.
- 29 Klamath River Basin Compact, 1957b, Articles, Oregon State Legislature, Oregon Revised Statutes, 2011
30 Edition, Chapter 542.610 thru 542.750. Viewed online at:
31 <http://landru.leg.state.or.us/ors/542.html>. Accessed on: Mar 15, 2013. Last update: 2011.
- 32 Knowles N, Dettinger M, Cayan D, 2007. Trends in Snowfall versus Rainfall for the Western United
33 States, 1949-2001. Prepared for California Energy Commission Public Interest Energy Research
34 Program, Project Report CEC-500-2007-032.
- 35 Laguna de Santa Rosa Foundation, Laguna de Santa Rosa Watershed Management Plan. Laguna de Santa
36 Rosa Foundation, 900 Sanford Road Santa Rosa, CA 95401, 707.527.9277. Viewed online at:

- 1 http://www.lagunadesantarosa.com/programs_rp_mp.shtml. Accessed: Aug 30, 2012. Last
2 update: 2011.
- 3 Lenihan JM, Bachelet D, Drapek R, Neilson RP, 2006, The response of vegetation distribution,
4 ecosystem productivity, and fire in California to future climate scenarios simulated by the mc1
5 dynamic vegetation model. California Climate Change Center, California Energy Commission,
6 Publication #CEC-500-2005-191-SF. Viewed online at:
7 <http://www.energy.ca.gov/2005publications/CEC-500-2005-191/CEC-500-2005-191-SF.PDF>.
8 Accessed: Feb 6, 2013. Last update: Feb 2006.
- 9 Mantua NJ, Hare SR, Zhang Y, Wallace JM, Francis RC, A Pacific Interdecadal Climate Oscillation with
10 Impacts on Salmon Production, 1997, Bulletin of the American Meteorological Society, Volume
11 78, Number 6, June 1997.
- 12 Markstrom SL, Hay LE, Ward-Garrison CD, Risley JC, Battaglin WA, Bjerklie DM, Chase KJ,
13 Christiansen DE, Dudley RW, Hunt RJ, et al., 2012, Integrated watershed-scale response to
14 climate change for selected basins across the United States: U.S. Geological Survey Scientific
15 Investigations Report 2011-5077, 143 pp. Viewed online at:
16 http://pubs.usgs.gov/sir/2011/5077/SIR11-5077_508.pdf. Accessed: Feb 6, 2013. Last update:
17 2012.
- 18 Mathany TM, Dawson BJ, Shelton JL, Belitz K, 2011, Ground water quality data in the northern Coast
19 Ranges study unit, 2009, Results from the California GAMA Program, U.S. Geological Survey
20 Data-Series 609, 92 p. Viewed online at: <http://pubs.usgs.gov/ds/609/pdf/ds609.pdf>. Accessed on:
21 Jul 17, 2012. Last update: Oct 19, 2011.
- 22 Mendocino Land Trust, Strategic Plan for the Big River Program of the Mendocino Land Trust, Adopted
23 by the Mendocino Land Trust Board of Trustees January 21, 2005. Viewed online at:
24 http://www.mendocinolandtrust.org/?download=Big_River_Strategic_Plan_web_final.pdf.
25 Accessed: Aug 23, 2012. Last update: 2012.
- 26 Mid Klamath Watershed Council, Overview of the Mid Klamath Sub-basin, 2006. Mid Klamath
27 Watershed Council, P.O. Box 409, Orleans, California 95556, 530.627.3020. Viewed online at:
28 <http://mkwc.org/midklamath/index.html>. Accessed: Feb 7, 2013. Last update: 2006.
- 29 Milly PCD, Betancourt J, Falkenmark M, Hirsch RM, Kundzewicz ZW, Lettenmaier DP, Stouffer RJ,
30 Stationarity is Dead: Whither Water Management?, Science 319:573-574. Viewed online at:
31 http://www.paztcn.wr.usgs.gov/julio_pdf/milly_et_al.pdf. Accessed: Feb 7, 2013. Last update:
32 Feb 2008.
- 33 National Agricultural Statistics Service, U.S. Department of Agriculture, California Grape Acreage, 2007
34 Crop. Viewed online at:
35 http://www.nass.usda.gov/Statistics_by_State/California/Publications/Grape_Acreage/200704gab
36 [tb00.pdf](http://www.nass.usda.gov/Statistics_by_State/California/Publications/Grape_Acreage/200704gabt00.pdf). Accessed: Aug 13, 2012. Last update: Apr 2008.

- 1 National Agricultural Statistics Service, USDA, Fruits and Nuts: 1992 and 1987; Viewed online at,
2 http://www.agcensus.usda.gov/Publications/1992/Volume_1_Chapter_2_County_Tables/California/ca2_31.pdf;
3 Accessed on Aug 13, 2012; Last update, Jul 19, 1994.
- 4 National Agricultural Statistics Service, U.S. Department of Agriculture, County Summary Highlights:
5 2002. Viewed online at:
6 http://www.agcensus.usda.gov/Publications/2002/Volume_1_Chapter_2_County_Level/California/st06_2_001_001.pdf. Accessed on: Aug 10, 2012. Last update: Mar 29, 2012.
7
- 8 National Agricultural Statistics Service, U.S. Department of Agriculture, County Summary Highlights:
9 2007. Viewed online at:
10 http://www.agcensus.usda.gov/Publications/2007/Full_Report/Volume_1_Chapter_2_County_Level/California/st06_2_001_001.pdf. Accessed on: Aug 10, 2012. Last update: Mar 29, 2012.
11
- 12 National Forest Growth, 2009, Siskiyou County Board of Supervisors. Viewed online at:
13 http://users.sisqtel.net/armstrng/national_forest_growth.htm. Accessed on: Mar 13, 2013. Last
14 update: [Unknown].
- 15 National Marine Fisheries Service, Southwest Region, Biological Opinion, Operation of the Klamath
16 Project between 2010 and 2018, Consultation provided for U.S. Bureau of Reclamation. Viewed
17 online at: http://swr.nmfs.noaa.gov/klamath/FINAL-Klamath_Ops_031510.pdf. Accessed: Sept
18 12, 2012. Last update: Mar 2010.
- 19 National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Southwest
20 Regional Office, Draft Southern Oregon/ Northern California Coast Coho Salmon Recovery Plan,
21 Bear River Population, Jan 2012. Viewed online at:
22 http://swr.nmfs.noaa.gov/recovery/soncc_draft/Chapter_28_Bear_River_Population.pdf.
23 Accessed: Sept 11, 2012. Last update: Jan 2012.
- 24 National Oceanic and Atmospheric Administration, National Weather Service, Northwest California
25 Rainfall, Eureka Weather Forecast Office, 300 Startare Drive, Eureka, California, 95501,
26 707.443.6484. Viewed online at: <http://www.wrh.noaa.gov/eka/rivers/>. Accessed: Jun 26, 2012.
27 Last update: [Unknown].
- 28 National Oceanic and Atmospheric Administration, Tides & Currents, Sea Levels Online. Sea Level
29 Variations of the United States Derived from National Water Level Observation Network
30 Stations. Viewed online at: <http://tidesandcurrents.noaa.gov/sltrends/sltrends.shtml>. Accessed:
31 Feb 7, 2013. Last update: Oct 2012.
- 32 National Research Council of the National Academies (NRC). 2012. Sea-Level Rise for the Coasts of
33 California, Oregon, and Washington: Past, Present, and Future.
- 34 North Coast Integrated Regional Water Management Plan, Phase I, Regional Framework for Integrated
35 Planning in the North Coast, 3-6. Viewed online at:
36 http://www.northcoastirwmp.net/docManager/1000006299/NCIRWMP_Phase_I_maps_2007.pdf
37 . Accessed on: Aug 31, 2012. Last update: Jul 2007.

- 1 North Coast Integrated Regional Water Management Plan, Klamath Watershed Management Area.
2 Viewed online at: <http://www.northcoastirwmp.net/Content/10301/preview.html#map>. Accessed:
3 Jul 18, 2012. Last update: 2007.
- 4 North Coast Integrated Regional Water Management Plan, Mattole Integrated Water Management
5 Program. Viewed online at: <http://www.northcoastirwmp.net/Content/10375/preview.html>.
6 Accessed: Aug 7, 2012. Last update: 2007.
- 7 North Coast Integrated Regional Water Management Plan, Salt River Ecosystem Restoration. Viewed
8 online at: <http://www.northcoastirwmp.net/Content/10394/preview.html>. Accessed: Aug 7, 2012.
9 Last update: 2007.
- 10 North Coast Integrated Regional Water Management Plan, Phase I, 2007. Viewed online at:
11 <http://www.northcoastirwmp.net/Content/10319/preview.html>. Accessed: Jul 24, 2012. Last
12 update: 2007.
- 13 North Coast Regional Water Quality Control Board, 1993, Water Quality Control Plan for the North
14 Coast Region. Viewed online at:
15 http://www.krisweb.com/biblio/ncc_swrcb_ncrwqcb_2001_wqcontrolplan.pdf. Accessed: Nov
16 2012. Last update: June 2001.
- 17 North Coast Regional Water Quality Control Board, 2007, Water Quality Control Plan For The North
18 Coast Region, North Coast Regional Water Quality Control Board, 5550 Skylane Blvd., Suite A,
19 Santa Rosa, CA 95403, Telephone: (707) 576-2220. Viewed online at:
20 http://www.northcoastirwmp.net/docManager/1000006306/ncrwqcb_basin_plan.pdf. Accessed:
21 Jul 20, 2012. Last update: Jan 2007.
- 22 North Coast Regional Water Quality Control Board, Salmon River, Siskiyou County, California, Total
23 Maximum Daily Load for Temperature and Implementation Plan. Viewed online at:
24 http://www.swrcb.ca.gov/northcoast/water_issues/programs/tmdls/salmon_river/062405/part_1_s
25 [almon_temperature_tmdl_report_adopted.pdf](http://www.swrcb.ca.gov/northcoast/water_issues/programs/tmdls/salmon_river/062405/part_1_salmon_temperature_tmdl_report_adopted.pdf). Accessed: Jun 29, 2012. Last update: June 2005.
- 26 North Coast Regional Water Quality Control Board, 2011, Water Quality Control Plan for the North
27 Coast Region. Viewed online at:
28 http://www.swrcb.ca.gov/northcoast/water_issues/programs/basin_plan/083105-
29 [bp/basin_plan.pdf](http://www.swrcb.ca.gov/northcoast/water_issues/programs/basin_plan/083105-bp/basin_plan.pdf). Accessed on: Sept 7, 2012. Last update: May 2011.
- 30 North Marin Water District 2013a, Novato History. Viewed online at: [_](http://www.nmwd.com/about_history_novato.php)
31 http://www.nmwd.com/about_history_novato.php. Accessed: Jan 24, 2013. Last update: 2013.
- 32 North Marin Water District 2013b, Sources of Water. Viewed online at:
33 <http://www.nmwd.com/services.php>. Accessed on: Jan 24, 2013. Last update: 2013.
- 34 Northwestern Bands of Shoshone Indians v. United States, 1945, U.S. Supreme Court. Viewed online at:
35 <http://caselaw.lp.findlaw.com/cgi-bin/getcase.pl?navby=case&court=us&vol=324&invol=335>.
36 Accessed on: Mar 14, 2013. Last update: Mar 12, 1945.

- 1 Oregon backs Klamath Tribes water rights, March 10, 2013. Viewed online at: [http://www.times-](http://www.times-standard.com/news/ci_22759422/oregon-backs-klamath-tribes-water-rights-effect-lower)
2 [standard.com/news/ci_22759422/oregon-backs-klamath-tribes-water-rights-effect-lower.](http://www.times-standard.com/news/ci_22759422/oregon-backs-klamath-tribes-water-rights-effect-lower)
3 Accessed on: Jul 17, 2013. Last update: Mar 2013.
- 4 Oregon State University, Parameter-elevation Regressions on Independent Slopes Model (PRISM)
5 Climate Group, PRISM Data Explorer. Wayne Gibson, Northwest Alliance for Computational
6 Science & Engineering, 2000 Kelley Engineering Center, OSU, Corvallis, Oregon, 97331,
7 541.737.2531. Viewed online at: <http://prismmap.nacse.org/nn/index.phtml>. Accessed on: Jul 31,
8 2012. Last update: 2012.
- 9 Palomar College, Behavioral Sciences Department, Basic Principles of Genetics, San Marcos, California,
10 Dr. Dennis O'Neil, 1998 – 2012. Viewed online at:
11 <http://anthro.palomar.edu/mendel/Default.htm>. Accessed: Jul 25, 2012. Last update: 2012.
- 12 Petaluma 360.com, Press Democrat opinion article, City's water supply is subject of study (Petaluma).
13 Viewed online at:
14 [http://www.pressdemocrat.com/article/20120107/COMMUNITY/120109799/1042/opinion.](http://www.pressdemocrat.com/article/20120107/COMMUNITY/120109799/1042/opinion)
15 Accessed on: Jan 24, 2013. Last update: Jan 2012.
- 16 Pierce, D.W., T. Das, D.R. Cayan, E.P. Maurer, N.L. Miller, Y. Bao, M. Kanamitsu, K. Yoshimura, M.A.
17 Snyder, L.C. Sloan, G. Franco, and M. Tyree. Probabilistic estimates of future changes in
18 California temperature and precipitation using statistical and dynamical downscaling. Climate
19 Dynamics, 2012. Springer-Verlag, DOI 10.1007/s00382-012-1337-9.
20 http://meteora.ucsd.edu/cap/pdf/Pierce_etal_2012_CD.pdf.
- 21 Potter Valley Irrigation District, Flows in the Eel River and the Potter Valley Project, Potter Valley
22 Irrigation District, P.O. Box 186, 10170 Main Street, Potter Valley, CA 95469, (707) 743-1109.
23 Viewed online at: <http://www.pottervalleywater.org/flows.html>. Accessed: Sept 11, 2012. Last
24 update: [Unknown].
- 25 Potter Valley Irrigation District, History of the Potter Valley Project, Potter Valley Irrigation District,
26 P.O. Box 186, 10170 Main Street, Potter Valley, California 95469, 707.743.1109. Viewed online
27 at: <http://www.pottervalleywater.org/history.html>. Accessed on: Jul 23, 2012. Last update:
28 [Unknown].
- 29 Qian, Yun, Steven J. Ghan, and L. Ruby Leung. 2010. Downscaling Hydroclimatic Changes Over the
30 Western US Based on CAM Subgrid Scheme and WRF Regional Climate Simulations.
31 International Journal of Climatology, April.
- 32 Redwood Creek Watershed Group, Redwood Creek Integrated Watershed Strategy, Jun 2006. Viewed
33 online at: <http://www.nps.gov/redw/parkmgmt/upload/RWC%20IWS%20Final.pdf>. Accessed on:
34 Feb 8, 2013. Last update: Jun 2006.
- 35 Request for opinion on "Rancheria Act" of August 18, 1958, 1960, Requested by Solicitor to
36 Commissioner of Indian Affairs, pp 1882-1886. Viewed online at:

- 1 http://thorpe.ou.edu/sol_opinions/p1876-1900.htm. Accessed on: Mar 14, 2013. Last update: Aug
2 1, 1960.
- 3 Rosenberg N J, Blad BL, Verma SB, Microclimate: The Biological Environment, Sept 1983, second
4 edition, ISBN: 978-0-471-06066-6. Wiley Press.
- 5 Round Valley Indian Tribes, A Sovereign Nation of Confederated Tribes, Round Valley Indian Tribes,
6 77826 Covelo Road, Covelo, California 95428, (707) 983-6126. Viewed online at:
7 <http://www.rvit.org/index.html>. Accessed: Sept 6, 2012. Last update: 2011.
- 8 Round Valley Indian Tribes, Population Projections 2010 – 2025. Prepared by: The Center For Applied
9 Research, 1133 Filbert Street, San Francisco, CA 94109, (415) 310-1108. Viewed online at:
10 <http://www.rvitplanning.com/RVIT%20Population%20Projections.pdf>. Accessed on: Sept 6,
11 2012. Last update: Jun 2010.
- 12 Russell N, Griggs G. 2012. Adapting to Sea Level Rise: A Guide for California’s Coastal Communities.
13 University of California, Santa Cruz for the California Energy Commission Public Interest
14 Environmental Research Program 2012
- 15 Salmon Creek Watershed Council, Salmon Creek Estuary Habitat Structures Project (Sonoma County).
16 Viewed online at: <http://www.salmoncreekwater.org/SalmonCreekEstuaryProject.pdf>. Accessed
17 on: Aug 23, 2012. Last update: 2010.
- 18 Salmon River Restoration Council, A Quick Look at the Salmon River Watershed (Siskiyou County),
19 P.O. Box 1089, Sawyers Bar, California 96027, 530.462.4665. Viewed online at:
20 <http://www.srrc.org/watershed/index.php>. Accessed on: Jun 28, 2012. Last update: [Unknown].
- 21 Scott River Water Trust, 2013, Scott River Water Trust, P.O. Box 591, Etna, CA 96027. Viewed online at:
22 <http://www.scottwatertrust.org/>. Accessed on: Mar 13, 2013. Last update: 2013.
- 23 Scott Valley Ground Water Management Plan, 2012, Siskiyou County Board of Supervisors. Viewed
24 online at:
25 [http://www.co.siskiyou.ca.us/BOS/DOCS/agenda/2013/Questys/MG23893/AS23903/AI24176/D
26 O24177/DO_24177.pdf](http://www.co.siskiyou.ca.us/BOS/DOCS/agenda/2013/Questys/MG23893/AS23903/AI24176/D024177/DO_24177.pdf). Accessed on: Mar 12, 2013. Last update: Oct 22, 2012.
- 27 Shasta River Coordinated Resources Management and Planning, Shasta River CRMP, P.O. Box 459,
28 Montague, California 96064, 530.842.6121. Viewed online at:
29 <http://www.snowcrest.net/shastacrmf/index.html>. Accessed on: Jun 27, 2012. Last update: 2000.
- 30 Shasta River Water Association, 2013, Shasta Valley Resource Conservation District, 215 Executive
31 Court, Suite A, Yreka, CA 96097. Viewed online at: [http://svrcd.org/wordpress/projects/shasta-
32 water-association/](http://svrcd.org/wordpress/projects/shasta-water-association/). Accessed on: Mar 13, 2013. Last update: [Unknown].
- 33 Shasta Valley Resource Conservation District, Shasta River Water Trust, 215 Executive Court, Suite A,
34 Yreka, CA 96097, (530) 842-6121, ext. 106. Viewed online at:

- 1 <http://svrcd.org/wordpress/projects/shasta-river-water-trust/>. Accessed: Aug 23, 2012. Last
2 update: Aug 2012.
- 3 Siskiyou County Code of Ordinances, 2012, Title 3, Chapters 13 and 19. Viewed online at:
4 http://www.municode.com/library/CA/Siskiyou_County. Accessed on: Mar 15, 2013. Last
5 update: 2012.
- 6 Siskiyou County Flood Control and Water Conservation District, 1959, 2010 California Code, Public
7 Contract Code. Viewed online at: [http://law.justia.com/codes/california/2010/pcc/21400-
8 21401.html](http://law.justia.com/codes/california/2010/pcc/21400-21401.html). Accessed on: Mar 15, 2013. Last update: 2010.
- 9 Siskiyou County Flood Control and Water Conservation District Act, 1957, Appendix 89 to the California
10 Water Code. Viewed online at: <http://www.klamathbasin-crisis.org/compact/AppendixB.pdf>.
11 Accessed on: Mar 15, 2013. Last update: (Unknown).
- 12 Siskiyou County Ordinances, 2009, Municode Library, Siskiyou County, CA, Code of Ordinances, Title
13 10, Chapter 13, Demolition, Deconstruction, Removal and Reclamation. Viewed online at:
14 <http://library.municode.com/index.aspx?clientId=16630>. Accessed on: Mar 12, 2013. Last update:
15 Apr 7, 2009.
- 16 Siskiyou Resource Conservation District, Scott River Watershed Council, Oct 2005 Update, Initial Phase
17 of the Scott River Watershed Council Strategic Action Plan, Siskiyou Resource Conservation
18 District, P.O. Box 268, Etna, California 96027.
- 19 Smith River Advisory Council, Smith River Anadromous Fish Action Plan, version 1.0. Viewed online
20 at: <http://www.smithriveralliance.org/resources/library/SRAFAP.pdf>. Accessed on: Sept 7, 2012.
21 Last update: Mar 2002.
- 22 Somach Simmons & Dunn, Environmental Law and Policy Alerts, January 8, 2013, Siskiyou County
23 Farm Bureau v. California Department of Fish & Game, Siskiyou County Superior Court Case
24 No. SC SC CV 11-00418. Viewed online at: <http://www.somachlaw.com/alerts.php?id=201>.
25 Accessed on May 31, 2013. Last update: Jan 8, 2013.
- 26 Sonoma County Agricultural Commissioner, 2007 Sonoma County Agricultural Crop Report, Sonoma
27 County Agricultural Commissioner, 133 Aviation Blvd, Suite 110, Santa Rosa, CA 95403, (707)
28 565-2371. Viewed online at: http://www.sonoma-county.org/agcomm/pdf/2007_crop_report.pdf.
29 Accessed on: Aug 13, 2012. Last update: Jun 2008.
- 30 Sonoma County Water Agency, Russian River Action Plan (KRIS Edition) 2nd Edition, Sonoma County
31 Water Agency, 2003404 Aviation Blvd, Santa Rosa, California 95403, 707.526.5370. Viewed
32 online at: http://www.krisweb.com/biblio/russian_scwa_scwa_2002_actionplan.pdf. Accessed:
33 Feb 8, 2013. Last update: Mar 2003.
- 34 SPOA Settlement, 2013. Viewed online at:
35 [http://www.yuroktribe.org/resort/documents/1SPOA_REFERENDUM_PRESS_RELEASE_FIN
AL1.pdf](http://www.yuroktribe.org/resort/documents/1SPOA_REFERENDUM_PRESS_RELEASE_FIN
36 AL1.pdf). Accessed on Jul 9, 2013. Last update: Jan 2013.

- 1 State Water Resources Control Board, Division of Water Rights, Policy for Maintaining Instream Flows
2 in Northern California Coastal Streams, State Water Resources Control Board, P.O. Box 100,
3 Sacramento, CA 95812-0100, (916) 341-5250. Viewed online at:
4 [http://www.waterboards.ca.gov/waterrights/water_issues/programs/instream_flows/docs/ab2121_](http://www.waterboards.ca.gov/waterrights/water_issues/programs/instream_flows/docs/ab2121_0210/adopted050410instreamflowpolicy.pdf)
5 [0210/adopted050410instreamflowpolicy.pdf](http://www.waterboards.ca.gov/waterrights/water_issues/programs/instream_flows/docs/ab2121_0210/adopted050410instreamflowpolicy.pdf). Accessed: Aug 23, 2012; Last Update: Oct 2010.
- 6 State Water Resources Control Board, National Pollutant Discharge Elimination System. Viewed online
7 at: http://www.waterboards.ca.gov/water_issues/programs/npdes/. Accessed on: Jan 31, 2013.
8 Last update: Nov 28, 2012.
- 9 State Water Resources Control Board, Storm Water Program, Glossary (MS4). Viewed online at:
10 [http://www.waterboards.ca.gov/water_issues/programs/stormwater/docs/phsii2012_4th/att_i_glos](http://www.waterboards.ca.gov/water_issues/programs/stormwater/docs/phsii2012_4th/att_i_glossary_final.pdf)
11 [sary_final.pdf](http://www.waterboards.ca.gov/water_issues/programs/stormwater/docs/phsii2012_4th/att_i_glossary_final.pdf). Accessed on: Jan 31, 2013. Last update on: Jan 24, 2013.
- 12 Stories of Klamath Basin Settlers, 2012, Klamath Basin Crisis, PO Box 314, Tulelake, CA 96134.
13 Viewed online at: <http://www.klamathbasincrisis.org/stories/storiestoc.htm>. Accessed on: Mar 15,
14 2013. Last update: Dec 27, 2012.
- 15 Summa Corporation v. California Ex Rel. Lands Commission, 1984, U.S. Supreme Court. Viewed online
16 at: <http://caselaw.lp.findlaw.com/cgi-bin/getcase.pl?court=US&vol=466&invol=198>. Accessed
17 on: Mar 13, 2013. Last update: Apr 17, 1984.
- 18 SUPER et al. v. WORK, Secretary of the Interior, et al, 1926, Marcia Armstrong, Siskiyou County
19 Supervisor, District 5. Viewed online at: <http://users.sisqtel.net/armstrng/SupervWork.htm>.
20 Accessed on: Mar 14, 2013. Last update: Apr 16, 1926.
- 21 Tee-Hit-Ton Indians v. United States, 1955, U.S. Supreme Court. Viewed online at:
22 <http://caselaw.lp.findlaw.com/scripts/getcase.pl?court=us&vol=348&invol=272>. Accessed on:
23 Mar 13, 2013. Last update: Feb 7, 1955.
- 24 Timber Harvest Levels on the Major National Forests in Siskiyou County, 1978-2009, Siskiyou County
25 Board of Supervisors. Viewed online at:
26 <http://users.sisqtel.net/armstrng/timber%20harvest%20NF.htm>. Accessed on Mar 13, 2013. Last
27 update: [Unknown].
- 28 Treaty of Guadalupe Hidalgo, 1848, Library of Congress, Hispanic Division Area Studies. Viewed online
29 at: <http://www.loc.gov/rr/hispanic/ghtreaty>. Accessed on: Mar 13, 2013. Last update: Feb 14,
30 2011.
- 31 Tribes of the Lower Klamath River System - Question of Treaties and Land Title, [date unkown], Marcia
32 Armstrong, Siskiyou County Supervisor, District 5. Viewed online at:
33 <http://users.sisqtel.net/armstrng/Indtreaties.htm>. Accessed on: Mar 14, 2013. Last update:
34 (Unknown).
- 35 Trinity River Restoration Program, Trinity River Restoration Program 2011 Annual Report, Trinity River
36 Restoration Program, P.O. Box 1300, 1313 South Main Street, Weaverville, California 96093,

- 1 530.623.1800. Viewed online at: <http://odp.trrp.net/FileDatabase/Documents/TRRP->
2 [2011_annual_rept_final1.pdf](http://odp.trrp.net/FileDatabase/Documents/TRRP-2011_annual_rept_final1.pdf). Accessed: Sept 11, 2012. Last update: May 2012.
- 3 Trinity River Restoration Program 2011, Typical Releases from Lewiston Dam. Viewed online at:
4 http://www.trrp.net/?page_id=391. Accessed on: Jul 17, 2012. Last update: [Unknown].
- 5 U.S. Army Corps of Engineers, San Francisco District, Russian River Ecosystem Restoration, U.S. Army
6 Corps of Engineers, Deputy District Engineer for Civil Works, Arijs Rakstins, (415) 503-6720.
7 Viewed online at:
8 <http://www.spn.usace.army.mil/projects/russianriverecosystemrestorationgi.html>. Accessed: Aug
9 30, 2012. Last update: Jan 2010.
- 10 U.S. Army Corps of Engineers, San Francisco District, Coyote Valley Dam, Lake Mendocino. Viewed
11 online at: <http://www.spn.usace.army.mil/projects/lakemendocino&m.html>. Accessed on: Sept
12 12, 2012. Last update: Jan 2010.
- 13 U.S. Army Corps of Engineers, San Francisco District, Environmental Assessment, Warm Springs Dam
14 and Lake Sonoma Operational Requirements. Viewed online at:
15 [http://www.spn.usace.army.mil/project_programs/Warm%20Springs%20Dam%20Inspection%20](http://www.spn.usace.army.mil/project_programs/Warm%20Springs%20Dam%20Inspection%20Deviations%20Final%20Draft_Redact.pdf)
16 [Deviations%20Final%20Draft_Redact.pdf](http://www.spn.usace.army.mil/project_programs/Warm%20Springs%20Dam%20Inspection%20Deviations%20Final%20Draft_Redact.pdf). Accessed on: Sept 12, 2012. Last update: May 2010.
- 17 U.S. Bureau of Reclamation, 2011a, West-Wide Climate Risk Assessments: Bias-Corrected and Spatially
18 Downscaled Surface Water Projections, Technical Memorandum No. 86-68210-2011-01. Viewed
19 online at: <http://www.usbr.gov/WaterSMART/docs/west-wide-climate-risk-assessments.pdf>.
20 Accessed on: Feb 11, 2013. Last update: Mar 2011.
- 21 U.S. Bureau of Reclamation, 2011b, Klamath Facilities Removal Draft Environmental Impact Statement/
22 Environmental Impact Report, State Clearinghouse # 2010062060. Viewed online at:
23 <http://www.defendruralamerica.com/files/EISEIRAppendix.pdf>. Accessed: Feb 11, 2013. Last
24 update: Sept 2011.
- 25 U.S. Bureau of Reclamation, Basin Report: Klamath River, U.S. Bureau of Reclamation, 6600 Washburn
26 Way, Klamath Falls, OR 97603, 541.884.9053. Viewed online at:
27 <http://www.usbr.gov/climate/SECURE/factsheets/klamath.html>. Accessed: Jun 26, 2012. Last
28 update: Apr 25, 2011.
- 29 U.S. Bureau of Reclamation, Klamath Project 2012 Operations Plan, U.S. Bureau of Reclamation,
30 Klamath Basin Area Office, 6600 Washburn Way Klamath Falls OR 97603 (541) 883-6935.
31 Viewed online at: http://www.usbr.gov/mp/kbao/docs/summer_operations.pdf. Accessed on: Sept
32 12, 2012. Last update: Apr 2012.
- 33 U.S. Court of Appeals, for the District of Columbia Circuit, Arizona Public Service Company v. U.S.
34 Environmental Protection Agency. Viewed online at:
35 [http://caselaw.lp.findlaw.com/scripts/getcase.pl?navby=search&case=/data2/circs/DC/981196a.ht](http://caselaw.lp.findlaw.com/scripts/getcase.pl?navby=search&case=/data2/circs/DC/981196a.html)
36 [ml](http://caselaw.lp.findlaw.com/scripts/getcase.pl?navby=search&case=/data2/circs/DC/981196a.html). Accessed on: Mar 12, 2013. Last update: May 5, 2000.

- 1 U.S. Court of Appeals, Ninth Circuit, State of Montana v. U.S. Environmental Protection Agency, 1998.
2 Viewed online at: *Montana v. US EPA*. Accessed on: Mar 12, 2013. Last update: Mar 3, 1998.
- 3 U.S. Court of Appeals, Seventh Circuit, Wisconsin v. Environmental Protection Agency. Viewed online
4 at: <http://caselaw.lp.findlaw.com/cgi-bin/getcase.pl?court=7th&navby=docket&no=992618>.
5 Accessed on: Mar 12, 2013. Last update: Sept 21, 2001.
- 6 U.S. Department of the Interior, Biological/ Conference Opinion Regarding the Effects of the U.S.
7 Bureau of Reclamation's Proposed 10-Year Operation Plan (Apr 1, 2008 – Mar 31, 2018) for the
8 Klamath Project and its Effects on the Endangered Lost River and Shortnose Suckers, U.S.
9 Department of the Interior, Fish and Wildlife Service, Klamath Falls Fish and Wildlife Office,
10 1936 California Avenue, Klamath Falls, Oregon 97601, (541) 885-8481. Viewed online at:
11 [http://www.fws.gov/klamathfallsfwo/news/2008%20BO/2008_Klamath_Project_Biological_Opi
12 nion_\(Final\).pdf](http://www.fws.gov/klamathfallsfwo/news/2008%20BO/2008_Klamath_Project_Biological_Opinion_(Final).pdf). Accessed on: Sept 12, 2012. Last update: Apr 2008.
- 13 U.S. Department of the Interior Record of Decision, Trinity River Mainstem Fishery Restoration, Final
14 Environmental Impact Statement/ Environmental Impact Report, Dec 2000. Viewed online at:
15 [http://www.schlosserlawfiles.com/TrinityRiver/ROD12-19-00\(b\).pdf](http://www.schlosserlawfiles.com/TrinityRiver/ROD12-19-00(b).pdf). Accessed on: Jul 17, 2012.
16 Last update: Dec 19, 2000.
- 17 U.S. Environmental Protection Agency 2011a, List of Tribes Eligible for Section 319 Funding. Viewed
18 online at: <http://water.epa.gov/polwaste/nps/tribal/upload/tribes319funding.pdf>. Accessed on: Jul
19 25, 2012. Last update: Oct 14, 2011.
- 20 U.S. Environmental Protection Agency 2011b, Region 9, Water Division, Tribal Office Water Pollution
21 Control Program Grants: Clean Water Act §106 (December 1, 2011). Viewed online at:
22 [http://www.epa.gov/region9/water/tribal/pdf/cwa-reporting/2012/FundingOpp-Tribal-
23 WtrPollutCtrl-ProgGrants2012.pdf](http://www.epa.gov/region9/water/tribal/pdf/cwa-reporting/2012/FundingOpp-Tribal-WtrPollutCtrl-ProgGrants2012.pdf). Accessed on: Jul 25, 2012. Last update: Dec 1, 2011.
- 24 U.S. Environmental Protection Agency, Handbook for Developing and Managing Tribal Nonpoint Source
25 Pollution Programs under Section 319 of the Clean Water Act. Office of Water Nonpoint Source
26 Control Branch, 4503(T) 1200 Pennsylvania Avenue, NW Washington, DC 20460. Viewed
27 online at:
28 [http://water.epa.gov/polwaste/nps/tribal/upload/2010_02_19_nps_tribal_pdf_tribal_handbook201
29 0.pdf](http://water.epa.gov/polwaste/nps/tribal/upload/2010_02_19_nps_tribal_pdf_tribal_handbook2010.pdf). Accessed: Sept 6, 2012. Last update: Feb 2010.
- 30 U.S. Environmental Protection Agency, 2012, Water: Small Systems and Capacity Development, Basic
31 Information. Viewed online at:
32 <http://water.epa.gov/type/drink/pws/smallsystems/basicinformation.cfm>. Accessed: Oct 3, 2012.
33 Last update: Mar 6, 2012.
- 34 U.S. Environmental Protection Agency, Region 9 and California Department of Water Resources, 2011.
35 Climate Change Handbook for Regional Water Planning.
36 <http://www.water.ca.gov/climatechange/CCHandbook.cfm>.

- 1 U.S. Fish and Wildlife Service and Hoopa Valley Tribe, 1999, Trinity River Flow Evaluation. Viewed
2 online at:
3 [http://odp.trrp.net/FileDatabase/Documents/USFWS%20HVT%201999%20\(June%20FINAL%20-%20locked\)%20Trinity%20River%20Flow%20Evaluation%20Final%20Report1.pdf](http://odp.trrp.net/FileDatabase/Documents/USFWS%20HVT%201999%20(June%20FINAL%20-%20locked)%20Trinity%20River%20Flow%20Evaluation%20Final%20Report1.pdf).
4
5 Accessed: Mar 7, 2013. Last update: Feb 22, 2013.
- 6 U.S. Fish and Wildlife Service, Klamath Basin National Wildlife Refuge Complex. Viewed online at:
7 <http://www.fws.gov/klamathbasinrefuges>. Accessed: Aug 23, 2012. Last update: Apr 4, 2012.
- 8 U.S. Fish and Wildlife Service, U.S. Bureau of Reclamation, Hoopa Valley Tribe, Trinity County, 2000,
9 Trinity River Mainstem Fishery Restoration Environmental Impact Statement. Viewed online at:
10 <http://odp.trrp.net/Data/Documents/Details.aspx?document=1238>. Accessed: Mar 7, 2013. Last
11 update: Apr 12, 2012.
- 12 U.S. Geological Survey, Smith River Basin Water Data Report, 2009, 11532500 Smith River near
13 Crescent City, California. Viewed online at:
14 <http://wdr.water.usgs.gov/wy2009/pdfs/11532500.2009.pdf>. Accessed: Jun 26, 2012. Last update:
15 2009.
- 16 U.S. Geological Survey c2012, Water Resources Data, USGS Surface-Water Annual Statistics for
17 California USGS 11525500 Trinity R A Lewiston, CA. Viewed online at:
18 http://waterdata.usgs.gov/ca/nwis/annual/?search_site_no=11525500&agency_cd=USGS&referre
19 [d_module=sw&format=sites_selection_links](http://waterdata.usgs.gov/ca/nwis/annual/?search_site_no=11525500&agency_cd=USGS&referred_module=sw&format=sites_selection_links). Accessed on: Feb 11, 2013. Last update: Jan 2013.
- 20 U.S. Nuclear Regulatory Commission, Humboldt Bay Power Plant. Viewed online at:
21 [http://www.nrc.gov/info-finder/decommissioning/power-reactor/humboldt-bay-nuclear-power-](http://www.nrc.gov/info-finder/decommissioning/power-reactor/humboldt-bay-nuclear-power-plant-unit-3.html)
22 [plant-unit-3.html](http://www.nrc.gov/info-finder/decommissioning/power-reactor/humboldt-bay-nuclear-power-plant-unit-3.html). Accessed on: Sept 24, 2012. Last update: Mar 2012.
- 23 United States v. Title Insurance & Trust Company, 1924, U.S. Supreme Court. Viewed online at:
24 <http://caselaw.lp.findlaw.com/scripts/getcase.pl?navby=search&court=US&case=/us/265/472.htm>
25 [l](http://caselaw.lp.findlaw.com/scripts/getcase.pl?navby=search&court=US&case=/us/265/472.html). Accessed on: Mar 13, 2013. Last update: Jun 19, 1924.
- 26 Veteran Homesteaders and Settlers in the Klamath Basin, 2001, Klamath Basin Crisis, PO Box 314,
27 Tulelake, CA 96134. Viewed online at:
28 <http://www.klamathbasin crisis.org/Homesteaders/homesteaders.htm>. Accessed on: Mar 15, 2013.
29 Last update: 2001.
- 30 Western Region Climate Center (WRCC). 2012. Climate Variability in the State of California. Journal of
31 Applied Meteorology and Climatology, 48, 1527-1541. (On-line data citation, accessed
32 10/3/2012: http://www.wrcc.dri.edu/monitor/cal-mon/frames_version.html).
- 33 Wilkinson, Robert. 2000. Methodology for Analysis of the Energy Intensity of California's Water
34 Systems, and An Assessment of Multiple Potential Benefits through Integrated Water-Energy
35 Efficiency Measures. Exploratory Research Project supported by Ernest Orlando Lawrence
36 Berkeley Laboratory, California Institute for Energy Efficiency.

1 Wilkinson, Robert C., Fawzi Karajeh, and Jule Mottin. 2005. Water Sources “Powering” Southern
2 California: Imported Water, Recycled Water, Ground Water, and Desalinated Water. An Analysis
3 of the Energy Intensity of Water Supplies for the West Basin and Central Basin Municipal Water
4 Districts.

5 Winters Doctrine, 1908, Office of the Solicitor, Department of the Interior, 1955, Brigham Young
6 University Law Review, 1975. Viewed online at:
7 <http://www.law2.byu.edu/lawreview/archives/1975/3/ran.pdf>. Accessed on: Mar 14, 2013. Last
8 update: 1975.

9 Yurok Tribe, 2006a, Culture and History. Viewed online at:
10 <http://www.yuroktribe.org/culture/culture.htm>. Accessed Jul 8, 2013. Last update: 2006.

11 Yurok Tribe, 2006b, Background Information. Viewed online at:
12 <http://www.yuroktribe.org/culture/history/history.htm>. Accessed Jul 8, 2013. Last update: 2006.

13 Personal Communications

14 Ford G, Staff Environmental Scientist, Water Branch, California Department of Fish and Game, Email on
15 Jul 16, 2012, Internal memo to B-160 Regional Report authors on priority areas and needs
16 specific to the North Coast Hydrologic Region, from a Department of Fish and Game perspective
17 for California, in relation to California water supply, Communicated thru Lew Moeller.

18
19

Table NC-1 Alluvial Groundwater Basins and Subbasins within the North Coast Hydrologic Region

Basin/Subbasin	Basin Name	Basin/Subbasin	Basin Name
1-1	Smith River Plain	1-33	Larabee Valley
1-2	Klamath River Valley	1-34	Dinsmores Town Area
1-2.01	Tule Lake	1-35	Hyampom Valley
1-2.02	Lower Klamath	1-36	Hettenshaw Valley
1-3	Butte Valley	1-37	Cottoneva Creek Valley
1-4	Shasta Valley	1-38	Lower Laytonville Valley
1-5	Scott River Valley	1-39	Branscomb Town Area
1-6	Hayfork Valley	1-40	Ten Mile River Valley
1-7	Hoopa Valley	1-41	Little Valley
1-8	Mad River Valley	1-42	Sherwood Valley
1-8.01	Mad River Lowland	1-43	Williams Valley
1-8.02	Dows Prairie School Area	1-44	Eden Valley
1-9	Eureka Plain	1-45	Big River Valley
1-10	Eel River Valley	1-46	Navarro River Valley
1-11	Covelo Round Valley	1-48	Gravelly Valley
1-12	Laytonville Valley	1-49	Annapolis Ohlson Ranch Formation Highlands
1-13	Little Lake Valley	1-50	Knights Valley
1-14	Lower Klamath River Valley	1-51	Potter Valley
1-15	Happy Camp Town Area	1-52	Ukiah Valley
1-16	Seiad Valley	1-53	Sanel Valley
1-17	Bray Town Area	1-54	Alexander Valley
1-18	Red Rock Valley	1-54.01	Alexander Area
1-19	Anderson Valley	1-54.02	Cloverdale Area
1-20	Garcia River Valley	1-55	Santa Rosa Valley
1-21	Fort Bragg Terrace Area	1-55.01	Santa Rosa Plain
1-22	Fairchild Swamp Valley	1-55.02	Healdsburg Area
1-25	Prairie Creek Area	1-55.03	Rincon Valley
1-26	Redwood Creek Area	1-56	McDowell Valley
1-27	Big Lagoon Area	1-57	Bodega Bay Area
1-28	Mattole River Valley	1-59	Wilson Grove Formation Highlands
1-29	Honeydew Town Area	1-60	Lower Russian River Valley
1-30	Pepperwood Town Area	1-61	Fort Ross Terrace Deposits
1-31	Weott Town Area	1-62	Wilson Point Area
1-32	Garberville Town Area		

**Table NC-2 Number of Well Logs by County and Use for the North Coast Hydrologic Region
(1977 - 2010)**

County	Total Number of Well Logs by Well Use						Total Well
	Domestic	Irrigation	Public Supply	Industrial	Monitoring	Other	
Del Norte	980	30	20	5	178	57	1,270
Humboldt	647	29	51	7	1,421	189	2,344
Mendocino	5,771	157	119	20	852	163	7,082
Siskiyou	5,120	445	86	20	663	358	6,692
Sonoma	10,750	1,215	366	95	2,878	529	15,833
Trinity	1,442	23	47	3	163	56	1,734
Total Well Records	24,710	1,899	689	150	6,155	1,352	34,955

Table NC-3 CASGEM Groundwater Basin Prioritization for the North Coast Hydrologic Region

Basin Prioritization	Count	Basin/Subbasin Number	Basin Name	Subbasin Name	2010 Census Population
High	0	NA	NA	NA	NA
Medium	1	1-4	SHASTA VALLEY	SHASTA VALLEY	5,333
Medium	2	1-55.01	SANTA ROSA VALLEY	SANTA ROSA PLAIN	250,375
Medium	3	1-1	SMITH RIVER PLAIN		24,588
Medium	4	1-2.01	KLAMATH RIVER VALLEY	TULELAKE	2,261
Medium	5	1-52	UKIAH VALLEY		32,761
Medium	6	1-10	EEL RIVER VALLEY		21,558
Medium	7	1-5	SCOTT RIVER VALLEY		3,520
Medium	8	1-3	BUTTE VALLEY		1,464
Low	2	<i>See Water Plan Update 2013 Vol. 4 Reference Guide – California's Groundwater Update 2013</i>			
Very Low	53	<i>See Water Plan Update 2013 Vol. 4 Reference Guide – California's Groundwater Update 2013</i>			
Totals:	63	Population of GW Basin Area:			550,630

Table NC-4 Groundwater Level Monitoring Wells by Monitoring Entity in the North Coast Hydrologic Region

State and Federal Agencies	Number of Wells
DWR	123
USGS	37
Total State and Federal Wells:	160
Monitoring Cooperators	
N/A	
Total Cooperator Wells:	0
CASGEM Monitoring Entities	
Siskiyou County Public Health and Community Development	5
Sonoma County PRMD	14
Tulelake Irrigation District	15
Total CASGEM Monitoring Entities:	34
Grand Total:	194

Table NC-5 Sources of Groundwater Quality Information for the North Coast Hydrologic Region

Agency	Links to Information
State Water Resources Control Board (http://www.waterboards.ca.gov/)	<p>Groundwater (http://www.waterboards.ca.gov/water_issues/programs/#groundwater)</p> <ul style="list-style-type: none"> • Communities that Rely on a Contaminated Groundwater Source for Drinking Water (http://www.waterboards.ca.gov/water_issues/programs/gama/ab2222/index.shtml) • Hydrogeologically Vulnerable Areas (http://www.waterboards.ca.gov/gama/docs/hva_map_table.pdf) • Aquifer Storage and Recovery (http://www.waterboards.ca.gov/water_issues/programs/asr/index.shtml) <p>GAMA (http://www.waterboards.ca.gov/gama/index.shtml)</p> <ul style="list-style-type: none"> • GeoTracker GAMA (Monitoring Data) (http://www.waterboards.ca.gov/gama/geotracker_gama.shtml) • Domestic Well Project (http://www.waterboards.ca.gov/gama/domestic_well.shtml) • Priority Basin Project (http://www.waterboards.ca.gov/water_issues/programs/gama/sw_basin_assesmt.shtml) • Special Studies Project (http://www.waterboards.ca.gov/water_issues/programs/gama/special_studies.shtml) • California Aquifer Susceptibility Project (http://www.waterboards.ca.gov/water_issues/programs/gama/cas.shtml) <p>Contaminant Sites</p> <ul style="list-style-type: none"> • Land Disposal Program (http://www.waterboards.ca.gov/water_issues/programs/land_disposal/) • Department of Defense Program (http://www.waterboards.ca.gov/water_issues/programs/dept_of_defense/) • Underground Storage Tank Program (http://www.waterboards.ca.gov/ust/index.shtml) • Brownfields (http://www.waterboards.ca.gov/water_issues/programs/brownfields/)
California Department of Public Health (http://www.cdph.ca.gov/Pages/DEFULT.aspx)	<p>Division of Drinking Water and Environmental Management (http://www.cdph.ca.gov/programs/Pages/DDWEM.aspx)</p> <ul style="list-style-type: none"> • Drinking Water Source Assessment and Protection (DWSAP) Program (http://www.cdph.ca.gov/certlic/drinkingwater/Pages/DWSAP.aspx) • Chemicals and Contaminants in Drinking Water (http://www.cdph.ca.gov/certlic/drinkingwater/Pages/Chemicalcontaminants.aspx) • Chromium-6 (http://www.cdph.ca.gov/certlic/drinkingwater/Pages/Chromium6.aspx) • Groundwater Replenishment with Recycled Water (http://www.cdph.ca.gov/HealthInfo/environhealth/water/Pages/Waterrecycling.aspx)

Agency	Links to Information
California Department of Water Resources (http://www.water.ca.gov/)	Groundwater Information Center (http://www.water.ca.gov/groundwater/index.cfm) <ul style="list-style-type: none"> • Bulletin 118 Groundwater Basins (http://www.water.ca.gov/groundwater/bulletin118/gwbasin_maps_descriptions.cfm) • California Statewide Groundwater Elevation Monitoring (CASGEM) (http://www.water.ca.gov/groundwater/casgem/) • Groundwater Level Monitoring (http://www.water.ca.gov/groundwater/data_and_monitoring/gw_level_monitoring.cfm) • Groundwater Quality Monitoring (http://www.water.ca.gov/groundwater/data_and_monitoring/gw_quality_monitoring.cfm) • Well Construction Standards (http://www.water.ca.gov/groundwater/well_info_and_other/well_standards.cfm) • Well Completion Reports (http://www.water.ca.gov/groundwater/well_info_and_other/well_completion_reports.cfm)
California Department of Toxic Substances Control (http://www.dtsc.ca.gov/)	EnviroStor (http://www.envirostor.dtsc.ca.gov/public/)
California Department of Pesticide Regulation (http://www.cdpr.ca.gov/)	Groundwater Protection Program (http://www.cdpr.ca.gov/docs/emon/grndwtr/index.htm) <ul style="list-style-type: none"> • Well Sampling Database (http://www.cdpr.ca.gov/docs/emon/grndwtr/gwp_sampling.htm) • Groundwater Protection Area Maps (http://www.cdpr.ca.gov/docs/emon/grndwtr/gwpa_maps.htm)
U.S. Environmental Protection Agency (http://www.epa.gov/safewater/)	US EPA STORET Environmental Data System (http://www.epa.gov/storet/)
U.S. Geological Survey (http://ca.water.usgs.gov/)	USGS Water Data for the Nation (http://waterdata.usgs.gov/nwis)

Table NC-6 Federally Recognized Tribes in North Coast Hydrologic Region

Name of Tribe	Cultural Affiliation
Bear River Band of the Rohnerville Rancheria	Wiyot, Mattole
Big Lagoon Rancheria	Yurok, Tolowa
Blue Lake Rancheria	Wiyot, Yurok, Hupa
Cahto Indian Tribe of the Laytonville Rancheria	Cahto, Pomo
Cher-Ae Heights Indian Community of the Trinidad Rancheria	Yurok, Wiyot, Tolowa
Cloverdale Rancheria of Pomo Indians of California	Pomo
Coyote Valley Band of Pomo Indians of California	Pomo
Dry Creek Rancheria Band of Pomo Indians of California	Pomo
Elk Valley Rancheria	Tolowa
Federated Indians of Graton Rancheria	Coast Miwok, Southern Pomo
Guidiville Rancheria of California	Pomo
Hoopa Valley Tribe	Hupa
Hopland Band of Pomo Indians of the Hopland Rancheria	Pomo
Karuk Tribe	Karuk
Kashia Band of Pomo Indians of the Stewarts Point Rancheria	Pomo
Lytton Rancheria of California	Pomo
Manchester Band of Pomo Indians of the Manchester-Point Arena Rancheria	Pomo
Pinoleville Pomo Nation	Pomo
Pit River Tribe (Eleven Bands, includes XL Ranch, Big Bend, Likely, Lookout, Montgomery Creek and Roaring Creek Rancherias)	Achomawi (Achumawi, Ajumawi), Aporidge, Astariwawi (Astarawi), Atsuge (Atsugewi), Atwamsini, Hanhawi (Hammawi), Hewisedawi, Ilmawi, Itsatawi, Kosalextawi (Kosalektawi), Madesi
Potter Valley Tribe	Pomo
Quartz Valley Indian Community of the Quartz Valley Indian Reservation	Klamath, Karuk, Shasta
Redwood Valley Rancheria of Pomo Indians	Pomo
Resighini Rancheria	Yurok
Round Valley Indian Tribes of the Round Valley Reservation	Wailacki, Yuki, Pomo, Concow, Nomlacki, Pit River
Sherwood Valley Rancheria of Pomo Indians	Pomo
Smith River Rancheria	Tolowa
Wiyot Tribe	Wiyot
Yurok Tribe of the Yurok Reservation	Yurok

Sources: Department of the Interior, Bureau of Indian Affairs. 2012. Indian Entities Recognized and Eligible To Receive Services From the Bureau of Indian Affairs. [Notice in Federal Register, Vol. 77, No. 155, August 10, 2012.] Viewed online at: <http://www.gpo.gov/fdsys/pkg/FR-2012-08-10/pdf/2012-19588.pdf>, Accessed on August 22, 2012.

Table NC-7 California Native American Tribes (Non-Recognized) in North Coast Hydrologic Region

California Native American Tribe	Cultural Affiliation
Melochundum Band of Tolowa Indians	Tolowa
Eel River Nation of Sovereign Wailaki	Eel River Athapaskans
SheBelNa Band of Mendocino Coast Pomo Indians	Pomo
Noyo River Indian Community	Sinkyone
Yokayo Tribe of Indians	Pomo
Shasta Tribe (Shasta Nation)	Konomihu, New River Indians, Okwanuchu
Mishewal-Wappo Tribe of Alexander Valley	Wappo Indians
Tsnungwe Council	Hupa, South Fork Hupa
Nor-Rel-Muk Nation (formerly Hayfork Band; formerly Nor-El-Muk Band of Wintu Indians)	Wintu

Source: California Native American Heritage Commission, Department of Water Resources

Table NC-8 North Coast Hydrologic Region Average Annual Groundwater Supply by Planning Area (PA) and by Type of Use (2005-2010)

North Coast Hydrologic Region		Agriculture Use Met by Groundwater		Urban Use Met by Groundwater		Managed Wetlands Use Met by Groundwater		Total Water Use Met by Groundwater	
PA Number	PA Name	TAF	%	TAF	%	TAF	%	TAF	%
101	Upper Klamath	182.6	36%	7.4	66%	2.5	1%	192.5	26%
102	Lower Klamath	8.2	30%	5.9	51%	0.0	0%	14.0	36%
103	Coastal	63.9	78%	18.0	37%	0.0	0%	81.9	63%
104	Russian River	46.7	63%	28.7	40%	0.0	0%	75.4	52%
2005-10 Annual Average HR Total:		301.3	44%	60.0	42%	2.5	1%	363.8	35%

Note: 1) TAF = thousand acre-feet

2) Percent use is the percent of the total water supply that is met by groundwater, by type of use.

3) 2005-10 Precipitation equals 99% of the 30-yr average for the North Coast Region

Table NC-9 North Coast Hydrologic Region Average Annual Groundwater Supply by County and by Type of Use (2005-2010)

North Coast Hydrologic Region County	Agriculture Use Met by Groundwater		Urban Use Met by Groundwater		Managed Wetlands Use Met by Groundwater		Total Water Use Met by Groundwater	
	TAF	%	TAF	%	TAF	%	TAF	%
Del Norte	4.3	49%	1.5	37%	0.0	0%	5.8	45%
Siskiyou	175.4	41%	11.8	56%	2.5	2%	189.8	32%
Trinity	2.8	34%	1.5	41%	0.0	0%	4.3	36%
Humboldt	58.8	91%	18.1	42%	0.0	0%	76.9	71%
Mendocino	24.3	48%	7.3	43%	0.0	0%	31.6	47%
Sonoma	43.7	75%	29.3	36%	0.0	0%	73.0	53%
2005-10 Annual Ave. Total:	309.3	50%	69.5	41%	2.5	2%	381.3	41%

Note: 1) TAF = thousand acre-feet

2) Percent use is the percent of the total water supply that is met by groundwater, by type of use.

3) 2005-10 Precipitation equals 99% of the 30-yr average for the North Coast Region

**Table NC-10 Summary of Community Water System Inventory
within the North Coast Hydrologic Region**

Water System Size	Number of Community Systems	Percent of Community Systems in Region	Population Served	Percent of Population Served
Large (> 10,000 population)	11	4	359,575	66
Medium (3,301 – 10,000 pop)	16	6	95,992	18
Small (500 – 3,300 pop)	40	15	57,482	11
Very small < 500 pop)	193	74	28,116	5
Community water systems that primarily provide wholesale water	2	1	---	---
Total	262		541,165	

Source: Water Boards 2012 Draft Report on "Communities that Rely on Contaminated Groundwater"

Note: Sonoma County Water Agency's (System No. 4910020) service area is in both the North Coast and San Francisco Bay Regions. To avoid duplication, it is only included in the North Coast Region.

Table NC-11 North Coast Hydrologic Water Balance Summary, 2001-2010

Table NC-11 North Coast Hydrologic Region water balance for 2001-2010 (in TAF)

North Coast (TAF)	Water Year (Percent of Normal Precipitation)									
	2001 (60%)	2002 (93%)	2003 (99%)	2004 (88%)	2005 (119%)	2006 (135%)	2007 (80%)	2008 (82%)	2009 (76%)	2010 (103%)
Water Entering the Region										
Precipitation	31,254	50,520	53,304	47,461	64,296	72,720	43,139	44,265	40,870	55,352
Inflow from Oregon/Mexico	988	995	1,000	973	909	2,241	1,145	1,182	966	874
Inflow from Colorado River	0	0	0	0	0	0	0	0	0	0
Imports from Other Regions	2	2	2	2	2	2	2	2	2	1
Total	32,244	51,517	54,306	48,436	65,207	74,963	44,286	45,449	41,838	56,227
Water Leaving the Region										
Consumptive Use of Applied Water * (Ag, M&I, Wetlands)	647	876	756	800	617	796	830	778	747	723
Outflow to Oregon/Nevada/Mexico	66	100	72	85	67	123	98	97	53	85
Exports to Other Regions	703	671	895	1,023	498	1,386	648	587	567	299
Statutory Required Outflow to Salt Sink	8,021	18,095	24,375	19,261	17,294	33,462	15,689	16,116	13,583	18,963
Additional Outflow to Salt Sink	122	85	79	75	87	139	95	92	91	74
Evaporation, Evapotranspiration of Native Vegetation, Groundwater Subsurface Outflows, Natural and Incidental Runoff, Ag Effective Precipitation & Other Outflows	23,323	31,929	27,956	27,608	46,660	39,299	27,619	28,415	27,276	35,636
Total	32,882	51,755	54,133	48,852	65,222	75,204	44,979	46,084	42,317	55,780
Change in Supply										
[+] Water added to storage										
[-] Water removed from storage										
Surface Reservoirs	-491	14	414	-166	170	12	-434	-378	-246	667
Groundwater **	-147	-252	-241	-250	-185	-253	-259	-257	-233	-220
Total	-638	-238	173	-416	-15	-241	-693	-635	-479	447
Applied Water * (Ag, Urban, Wetlands) (compare with Consumptive Use)	1,018	1,401	1,220	1,279	1,050	1,375	1,332	1,295	1,212	1,144
* Definition: Consumptive use is the amount of applied water used and no longer available as a source of supply. Applied water is greater than consumptive use because it includes consumptive use, reuse, and outflows.										
** Definition: Change in Supply: Groundwater – The difference between water extracted from and water recharged into groundwater basins in a region. All regions and years were calculated using the following equation:										
change in supply: groundwater = intentional recharge + deep percolation of applied water + conveyance deep percolation and seepage - withdrawals										
This equation does not include unknown factors such as natural recharge and subsurface inflow and outflow. For further details, refer to Volume 4, Reference Guide – <i>California's Groundwater Update 2013</i> and Volume 5 Technical Guide.										
n/a = not applicable										

Table NC-12 Trinity River ROD Water Year Types

Water Year Type	Frequency of Occurrence	Volume (acre-feet)	Peak Release (cubic feet per second)
Critically dry	12%	369,000	1,500
Dry	28%	453,000	4,500
Normal	20%	647,000	6,000
Wet	28%	701,000	8,500
Extremely wet	12%	815,000	11,000

Table NC-13 Trinity River Release and Diversion Summary Since 2000

Water Year	Forecast Water Year Type	Actual Water Year Type	Restoration Water Allocation	Actual Restoration Release	Safety of Dams			Total Release to Trinity River	Peak Release Magnitude	Total Diversion Through Carr Tunnels	Total Inflow into Trinity Reservoir	Notes
					Releases	Releases	Releases					
				(acre-feet)				(cfs)	(acre-feet)			
	[G]	[I]		[A]	[A,B]	[A,B]	[A,B]	[A]	[A]	[C]	[C,F]	
2000	Wet	Wet	340,000	359,600	200,400	-	-	560,000	5,310	1,108,600	1,660,200	[D]
2001	Dry	Dry	369,000	379,600	-	4,200	-	383,800	1,760	669,400	786,700	[E]
2002	Normal	Normal	470,000	482,700	-	-	-	482,700	6,040	629,000	1,243,800	[E]
2003	Wet	Wet	453,000	448,100	68,300	5,700	34,000	556,100	2,610	857,600	1,795,700	[E]
2004	Wet	Wet	646,900	651,000	81,100	-	36,200	768,300	6,200	987,500	1,443,000	[E]
2005	Normal	Wet	647,000	647,600	-	3,600	-	651,200	6,970	466,700	1,412,400	[H]
2006	Ex. Wet	Ex. Wet	815,000	809,900	406,300	-	-	1,216,200	10,100	1,350,600	2,396,500	[H]
2007	Dry	Dry	453,000	453,700	-	4,100	-	457,800	4,750	614,400	715,300	[H]
2008	Normal	Dry	647,000	648,700	-	-	-	648,700	6,470	555,000	835,800	[H]
2009	Dry	Dry	453,000	445,500	-	11,100	-	456,600	4,410	539,200	797,200	[H]
2010	Normal	Wet	647,000	656,700	-	-	-	656,700	6,840	274,700	1,538,000	[H]
TOTAL			5,293,900	5,326,400	756,100	28,700	70,200	6,181,400		7,778,000	13,086,600	

Source: Trinity River Restoration Program, Water Year Summaries, 2011. Viewed online at: <http://www.trrp.net/restore/flows/water-year-summaries/>. Accessed Aug 13, 2013.

Notes: All water volume values are rounded to the nearest 100 acre-feet.

A – Computed from daily average flow record reported by the US Geological Survey for the Trinity River at Lewiston Streamgage # 11525500. The accuracy of the flow records ranges from +/- 5 percent to +/- 15 percent on a time variable basis.

B – Volume estimate for flows above the summer or winter baseflow release for restoration.

C – Computed from daily average record provided by the Bureau of Reclamation. Reported negative daily inflow values included “as is” in calculations.

D – Water allocation prior to implementation of the 2000 Trinity River Mainstem Fishery Restoration Record of Decision.

E – Water allocation limited by Court order 2001-2004. Court ordered volumes varied by year.

F – Long-term average annual inflow to Trinity Reservoir (acre-feet/year) from 1911-2007 as provided by the U.S. Bureau of Reclamation = 1,254,000

G – Water year type based on the April forecast (50% exceedance) from the Bulletin 120 – Water Conditions in California by the California Department of Water Resources

H – Restoration water allocation as prescribed by the Trinity River Mainstem Fishery Restoration 2000 Record of Decision.

Table NC-14 Summary of Community Drinking Water Systems in the North Coast Hydrologic Region that Rely on One or More Contaminated Groundwater Well that Exceeds a Primary Drinking Water Standard

• Community Drinking Water Systems and Groundwater Wells Grouped by Water System Population	• No. of Affected Community Drinking Water Systems	• No. of Affected Community Drinking Water Wells
Small System ≤ 3,300	11	14
Medium System 3,301 – 10,000	2	4
Large System > 10,000	2	3
Total	15	21

Source: Water Boards 2012 Draft Report on "Communities that Rely on Contaminated Groundwater"

Table NC-15 Summary of Contaminants Affecting Community Drinking Water Systems in the North Coast Hydrologic Region

Principal Contaminant (PC)	Community Drinking Water Systems where PC exceeds the Primary MCL	No. of Community Drinking Water Wells where PC exceeds the Primary MCL
Arsenic	12	16
Trichloroethylene (TCE)	2	2
Nitrate	1	3
1,1-Dichloroethylene (1,1-DCE)	1	1

Source: Water Boards 2012 Draft Report on "Communities that Rely on Contaminated Groundwater"

Table NC-16 North Coast Hydrologic Region Water Management Agencies

Name	County	Type	Statutory Authority^a
Albion Mutual Water Company	Mendocino	Private water district	WS
Alderpoint County Water District	Humboldt	Special district	WS
Alexander Valley Acres Water Company	Sonoma	Private water district	WS
Arcata City Wastewater Treatment Plant	Humboldt	Municipal/domestic WWTF	WT
Armstrong Valley Water Company	Sonoma	Ngo	WS
Austin Acres Mutual Water Company	Sonoma	Private water district	WS
Austin Creek Mutual (Springhill)	Sonoma	Private water district	WS
Belmont Terrace Mutual Water Company	Sonoma	Private water district	WS
Benbow Water Corporation	Humboldt	Investor-Owned Water Co. - Class D (<500 connections)	WS
Bennett Ridge Mutual Water Company	Sonoma	Private water district	WS
Bertsch-Oceanview C.S.D.	Del Norte	Private water district	WS
Big Lagoon CSD	Humboldt	Private water district	WS
Big Lagoon Park Water Co.	Humboldt	Private water district	WS
Big River Vista Mutual Water Company	Mendocino	Private water district	WS
Big Springs Irrig. District	Siskiyou	Private water district	WS
Big Springs Irrigation District	Siskiyou	Irrigation district	IWS
Blue Lake City Publicly Owned Treatment Works	Humboldt	Municipal/domestic WWTF	WT
Bodega Bay Public Utilities District	Sonoma	Special district	WS
Bodega Bay Wastewater Rec.Fac.	Sonoma	Municipal/domestic WWTF	WT
Bodega Water Company	Sonoma	Private water district	WS
Branger Mutual Water Company, Inc.	Sonoma	Private water district	WS
Brooktrails Township C.S.D.	Mendocino	Private water district	WS
Bucher Water Company	Sonoma	Private water district	WS
Bucktail Mutual Water Company	Trinity	Private water district	WS
Butte Valley Irrigation District	Siskiyou	Federal water contractors service areas	WS
California American Water	Humboldt / Sonoma	Investor-Owned Water Co. - Class A (>10,000 connections)	WS
California Water Service Company	Sonoma	Investor-Owned Water Co. - Class A (>10,000 connections)	WS
Calpella County Water District	Mendocino	Special district	WS
Calpella Cwd-Wastewater Treatment Plant	Mendocino	Municipal/domestic WWTF	WT
Carmet By the Sea Water Company	Sonoma	Private water district	WS
Cazadero Water Company	Sonoma	Investor-Owned Water Co. - Class D (<500 connections)	WS
CDC Pelican Bay Prison Wastewater Treatment Plant	Del Norte	Municipal/domestic WWTF	WT
Church Tree C.S.D.	Del Norte	Private water district	WS
City of Arcata	Humboldt	City	WS
City Of Blue Lake W.S.A.	Humboldt	Private water district	WS
City Of Cloverdale W.S.A.	Sonoma	Private water district	WS
City of Cotati	Sonoma	City	WS

Name	County	Type	Statutory Authority^a
City of Cotati	Sonoma	Municipal/domestic WWTF	WT
City Of Dorris	Siskiyou	Private water district	WS
City Of Eureka W.S.A.	Humboldt	Private water district	WS
City Of Fort Bragg W.S.A.	Mendocino	Private water district	WS
City Of Fortuna W.S.A.	Humboldt	Private water district	WS
City Of Healdsburg W.S.A.	Sonoma	Private water district	WS
City of Rohnert Park	Sonoma	Municipal/domestic WWTF	WT
City Of Rohnert Park W.S.A.	Sonoma	Private water district	WS
City of Santa Rosa	Sonoma	City	WS
City Of Sebastopol W.S.A.	Sonoma	Private water district	WS
City of Trinidad C.S.D.	Humboldt	Private water district	WS
Clear Lake National Wildlife Refuge	Modoc	Federal water district	WS
Cloverdale City Wastewater Treatment Plant	Sonoma	Municipal/domestic WWTF	WT
College Of The Redwoods, Publicly Owned Treatment Works	Humboldt	Municipal/domestic WWTF	WT
Colonial Realty I.D.	Siskiyou	Federal water contractors service areas	WS
Copco Lake MWC	Siskiyou	Private water district	WS
Covelo Community Services District; Covelo City Publicly Owned Treatment Works	Mendocino	Municipal/domestic WWTF	WT
Crescent City Wastewater Treatment Plant	Del Norte	Municipal/domestic WWTF	WT
Crescent City Water District	Del Norte	Private water district	WS
Del Norte County Flood Control District	Del Norte	Flood control district	FC
Del Oro Water Co.-Ferndale	Humboldt	Private water district	WS
Dorris City Sewage Treatment Plant	Siskiyou	Municipal/domestic WWTF	WT
Elk County W.D.	Mendocino	Private water district	WS
Etna CSD	Siskiyou	Irrigation district	IWS
Etna Sewage Treatment Plant	Siskiyou	Municipal/domestic WWTF	WT
Eureka City Elk River Wastewater Treatment Plant	Humboldt	Municipal/domestic WWTF	WT
Ferndale City Publicly Owned Treatment Works	Humboldt	Municipal/domestic WWTF	WT
Fieldbrook C.S.D.	Humboldt	Private water district	WS
Forestville County Water District	Sonoma	Special district	WS
Forestville Water District	Sonoma	Private water district	WS
Fort Bragg City Wastewater Treatment Plant	Mendocino	Municipal/domestic WWTF	WT
Fort Jones City Wastewater Treatment Plant	Siskiyou	Municipal/domestic WWTF	WT
Fortuna City Wastewater Treatment Plant	Humboldt	Municipal/domestic WWTF	WT
Francis Land and Water Company	Humboldt	Investor-Owned Water Co. - Class C (500-2,000 connections)	WS
Garberville Publicly Owned Treatment Works	Humboldt	Municipal/domestic WWTF	WT
Garberville Water Company	Humboldt	Ngo	WS
Gasquet C.S.D.	Del Norte	Private water district	WS
Geyserville Water Works	Sonoma	Investor-Owned Water Co. - Class D (<500 connections)	WS
Gill Creek Mutual Water Company	Sonoma	Private water district	WS

Name	County	Type	Statutory Authority^a
Graton Community Service District	Sonoma	Private water district	WS
Grenada I.D.	Siskiyou	Private water district	WS
Grenada Irrigation District	Siskiyou	Irrigation district	IWS
Grenada Sd Sewage Treatment Plant	Siskiyou	Municipal/domestic WWTF	WT
Happy Camp C.S.D.	Siskiyou	Private water district	WS
Happy Camp Wastewater Treatment Plant	Siskiyou	Municipal/domestic WWTF	WT
Hayfork Wastewater Facilities	Trinity	Municipal/domestic WWTF	WT
Healdsburg City Wastewater Treatment Plant	Sonoma	Municipal/domestic WWTF	WT
Hidden Valley Lake CSD	Humboldt	Private water district	WS
Hoopa Valley CSD	Humboldt	Private water district	WT
Hopland Public Utility Dist.	Mendocino	Private water district	WS
Hornbrook C.S.D.	Siskiyou	Private water district	WS
Humboldt Bay Municipal Water District	Humboldt	Special district	WS
Humboldt Bay Recreation & Conservation District	Humboldt	County-wide agency	WS
Humboldt Community Services District	Humboldt	Special district	WS
Humboldt County Flood Control District	Humboldt	Flood control district	FC
Hydesville County W.D.	Humboldt	Private water district	WS
Irish Beach Water District	Mendocino	Private water district	WS
Klamath C.S.D.	Del Norte	Private water district	WS
Klamath Sewage Treatment Plant	Del Norte	Municipal/domestic WWTF	WT
Lake Shastina Community Service District Sewage Treatment Plant	Siskiyou	Municipal/domestic WWTF	WT
Lake Shastina Mutual Water District	Siskiyou	Special district	WS
Laytonville Water District	Mendocino	Special district	WS
Lewiston Valley Water Co Publicly Owned Treatment Works	Trinity	Municipal/domestic WWTF	WT
Loleta C.S.D.	Humboldt	Private water district	WS
Loleta Publicly Owned Treatment Works	Humboldt	Municipal/domestic WWTF	WT
Lower Klamath National Wildlife Refuge	Siskiyou	Federal water district	WS
Lower Tule River Irrigation District	Siskiyou	Special district	WS
MacDoel Water Works	Siskiyou	Investor-Owned Water Co. - Class D (<500 connections)	WS
Manila Community Service District Wastewater Treatment Plant	Humboldt	Municipal/domestic WWTF	WT
Mayacama Golf Club, LLC	Sonoma	Investor-Owned Water Co. - Class D (<500 connections)	WS
McKinleyville C.S.D.	Humboldt	Private water district	WS
McKinleyville Wastewater Treatment Plant	Humboldt	Municipal/domestic WWTF	WT
Mendocino City Community Service District & High School	Mendocino	Municipal/domestic WWTF	WT
Mendocino County Russian River Flood Control and Water Conservation Improvement District	Mendocino	Flood control district	FC
Mendocino County Water Agency	Mendocino	Special district	WS

Name	County	Type	Statutory Authority^a
Mendocino County Water Works District, Gualala Community Services District, Gualala Wastewater Treatment Plant	Mendocino	Municipal/domestic WWTF	WT
Mendocino Inland Water and Power Commission	Mendocino	Special district	WS
Millview County W.D.	Mendocino	Private water district	WS
Miranda C.S.D.	Humboldt	Private water district	WS
Miranda Publicly Owned Treatment Works	Humboldt	Municipal/domestic WWTF	WT
Montague Sewage Treatment Plant	Siskiyou	Municipal/domestic WWTF	WT
Montague Water Conservation District	Siskiyou	Special district	WS
Montair Subdivision, Sewage Treatment Plant	Siskiyou	Municipal/domestic WWTF	WT
Myers Flat Mutual Water System	Humboldt	Special district	WS
Newell CWD Sewage Treatment Plant	Siskiyou	Municipal/domestic WWTF	WT
North Gualala Water Works	Mendocino	Investor-Owned Water Co. - Class C (500-2,000 connections)	WS
North Marin Water District	Marin	Special district	WS
Occidental Community Services District	Sonoma	Private water district	WS
Odd Fellows Wastewater Treatment Plant	Sonoma	Municipal/domestic WWTF	WT
Orick C.S.D.	Humboldt	Private water district	WS
P Canal	Siskiyou	Private water district	WS
Point Arena Wastewater Treatment Plant	Mendocino	Municipal/domestic WWTF	WT
Point Arena Water Works, Inc.	Mendocino	Investor-Owned Water C. - Class D (<500 connect)	WS
Potter Valley Irrigation Dist.	Mendocino	Private water district	WS
Redway Community Service District.	Humboldt	Private water district	WS
Redway Publicly Owned Treatment Works	Humboldt	Municipal/domestic WWTF	WT
Redwood Park Sewage Treatment Plant	Del Norte	Municipal/domestic WWTF	WT
Redwood Valley County Water District	Mendocino	Special district	WS
Redwood Water Company, Inc.	Sonoma	Investor-Owned Water Co. - Class D (<500 connections)	WS
Resort Improvement District #1	Humboldt	Private water district	WS
Rio Dell City Wastewater Treatment Plant	Humboldt	Municipal/domestic WWTF	WT
Riverview Acres Water Company	Trinity	Investor-Owned Water Co. - Class D (<500 connections)	WS
Rogina Water Company	Mendocino	Investor-Owned Water Co. - Class C (500-2,000 connections)	WS
Round Valley Community Sewer System	Mendocino	Municipal/domestic WWTF	WT
Russian River County W.D.	Sonoma	Private water district	WS
Santa Rosa City Wastewater Treatment Plant, Laguna	Sonoma	Municipal/domestic WWTF	WT
Scott Valley Irrigation District	Siskiyou	Irrigation district	IWS
SCWA Airport Water Reclamation Facility	Sonoma	Municipal/domestic WWTF	WT
SCWA Geyserville Community Service District	Sonoma	Municipal/domestic WWTF	WT
SCWA Occidental Community Service District	Sonoma	Municipal/domestic WWTF	WT
SCWA Russian River Community Service District	Sonoma	Municipal/domestic WWTF	WT

Name	County	Type	Statutory Authority^a
Sea Ranch Water Company	Sonoma	Investor-Owned Water Co. - Class C (500-2,000 connections)	WS
Sereno Del Mar Water Company	Sonoma	Investor-Owned Water C. Class D (<500 connect)	WS
Shelter Cove Publicly Owned Treatment Works	Mendocino	Municipal/domestic WWTF	WT
Siskiyou County Flood Control and Water Conservation District	Siskiyou	Flood control district	FC
Smith River C.S.D.	Del Norte	Private water district	WS
Sonoma County Mutual Water Company	Sonoma	Private water district	WS
Sonoma County Water Agency	Sonoma	Special district	WS
Sonoma County Water Agency	Sonoma	Flood control district	FC
SSU Wastewater Equalization Tank	Sonoma	Municipal/domestic WWTF	WT
Sweetwater Springs CWD - Guerneville	Sonoma	Private water district	WS
Tennant Community Service District St/Lf	Siskiyou	Municipal/domestic WWTF	WT
Town Of Windsor W.S.A	Sonoma	Private water district	WS
Trinity County Water Works District #1	Trinity	Municipal/domestic WWTF	WT
Trinity Village Water Company	Trinity	Investor-Owned Water Co. - Class D (<500 connections)	WS
Tulelake City Wastewater Treatment Plant	Siskiyou	Municipal/domestic WWTF	WT
Tulelake Irrigation District	Siskiyou/M odoc	Federal water district	WS
Tulelake National Wildlife Refuge	Siskiyou/M odoc	Federal water district	WS
Ukiah City Wastewater Treatment Plant	Mendocino	Municipal/domestic WWTF	WT
Ukiah Water District	Mendocino	Private water district	WS
USFS Orleans R.S. Sewage Treatment Plant	Humboldt	Municipal/domestic WWTF	WT
Weaverville C.S.D.	Trinity	Private water district	WS
Weaverville SD Wastewater Treatment Plant	Trinity	Municipal/domestic WWTF	WT
Weed Shastina Wastewater Treatment Plant	Siskiyou	Municipal/domestic WWTF	WT
Weed Wastewater Treatment Plant	Siskiyou	Municipal/domestic WWTF	WT
Wendell Water Company	Sonoma	Investor-Owned Water Co. - Class D (<500 connections)	WS
Weott C.S.D	Humboldt	Private water district	WS
Weott Wastewater Treatment Plant	Humboldt	Municipal/domestic WWTF	WT
Wesewage Treatment Plant Land App For Biosolid	Mendocino	Municipal/domestic WWTF	WT
Wesewage Treatment Plantort CWD	Mendocino	Municipal/domestic WWTF	WT
West Water Company	Sonoma	Investor-Owned Water Co. - Class D (<500 connections)	WS
Westhaven C.S.D.	Humboldt	Private water district	WS
Westport County Water District	Mendocino	Private water district	WS
Willits City Wastewater Treatment Plant	Mendocino	Municipal/domestic WWTF	WT
Willow County Water District	Mendocino	Private water district	WS
Willow Creek Community Services District	Humboldt	Special district	WS
Windsor, Town Of Wastewater Treatment Plant	Sonoma	Municipal/domestic WWTF	WT
Yokayo Water System	Mendocino	Private water district	WS

Name	County	Type	Statutory Authority ^a
Yreka City Wastewater Treatment Plant	Siskiyou	Municipal/domestic WWTF	WT
Yulupa Mutual Water Company	Sonoma	Private water district	WS
Yurok Tribe Public Utilities District	Humboldt	Private water district	WS

Source: County of Humboldt, Community Development Services, Integrated Regional Water Management (IRWM) Region Acceptance Process (RAP).

^a WS = water supply, WT = wastewater treatment, IWS = irrigation water supply, FC = flood control.

Table NC-17 State Funding Received

Funding Received	Description
\$3,394,652	Prop 1E 2011 City of Fortuna Rohner Creek Flood Control and Riparian Habitat Improvement Project
\$500,000	Prop 50 2011 planning grant for NCIRWM DAC Pilot Project to Humboldt County to administer for Region
\$24,831,579	Prop 50 2009 Round 1 Implementation Grant (State Water Board) for Humboldt County to administer for Region; 21 projects in 7 counties
\$2,176,860	Prop 50 2010 Supplemental for Coastal Implementation Projects administered by Humboldt County for Region
\$160,000	Prop 50 2007 Scott River IRWM Implementation Grant (delayed due to economic constraints at State level)
\$50,000	Prop 50 2007 Local Groundwater Assistance grant to Ukiah for groundwater management plan development
\$1,000,000	Prop 84 2010 planning grant for NCIRWMP Phase III work on plan
\$500,000	Prop 84 2011 DAC Pilot Project administered by Humboldt County for Region
\$8,221,061	Prop 84 2011 Round 1 Implementation grant for 19 projects in Region

Source: California Department of Water Resources, 2013.

Table NC-18 Groundwater Management Plans in the North Coast Hydrologic Region

Map Label	Agency Name	Date	Coverage Area		
			County	Basin Number	Basin Name
NC-1	Humboldt Bay Municipal WD	2006	Humboldt	1-8.01	Mad River Lowland Subbasin
	No signatories on file				
NC-2	Mendocino City CSD	2007	Mendocino	1-21	Fort Bragg Terrace Area Basin
	No signatories on file			–	Non-B118 Basin
SR-14	Glenn County	2009	Glenn	5.21.52	Colusa Subbasin
	Provident ID			5-21.58	West Butte Subbasin
	Glide WD			5.21.51	Corning Subbasin
	Willow Creek MWC			5.61	Chrome Town Basin
	California Water Service			5-62	Elk Creek Area Basin
	Princeton-Codora-Glenn ID, Provident ID			5-63	Stonyford Town Area
	Kanawha WD			5-88	Stony Gorge Reservoir Basin
	Glenn-Colusa ID			5-89	Squaw Flat Basin
	Orland-Artois WD			5-90	Funks Creek Basin
	Western Canal			–	Non-B118 Basin
SR-15	Lake County Watershed Protection District	2006	Lake	5-13	Upper Lake Valley
	No signatories on file			5-14	Scotts Valley
				5-16	High Valley
				5-17	Burns Valley
				5-18	Coyote Valley
				5-19	Collayomi Valley
				5-30	Lower Lake Valley
				5-31	Long Valley
				5-66	Clear Lake Cache Formation
				5-94	Middle Creek
				1-48	Gravelley Valley (NC)

Table NC-19 Assessment for SB 1938 GWMP Required Components, SB 1938 GWMP Voluntary Components, and Bulletin 118-03 Recommended Components

SB 1938 GWMP Required Components	Percent of Plans that Meet Requirement
Basin Management Objectives	25%
BMO: Monitoring/Management Groundwater Levels	100%
BMO: Monitoring Groundwater Quality	100%
BMO: Inelastic Subsidence	75%
BMO: SW/GW Interaction & Affects to Groundwater Levels & Quality	50%
Agency Cooperation	50%
Map	75%
Map: Groundwater basin area	100%
Map: Area of local agency	100%
Map: Boundaries of other local agencies	75%
Recharge Areas (1/1/2013)	Not Assessed
Monitoring Protocols	0%
MP: Changes in groundwater levels	100%
MP: Changes in groundwater quality	75%
MP: Subsidence	50%
MP: SW/GW Interaction & Affects to Groundwater Levels & Quality	25%
Met all required components, and subcomponents:	0%
SB 1938 Voluntary Components	Percent of Plans that Include Component
Saline Intrusion	50%
Wellhead Protection & Recharge	50%
Groundwater Contamination	75%
Well Abandonment & Destruction	75%
Overdraft	75%
Groundwater Extraction & Replenishment	50%
Monitoring	100%
Conjunctive Use Operations	25%
Well Construction Policies	75%
Construction and Operation	75%
Regulatory Agencies	75%
Land Use	25%
Bulletin 118-03 Recommended Components	Percent of Plans that Include Component
GMP Guidance	50%
Management Area	75%
BMOs, Goals, & Actions	50%
Monitoring Plan Description	75%
IRWM Planning	25%
GMP Implementation	75%
GMP Evaluation	75%

Table NC-20 Factors Contributing to Successful Groundwater Management Plan Implementation in the North Coast Hydrologic Region

Key Components	Respondents
Data collection and sharing	1
Outreach and education	-
Developing an understanding of common interest	1
Sharing of ideas and information with other water resource managers	1
Broad stakeholder participation	1
Adequate surface water supplies	1
Adequate regional and local surface storage and conveyance systems	1
Water budget	-
Funding	1
Time	1

Table NC-21 Factors Limiting Successful Groundwater Management Plan Implementation in the North Coast Hydrologic Region

Limiting Factors	Respondents
Funding for groundwater management projects	1
Funding for groundwater management planning	1
Unregulated Pumping	1
Groundwater Supply	-
Participation across a broad distribution of interests	-
Lack of Governance	-
Surface storage and conveyance capacity	-
Understanding of the local issues	-
Access to planning tools	1
Outreach and education	-
Data collection and sharing	1
Funding to assist in stakeholder participation	1

Table NC-22 Groundwater Ordinances that Apply to Counties in the North Coast Hydrologic Region

County	Groundwater Management	Guidance Committees	Export Permits	Recharge	Well Abandonment and Destruction	Well Construction Policies
Del Norte	-	-	-	-	Y	-
Glenn	Y	Y	-	-	Y	Y
Humboldt	-	-	-	-	-	Y
Lake	-	-	Y	-	Y	Y
Mendocino	-	-	-	-	Y	Y
Modoc	-	-	Y	-	-	Y
Siskiyou	-	Y	Y	-	Y	-
Sonoma	-	-	-	-	Y	Y
Trinity	-	-	-	-	-	Y

Table NC-23 Groundwater Adjudications in the North Coast Hydrologic Region

Court Judgment	North Coast HR Basin/Subbasin	Basin Number	County	Judgment Date
Scott River Stream System	Scott River Valley Basin	1-5	Siskiyou	1980

Note: Table represents information as of April, 2013

Table NC-24 Conceptual Growth Scenarios

Scenario	Population Growth	Development Density
LOP-HID	Lower than Current Trends	Higher than Current Trends
LOP-CTD	Lower than Current Trend	Current Trends
LOP-LOD	Lower than Current Trends	Lower than Current Trends
CTP-HID	Current Trends	Higher than Current Trends
CTP-CTD	Current Trends	Current Trends
CTP-LOD	Current Trends	Lower than Current Trends
HIP-HID	Higher than Current Trends	Higher than Current Trends
HIP-CTD	Higher than Current Trends	Current Trends
HIP-LOD	Higher than Current Trends	Lower than Current Trends

Source: California Department of Water Resources 2012.

Table NC-25 Growth Scenarios (Urban) — North Coast

Scenario ^a	2050 Population (thousand)	Population Change (thousand) 2006^b to 2050	Development Density	2050 Urban Footprint (thousand acres)	Urban Footprint Increase (thousand acres) 2006^c to 2050
LOP-HID	763.3 ^d	106.7	High	204.3	15.9
LOP-CTD	763.3	106.7	Current Trends	206.5	18.1
LOP-LOD	763.3	106.7	Low	208.5	20.1
CTP-HID	814.9 ^e	14.9	High	219.4	31.0
CTP-CTD	814.9	14.9	Current Trends	221.8	33.4
CTP-LOD	814.9	14.9	Low	224.6	36.2
HIP-HID	1,185.6 ^f	33.3	High	267.5	79.1
HIP-CTD	1,185.6	33.3	Current Trends	278.0	89.6
HIP-LOD	1,185.6	33.3	Low	288.6	100.2

Source: California Department of Water Resources 2012.

Notes:

^a See Table NC-24 for scenario definitions

^b 2006 population was 656.6 thousand.

^c 2006 urban footprint was 188.4 thousand acres.

^d Values modified by the California Department of Water Resources (DWR) from the Public Policy Institute of California.

^e Values provided by the California Department of Finance.

^f Values modified by DWR from the Public Policy Institute of California.

Table NC-26 Growth Scenarios (Agriculture) — North Coast

Scenario ^a	2050 Irrigated Land Area ^b (thousand acres)	2050 Irrigated Crop Area ^c (thousand acres)	2050	Change in Irrigated Crop Area (thousand acres) 2006 to 2050
			Multiple Crop Area ^d (thousand acres)	
LOP-HID	325.1	325.1	0.0	-37.8
LOP-CTD	324.7	324.7	0.0	-38.2
LOP-LOD	324.4	324.4	0.0	-38.5
CTP-HID	322.4	322.4	0.0	-40.5
CTP-CTD	322.0	322.0	0.0	-40.9
CTP-LOD	321.4	321.4	0.0	-41.5
HIP-HID	314.4	314.4	0.0	-48.4
HIP-CTD	312.3	312.3	0.0	-50.6
HIP-LOD	310.2	310.2	0.0	-52.7

Source: California Department of Water Resources 2012

Notes:

^a See Table NC-24 for scenario definitions

^b 2006 Irrigated land area was estimated by the California Department of Water Resources (DWR) to be 362.9 thousand acres.

^c 2006 Irrigated crop area was estimated by DWR to be 362.9 thousand acres.

^d 2006 multiple crop area was estimated by DWR to be 0.0 thousand acres.

Table NC-27 Resource Management Strategies Addressed in IRWMPs in the North Coast Hydrologic Region

Resource Management Strategy	IRWMP 1	IRWMP 2
Agricultural Water Use Efficiency		
Urban Water Use Efficiency		
Conveyance – Delta		
Conveyance – Regional/Local		
System Reoperation		
Water Transfers		
Conjunctive Management & Groundwater		
Desalination		
Precipitation Enhancement		
Recycled Municipal Water		
Surface Storage – CALFED		
Surface Storage – Regional/Local		
Drinking Water Treatment and Distribution		
Groundwater and Aquifer Remediation		
Match Water Quality to Use		
Pollution Prevention		
Salt and Salinity Management		
Agricultural Lands Stewardship		
Economic Incentives		
Ecosystem Restoration		
Forest Management		
Land Use Planning and Management		
Recharge Areas Protection		
Water-Dependent Recreation		
Watershed Management		
Flood Risk Management		
Flood Management		
Desalination (Brackish and Sea Water)		
Salt and Salinity Management		

Figure NC-1 North Coast Hydrologic Region



Figure NC-2 North Coast Hydrologic Region Watersheds



Figure NC-3 Alluvial Groundwater Basins and Subbasins within the North Coast Hydrologic Region



Figure NC-4 Number of Well Logs by County and Use for the North Coast Hydrologic Region (1977–2010)




Figure NC-5 Percentage of Well Logs by Use for the North Coast Hydrologic Region (1977–2010)




Figure NC-6 Number of Well Logs Filed per Year by Use for the North Coast Hydrologic Region (1977–2010)




Figure NC-7 CASGEM Groundwater Basin Prioritization for the North Coast Hydrologic Region




Figure NC-8 Monitoring Well Location by Agency, Monitoring Cooperator, and CASGEM Monitoring Entity in the North Coast Hydrologic Region





Figure NC-9 Percentage of Monitoring Wells by Use in the North Coast Hydrologic Region



PLACEHOLDER Figure NC-10 Total Population 1960-2010 (2050), North Coast Hydrologic Region

[figure to come]

PLACEHOLDER Figure NC-11 Population Growth Trends, North Coast Hydrologic Region

[figure to come]

Figure NC-12 Disadvantaged Communities (DAC) in the North Coast Hydrologic Region




Figure NC-13 North Coast Hydrologic Region Inflows and Outflows in 2010




Figure NC-14 Contribution of Groundwater to the North Coast Hydrologic Region Water Supply by Planning Area (2005-2010)




Figure NC-15 North Coast Hydrologic Region Annual Groundwater Water Supply Trend (2002-2010)




Figure NC-16 North Coast Hydrologic Region Annual Groundwater Supply Trend by Type of Use (2002-2010)




Figure NC-17 North Coast Hydrologic Region Water Balance by Water Year, 2001-2010

California's water resources vary significantly from year to year. Ten recent years show this variability for water use and water supply. Applied Water Use shows how water is applied to urban and agricultural sectors and dedicated to the environment and the Dedicated and Developed Water Supply shows where the water came from each year to meet those uses. Dedicated and Developed Water Supply does not include the approximately 125 million acre-feet (MAF) of statewide precipitation and inflow in an average year that either evaporates, are used by native vegetation, provides rainfall for agriculture and managed wetlands, or flow out of the state or to salt sinks like saline aquifers. Groundwater extraction includes annually about 2 MAF more groundwater used statewide than what naturally recharges – called groundwater overdraft. Overdraft is characterized by groundwater levels that decline over a period of years and never fully recover, even in wet years.




Figure NC-17 North Coast Hydrologic Region Water Balance by Water Year, 2001-2010 (continued)**Key Water Supply and Water Use Definitions**

Applied water. The total amount of water that is diverted from any source to meet the demands of water users without adjusting for water that is depleted, returned to the developed supply or considered irrecoverable (see water balance figure).

Consumptive use is the amount of applied water used and no longer available as a source of supply. Applied water is greater than consumptive use because it includes consumptive use, reuse, and outflows.

Instream environmental. Instream flows used only for environmental purposes.

Instream flow. The use of water within its natural watercourse as specified in an agreement, water rights permit, court order, FERC license, etc.

Groundwater Extraction. An annual estimate of water withdrawn from banked, adjudicated, and unadjudicated groundwater basins.

Recycled water. Municipal water which, as a result of treatment of waste, is suitable for a direct beneficial use or a controlled use that would not otherwise occur and is therefore considered a valuable resource.

Reused water. The application of previously used water to meet a beneficial use, whether treated or not prior to the subsequent use.

Urban water use. The use of water for urban purposes, including residential, commercial, industrial, recreation, energy production, military, and institutional classes. The term is applied in the sense that it is a kind of use rather than a place of use.

Water balance. An analysis of the total developed/dedicated supplies, uses, and operational characteristics for a region. It shows what water was applied to actual uses so that use equals supply.

North Coast Water Balance by Water Year Data Table (MAF)

	2001 (60%)	2002 (93%)	2003 (99%)	2004 (88%)	2005 (119%)	2006 (135%)	2007 (80%)	2008 (82%)	2009 (76%)	2010 (103%)
Applied Water Use										
Urban	149	152	155	157	155	153	167	160	161	153
Irrigated Agriculture	633	942	811	891	663	884	931	903	851	824
Managed Wetlands	254	345	304	301	291	399	311	311	281	225
Req Delta Outflow	0	0	0	0	0	0	0	0	0	0
Instream Flow	1,474	1,422	1,617	1,711	1,613	1,968	1,596	1,533	1,558	1,801
Wild & Scenic R.	6,548	16,697	22,783	17,752	15,866	31,583	14,132	14,632	12,075	17,274
Total Uses	9,057	19,557	25,669	20,811	18,587	34,987	17,137	17,539	14,926	20,276
Depleted Water Use (stippling)										
Urban	127	97	96	99	101	65	100	84	90	83
Irrigated Agriculture	516	708	612	656	485	667	694	665	621	611
Managed Wetlands	223	249	215	209	184	299	239	217	199	163
Req Delta Outflow	0	0	0	0	0	0	0	0	0	0
Instream Flow	1,474	1,398	1,593	1,509	1,428	1,879	1,557	1,484	1,508	1,689
Wild & Scenic R.	6,548	16,697	22,783	17,752	15,866	31,583	14,132	14,632	12,075	17,274
Total Uses	8,888	19,148	25,298	20,225	18,064	34,493	16,721	17,082	14,493	19,819
Dedicated and Developed Water Supply										
Instream	7,934	18,095	24,375	19,261	17,294	33,443	15,689	16,116	13,583	18,962
Local Projects	341	417	391	384	329	361	382	377	369	394
Local Imported Deliveries	18					38	41	3	2	1
Colorado Project	0	0	0	0	0	0	0	0	0	0
Federal Projects	238	336	274	304	241	338	336	316	297	173
State Project	0	0	0	0	0	0	0	0	0	0
Groundwater Extraction	453	386	364	391	298	388	396	395	365	340
Inflow & Storage	0	41	11	45	35	107	85	86	66	112
Reuse & Seepage	62	283	255	426	390	300	196	246	243	293
Recycled Water	12	0	0	0	0	12	12	0	1	1
Total Supplies	9,057	19,557	25,669	20,811	18,587	34,987	17,137	17,539	14,926	20,276

Figure NC-18 Groundwater Level Trends in Selected Wells in the North Coast Hydrologic Region





Figure NC-19 Flood Exposure to the 100-Year Floodplain, North Coast Hydrologic Region



Z:\Projects\102760_StatewideFloodPlan\HydroRegions\100_8x11.mxd

North Coast Key Results			
Total Population:	636,800	Transportation Facilities:	429
Population Exposed:	33,300	Transportation Segments (miles):	330
Percent of Population Exposed:	5	Essential Facilities:	45
Exposed Structures:	17,400	Lifeline Utilities:	10
Value of Exposed Structure and Contents:	\$3.0 Billion	Dept. of Defense Facilities:	-
Total Area (acres)	12.4 Million	Dept. of Defense Facilities (acres):	-
Exposed Area (acres)	400,800	High Potential Loss Facilities:	32
Percent of Area Exposed:	3	Native American Tribes:	4
Exposed Ag. Crops (acres):	108,300	Native American Tribal Lands (acres):	5,568
Percent of Ag. Crops Exposed:	25	Sensitive Animal Species Exposed:	117
Value of Exposed Ag. Crops:	\$84.0 Million	Sensitive Plant Species Exposed:	203

Statewide Flood Hazard Exposure Summary for the North Coast Hydrologic Region 100-year Floodplain

STATEWIDE FLOOD MANAGEMENT PLANNING PROGRAM

Jan 28, 2013

Figure NC-20 Flood Exposure to the 500-Year Floodplain, North Coast Hydrologic Region




Figure NC-21 Location of Groundwater Management Plans in the North Coast Hydrologic Region



Figure NC-22 Groundwater Adjudications in the North Coast Hydrologic Region





Figure NC-23 Integrated Regional Management Planning Areas in the North Coast Hydrologic Region






Figure NC-24 Change in North Coast Agricultural and Urban Water Demands for 117 Scenarios from 2006-2050 (thousand acre feet per year)



Climate

- Historical
- Future

Figure NC-25 Energy Intensity of War Water Extraction and Conveyance in the North Coast Hydrologic Region

Type of Water	Energy Intensity (yellow bulb = 1-500 kWh/AF)	% of regional water supply
Colorado (Project)	<i>This type of water not available</i>	0%
Federal (Project)	 <250 kWh/AF	21%
State (Project)	<i>This type of water not available</i>	0%
Local (Project)	 <250 kWh/AF	27%
Local Imports	<i>This type of water not available</i>	1%
Groundwater	 <250 kWh/AF	28%

Energy intensity per acre foot of water

Energy intensity (EI) in this figure is the total amount of energy required for the extraction and conveyance of one acre-foot of water and does not include treatment, distribution to point of use, or end use energy (e.g., water heating). These figures should be seen as ranges within which the EI of different sources of each water type would likely fall i.e., a water type with four bulbs should be interpreted to mean that most sources of that water type in the region would have an EI of between 1,501-2,000 kWh/ acre-ft of water. Smaller light bulbs represent an EI of greater than zero, and less than 250 kWh/acre-ft. EI of desalinated and recycled water is not shown, but is covered in Resource Management Strategies #XX and #YY respectively, Volume 3. (For detailed description of the methodology used to calculate EI in this figure, see Technical Guide, Volume 5 or References Guide, Volume 4 (TBD)).

1 **Box NC-1 California Statewide Groundwater Elevation Monitoring (CASGEM) Basin Prioritization**
2 **Data Considerations**

3 Senate Bill 7x 6 (SBx7 6; Part 2.11 to Division 6 of the California Water Code § 10920 et seq.) requires, as part of the
4 CASGEM program, DWR to prioritize groundwater basins to help identify, evaluate, and determine the need for additional
5 groundwater level monitoring by considering available data listed below:.

- 6 1. The population overlying the basin,
- 7 2. The rate of current and projected growth of the population overlying the basin,
- 8 3. The number of public supply wells that draw from the basin,
- 9 4. The total number of wells that draw from the basin,
- 10 5. The irrigated acreage overlying the basin,
- 11 6. The degree to which persons overlying the basin rely on groundwater as their primary source of water,
- 12 7. Any documented impacts on the groundwater within the basin, including overdraft, subsidence, saline intrusion, and
13 other water quality degradation, and
- 14 8. Any other information determined to be relevant by the DWR.

15 Using groundwater reliance as the leading indicator of basin priority, DWR evaluated California's 515 alluvial groundwater
16 basins and categorized them into five groups:

- 17 • Very High
- 18 • High
- 19 • Medium
- 20 • Low
- 21 • Very Low

1 **Box NC-2 Near-Coastal Issues**

2 Coastal regions in California share common concerns and issues. The update of the California Water Plan 2013 is
3 introducing a focus on near-coastal issues. The issues common to all coastal areas include increased coastal flooding
4 especially as it relates to climate change, sea level rise, and the potential degradation of aquifer water quality. Desalination
5 may be a future water supply source for drinking water, and impacts on adjacent water conditions and ecosystems are of
6 concern. Stormwater and wastewater management are significant near-coastal issues, including the impacts of runoff and
7 discharge on coastal water quality. Near coastal planners and resource managers have increased attention to ecological
8 linkages between freshwater flows, wetlands, and anadromous fish species. Conjunctive water management strategies as
9 applied in near coastal areas consider groundwater management for recharge and water supply for multiple land uses and
10 objectives.

11 Climate change is anticipated to have profound effects on the North Coast regions, as the effects of climate change will alter
12 rain patterns and intensity and well as temperatures. Because of the interrelationship of water supply, quality, floods and
13 flooding, land use and fisheries, coastal managers are relying on current science and recommended strategies for
14 adaptation and resource management. These shared concerns, issues, approaches and strategies are discussed below in
15 detail relevant to the North Coast region.

16 Find information on near-coastal issues in the North Coast region under the "Flood Management" and "Climate Change ... "
17 sections as well as "Recent Initiatives ..." and "Ecosystem Restoration." In Volume 4, Near-Coastal Issues are discussed in
18 an article "[title to be provided]."
19

1 **Box NC-3 Tribal Water Rights on the Klamath and Trinity Rivers**

2 Tribal water rights on the Klamath and Trinity rivers have had a long and complicated history. This history as it relates to
3 Tribal Water Rights has been well documented in a briefing paper presented at the 2009 California Tribal Water Summit by
4 the Hoopa Valley Tribe(cite below). This paper describes the legal history of the Klamath and Trinity rivers with details on
5 the impacts of the Trinity River Diversion for the Central Valley Project and issues of sustainable water quantity and quality
6 in the Klamath River. It further describes the provisions of the Trinity River Restoration Program which was enacted under
7 the Central Valley Project Improvement Act.

8 The paper also details three potential adverse effects of the proposed Klamath River Restoration Agreement (KRRRA) and
9 the Klamath Hydroelectric Settlement Agreement (KHSA) on the success of the Trinity River Restoration Program. The three
10 potential adverse effects are the cost of implementing the KBRA of over \$1 billion, the KBRA's guaranteed irrigation
11 diversions of water for the Klamath Irrigation District Project in Oregon on water availability, and the lengthy dam removal
12 planning process authorized by the KHSA which will delay any restoration for many years.

13 The conclusion made by the briefing paper is that Native American tribes have a key role in the sustainable use of water
14 both in terms of quantity and quality. Tribes must be accorded the respect due to a government and dealt with on a
15 government-to-government basis if successful accommodation of the competing interests is to be achieved.

16 _____
17 Add to citation in NC RR:

18 Hoopa Valley Indian Tribe, California Tribal Water Summit 2009, Tribal Water Authorities-Rivers, Dams & Fish, Section 9.
19 Hoopa Valley Tribe, P.O. Box 1348 Hoopa, CA 95546, Thomas P. Schlosser, October 21, 2009, PDF. Viewed online at:
20 http://www.waterplan.water.ca.gov/docs/tws/CTWS_BriefingPaper_Rivers2_Hoopa_v1.pdf. Accessed on: Feb 6, 2013. Last
21 update: Oct 2009.

Box NC-4 Other Groundwater Management Planning Efforts in the North Coast Hydrologic Region

The Integrated Regional Water Management plans, Urban Water Management plans, and Agriculture Water Management plans in the North Coast Hydrologic Region that also include components related to groundwater management are briefly discussed below.

Integrated Regional Water Management Plans

The North Coast Hydrologic Region is unique in that it is fully covered by one IRWM plan. Although the North Coast IRWM plan addresses groundwater resources in their goals and objectives, similar to other IRWM plans throughout the State, they do not actively manage local groundwater resources. Instead they defer groundwater management to local entities with groundwater management plans, and identify county, state, federal, and tribal entities that address groundwater management issues, such as County General Plans, the California Water Plan, the Environmental Protection Agency's Underground Injection Control Program, and Tribal/Reservation plans. Regional prioritization of groundwater management plan development and implementation of local groundwater management planning is one of the goals of this IRWM region.

Urban Water Management Plans

Urban Water Management plans are prepared by California's urban water suppliers to support their long-term resource planning and to ensure adequate water supplies are available to meet existing and future water uses. Urban use of groundwater is one of the few uses that meter and report annual groundwater extraction volumes. The groundwater extraction data is currently submitted with the Urban Water Management plan and then manually translated by DWR staff into a database. Online methods for urban water managers to directly enter their water use along with their plan updates is currently under evaluation and review by DWR. Because of the time-line, the plans could not be reviewed for assessment for Water Plan Update 2013.

Agricultural Water Management Plans

Agricultural Water Management plans are developed by water and irrigation districts to advance the efficiency of farm water management while benefitting the environment. New and updated Agricultural Water Management plans addressing several new requirements were submitted to DWR by December 31, 2012 for review and approval. These new or updated plans provide another avenue for local groundwater management, but because of the time-line, the plans could not be reviewed for assessment for Water Plan Update 2013.

Box NC-5 Statewide Conjunctive Management Inventory Effort in California

The effort to inventory and assess conjunctive management projects in California was conducted through literature research, personal communication, and documented summary of the conjunctive management projects. The information obtained was validated through a joint DWR-ACWA survey. The survey requested the following conjunctive use program information:

1. Location of conjunctive use project;
2. Year project was developed;
3. Capital cost to develop the project;
4. Annual operating cost of the project;
5. Administrator/operator of the project; and
6. Capacity of the project in units of acre-feet.

To build on the DWR/ACWA survey, DWR staff contacted by telephone and email the entities identified to gather the following additional information:

7. Source of water received;
8. Put and take capacity of the groundwater bank or conjunctive use project;
9. Type of groundwater bank or conjunctive use project;
10. Program goals and objectives; and
11. Constraints on development of conjunctive management or groundwater banking (recharge) program.

Statewide, a total of 89 conjunctive management and groundwater recharge programs were identified. Conjunctive management and groundwater recharge programs that are in the planning and feasibility stage are not included in the inventory.