

29.1 Introduction

Climate is the average weather over many years, measured most often in terms of temperature, precipitation, and wind. For example, the climate of California’s Central Valley is a Mediterranean climate, which is hot and dry during the summer and cool and damp in winter, with the majority of precipitation falling as rain in the winter months. Climate is unique to a particular location and changes on timescales of decades to centuries or millennia.

Climate change generally refers to a “statistically significant variation in either the mean state of the climate or in its variability, persisting for an extended period (typically decades or longer)” (World Meteorological Organization 2013). Although the climate can change, and has changed, in the past in response to natural drivers, recent climate change has been more rapid than previous episodes of climate change and has been unequivocally linked to increasing concentrations of greenhouse gases (GHGs) in Earth’s lower atmosphere and the rapid timescale on which these gases have accumulated (Intergovernmental Panel on Climate Change 2007a). The major causes of this rapid loading of GHGs into the atmosphere include the burning of fossil fuels since the beginning of the industrial revolution, agricultural practices, increases in livestock grazing, and deforestation. More background information on GHG emissions is provided in Chapter 22, *Air Quality and Greenhouse Gases*, Section 22.1.3.

Higher concentrations of heat-trapping GHGs in the atmosphere result in increasing global surface temperatures, a phenomenon commonly referred to as *global warming* or *climate change*. Higher global surface temperatures in turn result in changes to the Earth’s climate system, including: the jet stream; El Niño; the Indian monsoon; ocean temperature and acidity; the extent of alpine glaciers, sea ice and polar ice sheets; atmospheric water content; and the extent and health of boreal and tropical forests (Intergovernmental Panel on Climate Change 2007a, b). Some of the above changes will result in specific impacts at the state and local level.

29.2 Purpose

This EIR/EIS analyzes three fundamental questions relating to climate change. Two of them are analyzed in other chapters. The third is analyzed in this chapter.

1. What is the impact of the BDCP alternatives on climate change? i.e., how will GHG emissions from construction and operation activities associated with the project alternatives contribute to elevated GHG concentrations in the atmosphere?
2. How will the impacts of the BDCP alternatives on the study area for each resource (the area in which impacts may occur) be affected by climate change? i.e., are future changes in climate likely to exacerbate project impacts?
3. How will the BDCP alternatives affect the resiliency and adaptability of the Plan Area (the area covered by the BDCP) to the effects of climate change?

1 Question 1 is addressed in Chapter 22, *Air Quality and Greenhouse Gases* (Impacts AQ-15, AQ-16, and
2 AQ-18), through the calculation of GHG emissions inventories and identification of GHG mitigation
3 opportunities associated with the BDCP alternatives.

4 Question 2 is addressed throughout this document in each of the resource chapters. Under
5 discussion of the No Action Alternative, each resource chapter evaluates how the BDCP alternatives
6 would affect the specific resource in question. In each of these analyses, where the effects of the
7 BDCP alternatives are analyzed for 2025 and 2060 conditions, climate change is integrated into the
8 analysis. In these analyses, the BDCP alternatives are evaluated using a projection of future climate
9 that includes changes in temperature, precipitation, humidity, hydrology, and sea level rise (SLR).
10 Appendix 5A, *BDCP EIR/EIS Modeling Technical Appendix*, provides detailed information about the
11 development of the climate change projections. The interrelation between resource topics
12 addressed in this EIS/EIR and potential climate change effects under the No Action Alternative are
13 presented in Table 29-1. An 'X' in the table signifies that there is a clear connection between the
14 resource topic and a climate change effect under the No Action Alternative. Readers seeking
15 additional information about a specific climate change effect on a specific resource should reference
16 the resource specific chapter of this EIR/EIS. The potential climate effects under the No Action
17 Alternative listed in Table 29-1 are based on the California Natural Resource Agency's (CNRA)
18 climate adaptation guidance (California Natural Resources Agency 2009) that was adapted to be
19 specific to the Plan Area.

20 Question 2 also fulfills the requirements for climate change analysis outlined in the Delta Reform Act
21 of 2009 (Cal. Water Code, § 85000 *et seq.*). Within the Delta Reform Act, Water Code Section 85320
22 identifies the contents that the EIR portion of this Draft EIR/EIS must include for the BDCP to be
23 considered for inclusion in the Delta Plan prepared by the Delta Stewardship Council. Section
24 85320(b)(2)(C) of the Water Code directs that the EIR address "[t]he potential effects of climate
25 change, *possible sea level rise up to 55 inches* [140 centimeters], and possible changes in total
26 precipitation and runoff patterns on the conveyance alternatives and habitat restoration activities
27 considered in the [EIR]." (Italics added.). It should be noted, the California Ocean Protection Council
28 and other scientific bodies have projected that SLR will not reach 55 inches (140 centimeters) until
29 approximately the year 2100. SLR projections for 2025 and 2060 were developed based on research
30 available during the analysis design and based on the requirements of Water Code Section 85320,
31 which required that BDCP evaluate a sea level rise of 55 inches (well in excess of the expected sea
32 level described by any major study for 2060).

33 This information is provided to discuss the benefits of the BDCP alternatives in the face of expected
34 climate change.

35 This chapter is organized differently from the other resource chapters because analyzing how the
36 BDCP alternatives would improve the Plan Area's resiliency and adaptability to climate change is a
37 fundamentally different analysis than those presented in other resource chapters. Whereas the
38 other chapters are organized to identify effects of the action alternatives and how to mitigate these
39 impacts, this chapter's function is to analyze and disclose how the action alternatives affect the Plan
40 Area's resiliency and adaptability to expected climate change. The study area for this chapter,
41 therefore, is defined as the Plan Area, which is largely formed by the statutory borders of the Delta,
42 along with areas in Suisun Marsh and the Yolo Bypass.

43 This chapter addresses question 3: How will the BDCP alternatives affect the resiliency and
44 adaptability of the Plan Area to the effects of climate change? In this context, resiliency and

1 adaptability mean the ability of the Plan Area to remain stable or flexibly change, as the effects of
 2 climate change increase, in order to continue providing water supply benefits with sufficient water
 3 quality and supporting ecosystem conditions that maintain or enhance aquatic and terrestrial plant
 4 and animal species.

5 **Table 29-1. Linkages between Resource Areas Addressed in this EIR/EIS and Climate Change**

Resource Topic	Potential Climate Change Effect																	
	Increased air temperature	Increased water temperature	Increased soil temperature	Reduced precipitation/ runoff volumes	Shift from snowfall to rainfall	Early snowmelt	Changes in evapotranspiration	Increased frequency/severity of flood events	Increased frequency/severity of droughts	Increased frequency of extreme heat events	SLR	Increased salinity intrusion	Changes in erosion/ sedimentation rates	Decreased species populations and quality of species habitat	Changes in species geographic range/distribution	Spreading of pests and vector-borne diseases	Increased fire risk	Increased atmospheric CO ₂ concentrations and acidification
Water Supply (Ch. 5)	X	X	X	X	X	X	X	X	X	X	X	X	X				X	
Surface Water (Ch. 6)	X	X	X	X	X	X	X	X	X		X	X	X					
Groundwater (Ch. 7)	X	X	X	X	X	X	X	X	X		X	X						
Water Quality (Ch. 8)	X	X		X	X	X	X	X	X		X	X	X	X		X		X
Geology and Seismicity (Ch. 9)																		
Soils (Ch. 10)	X		X					X	X		X	X	X					X
Fish and Aquatic Resources (Ch. 11)	X	X		X	X	X		X	X	X	X	X	X	X	X		X	X
Terrestrial Biological Resources (Ch. 12)	X	X		X	X	X	X	X	X	X	X	X	X	X	X	X	X	X
Land Use (Ch. 13)	X		X				X	X			X						X	
Agricultural Resources (Ch. 14)	X		X	X			X	X	X	X	X	X					X	X
Recreation (Ch. 15)	X			X		X		X	X	X	X				X		X	
Socioeconomics (Ch. 16)	X		X	X				X	X	X	X			X		X	X	
Aesthetics and Visual Resources (Ch. 17)	X							X	X	X					X		X	
Cultural and Historic Resources (Ch. 18)			X					X	X		X		X				X	X
Transportation (Ch. 19)								X			X							
Public Services and Utilities (Ch. 20)	X							X	X	X	X					X	X	
Energy (Ch. 21)	X	X		X	X			X	X	X								
Air Quality and Greenhouse Gases (Ch. 22)	X	X						X		X	X						X	X
Noise (Ch. 23)																		
Hazards and Hazardous Materials (Ch. 24)	X							X		X	X						X	
Public Health (Ch. 25)	X							X	X	X	X	X				X	X	
Mineral Resources (Ch. 26)	X							X			X		X				X	X
Paleontological Resources (Ch. 27)	X							X	X				X				X	X
Environmental Justice (Ch. 28)	X							X	X	X	X					X	X	

6
 7 This resiliency and adaptation analysis focuses on the major impacts of climate change in the Plan
 8 Area and the clear and measurable ways that the BDCP alternatives will ameliorate these impacts or

1 add flexibility to the system so that the Plan Area can continue providing water supply benefits with
2 sufficient water quality and supporting ecosystem conditions that maintain or enhance aquatic and
3 terrestrial plant and animal species. No single project and indeed none of the BDCP alternatives
4 would be able to completely counteract all of the impacts of climate change; however, as shown
5 below the BDCP alternatives provide important added resilience and adaptability to many of the
6 expected changes. Impacts for which the BDCP alternatives provide no added resiliency or
7 adaptation benefit or for which the benefit is minimal, or not documentable are not discussed in this
8 chapter.

9 While there is a lot of overlap between the analysis provided here and that provided in the resource
10 effects chapters, the main difference is that this chapter focuses on both negative effects and benefits
11 and that it compares a climate changed future without the BDCP alternatives to a climate changed
12 future with the BDCP alternatives. Resource chapters include comparisons to the No Action/No
13 Project Alternative at 2060 (the NEPA point of comparison), which represents the net impact of the
14 project isolated from the effects of climate change. These chapters also compare the BDCP
15 alternatives to Existing Conditions (the CEQA baseline), which represents the net impact of the
16 project combined with the effects of climate change. The differences between these two
17 comparisons allow readers to determine the incremental effects attributable to climate change as
18 distinct from the impacts of the action alternatives. The resource chapters do not, however,
19 specifically contemplate the extent to which BDCP action alternatives would contribute to the
20 resiliency and adaptability of the Plan Area to the effects of climate change. Instead, this analysis is
21 included in this chapter.

22 29.3 Organization

23 This chapter presents (1) basic background on scientific efforts to evaluate the degree and impacts
24 of future climate changes (a detailed background discussion on climate change is provided in
25 Appendix 5A, *BDCP EIR/EIS Modeling Technical Appendix*); (2) a discussion of observed
26 climatological changes over the past several decades, and expected future changes during the rest of
27 this century globally, in California, and for the Plan Area; (3) an evaluation of the resiliency and
28 adaptability of the Plan Area to the major expected impacts of climate change; and (4) an evaluation
29 of the BDCP alternatives' compatibility with applicable plans and policies designed to adapt to
30 climate change or improve resilience to it.

31 29.4 Climate Change Background

32 A vast amount of scientific research on climate change, both its causes and effects, at all geographic
33 scales has been conducted during the last 50 years. The Intergovernmental Panel on Climate Change
34 (IPCC) was established by the United Nations Environment Program and the World Meteorological
35 Organization (WMO) to provide the world with a clear scientific view of the current state of
36 knowledge regarding climate change and its potential environmental and socioeconomic impacts
37 (Intergovernmental Panel on Climate Change 2011). IPCC, an organization of more than 800
38 scientists from around the world, regularly publishes summary documents, which analyze and
39 consolidate all recent peer-reviewed scientific literature, providing a consensus of the state of the
40 science. Thus, IPCC is viewed by governments, policymakers and scientists as the leading
41 international body on the science of climate change and its summaries are considered to be the best

1 available science. IPCC documents address change at the global and super-regional scales. Both IPCC
2 studies and California-specific studies (e.g., California Air Resources Board [CARB], California
3 Energy Commission [CEC], the California Department of Water Resources [DWR], CNRA, and U.S.
4 Department of the Interior, Bureau of Reclamation) that are based on IPCC data are referenced
5 throughout this chapter.

6 Scientific measurements have shown that changes in the *global* climate system are already
7 occurring. These include: rising air temperatures; rising ocean temperatures; rising ocean salinity;
8 rising global sea levels; changes in precipitation patterns; and increased intensity and frequency of
9 extreme events such as storms, droughts, and wildfires (Intergovernmental Panel on Climate Change
10 2007b; California Department of Water Resources 2009).

11 29.5 Environmental Setting/Affected Environment

12 The Plan Area has a predominantly Mediterranean climate characterized by hot, dry summers and
13 cool, rainy winters. From 1981–2010, average monthly temperatures in Sacramento ranged from
14 41.0°F (5°C) in December and January to 94.1°F (34.5°C) in July, with average monthly rainfall
15 ranging from a low of 0.02 inches (0.05 centimeters) in July to a high of 3.90 inches (9.9
16 centimeters) in February (Western Regional Climate Center 2012). Average air temperatures in the
17 mountainous regions of the watershed are typically 5–10 degrees lower than the temperature on
18 the valley floor.

19 Although the snow lines vary by storm event, portions of the Sacramento, San Joaquin, Mokelumne,
20 and Cosumnes River watersheds are above the snow line; consequently, much of their respective
21 runoff into the Delta is from snowmelt. Snow in higher elevations serves as an effective type of
22 natural storage because it typically melts gradually during the spring and summer. The snowline is
23 often around the elevation of 5,000 feet (1,524 meters) (U.S. Army Corp of Engineers 2002).

24 Annual precipitation in the Sacramento River watershed ranges from 80 to 90 inches (as liquid
25 water) (203 to 229 centimeters) of primarily snowfall in the mountainous regions, to 41 inches (104
26 centimeters) of rain in Redding and 19 inches (48 centimeters) in Sacramento. Average annual
27 precipitation for the entire watershed is approximately 36 inches (91 centimeters). Most
28 precipitation occurs between November and April, with little or no precipitation falling between
29 May and October (Huber-Lee et al. 2003). Precipitation that falls as rain in the project area can run
30 off into the rivers (and eventually into the Delta), infiltrate into the soils (recharging the
31 groundwater system) or evapotranspire. Factors such as spring temperatures and the nature of
32 precipitation (rain/snow elevations in storms) during the October to April period play an important
33 role in runoff timing.

34 The primary type of soil in the Plan Area is peat. These soils were developed by the formation of
35 mineral soils near the channels during flood conditions, and by the formation of organic soils on
36 marsh island interiors as plant residues accumulated faster than they could decompose. Prior to the
37 mid-1800s, the Delta was a vast marsh and floodplain, under which peat soils developed to a
38 thickness of up to 30 feet (9 meters) in some areas. In addition to peat, the Delta soils are composed
39 of mineral sediments from rivers (United States Geological Survey 2000).

40 The Plan Area has historically been affected by periodic extreme precipitation events. The majority
41 of these historical events have likely been caused by an atmospheric phenomenon called an
42 atmospheric river (AR) (Dettinger 2011). ARs are narrow corridors of water vapor transported in

1 the lower atmosphere that traverse long swaths of the Earth’s surface (Dettinger and Ralph 2011).
 2 These storms can deliver tremendous amounts of precipitation to California in a very short period of
 3 time. In addition, these storms tend to be warm (originating in the tropics) which results in higher
 4 snowlines and larger portions of the watershed contributing to direct runoff. More detailed
 5 information on surface water and climate and meteorological conditions in the Plan Area is provided
 6 in Chapter 6, *Surface Water*, and Chapter 22, *Air Quality and Greenhouse Gases*.

7 Because this chapter discusses how the BDCP alternatives affect the resiliency and adaptability of
 8 the Plan Area to the effects of climate change, this section also discusses expected changes to the
 9 environmental setting. The following background sections provide brief descriptions of (1) recent
 10 trends in key climate metrics such as temperature, precipitation, and sea level, and (2) projections of
 11 how the climate will change between now and 2100. Although the year 2100 is approximately 40
 12 years after the end of the 2060 time period analyzed in other chapters of this Draft EIR/EIS
 13 (reflecting the approximate end date of the 50-year permit term proposed for the BDCP), the year
 14 2100 was chosen in part because of language enacted by the California Legislature in the
 15 Sacramento-San Joaquin Delta Reform Act of 2009 (Cal. Wat. Code, § 85000 *et seq.*) requiring the EIR
 16 to address “[t]he potential effects of climate change, *possible sea level rise up to 55 inches* [140
 17 centimeters], and possible changes in total precipitation and runoff patterns on the conveyance
 18 alternatives and habitat restoration activities considered in the [EIR]” (Cal. Wat. Code, § 85320.
 19 *Italics added.*). It should be noted, the California Ocean Protection Council and other scientific bodies
 20 have projected that SLR will not reach 55 inches (140 centimeters) until approximately the year
 21 2100.

22 This information is provided at the global scale, at the state level, and for the Plan Area. Projections
 23 of future climate change are based on the level of GHGs already in the atmosphere, the current rate
 24 at which human activity releases GHGs to the atmosphere, and the future rate of GHG emissions,
 25 which in turn relies on predictions of future population, global economic growth, future available
 26 energy sources, and regulations. Consequently, future projections of climate are typically displayed
 27 as a range, with the lower end representing a minimum amount of estimated change based on past
 28 and current GHG emissions and the higher end representing a high degree of global economic
 29 growth and the absence of large-scale mitigation of GHG emissions.

30 **29.5.1.1 Global Climate Change Effects**

31 **Recent Trends**

32 The IPCC has found that, “[w]arming of the climate system is unequivocal, as is now evident from
 33 observations of increases in global average air and ocean temperatures, widespread melting of snow
 34 and ice, and rising global average sea level” (Intergovernmental Panel on Climate Change 2007a).
 35 Global annual surface temperatures have increased at a rate of 0.13°C (0.23°F) per decade during
 36 the period 1950–2000. This rate is double the rate observed during the period 1900–1950. Further,
 37 11 of the 12 years during the period 1995–2006 rank among the 12 warmest years in the
 38 instrumental record of global surface temperature (since 1850) (Intergovernmental Panel on
 39 Climate Change 2007a).

40 Much of the Western United States has experienced warming during the 20th century (roughly 2°F
 41 [1.1°C]) and is projected to experience further warming during the 21st century with central
 42 estimates varying from roughly 5 to 7°F (2.8°C to 3.8°C), depending on location (Bureau of
 43 Reclamation 2011). Historical trends in annual precipitation are less apparent. Future projections

1 suggest that the Northwestern and north-central portions of the United States gradually may
 2 become wetter (e.g., Columbia Basin and Missouri River basin) while the Southwestern and south-
 3 central portions gradually become drier (e.g., San Joaquin, Truckee, and Rio Grande River basins and
 4 the Middle to Lower Colorado River Basin). Areas in between have median projected changes closer
 5 to no change, meaning they have roughly equal chances of becoming wetter or drier (e.g., Klamath
 6 and Sacramento basins and the Upper Colorado Basin). These summary statements refer to median
 7 projected changes in temperature and precipitation, characterized generally across the Western
 8 United States. Projections show that there is significant variability and uncertainty about these
 9 projected conditions both geographically and with time (Bureau of Reclamation 2011).

10 Warming trends appear to have led to a shift in cool season precipitation towards more rain and less
 11 snow, which has caused increased rainfall-runoff volume during the cool season accompanied by
 12 less snowpack accumulation in some Western United States locations (Bureau of Reclamation 2011).
 13 Hydrologic analyses-based future climate projections suggest that warming and associated loss of
 14 snowpack will persist over much of the Western United States. However, there are some geographic
 15 contrasts. Snowpack losses are projected to be greatest where the baseline climate is closer to
 16 freezing thresholds (e.g., lower lying valley areas and lower altitude mountain ranges). It also
 17 appears that, in high altitude and high latitude areas, there is a chance that cool season snowpack
 18 actually could increase during the 21st century (e.g., Columbia headwaters in Canada, Colorado
 19 headwaters in Wyoming), because precipitation increases are projected and appear to offset the
 20 snow-reduction effects of warming in these locations (Bureau of Reclamation 2011).

21 During the same period over which global temperatures have increased, sea levels have risen on
 22 average 0.07 inches (0.18 centimeters) per year with SLR during the period 1993–2003 rising at a
 23 rate of 0.12 inches (0.31 centimeters) per year and increasing overall by about 6.7 inches (17
 24 centimeters) during the twentieth century (Intergovernmental Panel on Climate Change 2007a).
 25 Observed trends in SLR can be attributed to both thermal expansion of the world’s oceans and the
 26 melting of ice sheets (polar and alpine). Also during a similar period (1900–2007) measurements
 27 have shown increases in global ocean temperature (since 1961); a decline in the extent of mountain
 28 glaciers and global snow cover; increased atmospheric water vapor content; loss in mass of the
 29 polar ice sheets; decreased extent of Arctic sea ice; increased precipitation in the eastern portions of
 30 North and South America, northern Europe and northern and central Asia; drying conditions in the
 31 Sahel region of the Sahara Desert in Africa, the Mediterranean and southern Africa; strengthening in
 32 mid-latitude westerly winds (since 1960s); more intense and longer drought conditions in the
 33 tropics and sub-tropics (since the 1970s); increased frequency of extreme precipitation events over
 34 land areas; higher average night time temperatures; decreased frost days and increased frequency
 35 and duration of extreme heat events (since 1950s); and increased tropical cyclone activity in the
 36 North Atlantic (Intergovernmental Panel on Climate Change 2007a). There may also be additional
 37 synergistic impacts of extreme weather events, such as the SLR coupled with high tide and extreme
 38 storm surges. The above listed changes are in turn resulting in changes to the climate of California as
 39 the regional climate is moderated by sea surface temperature, westerly jet stream wind patterns, the
 40 El Niño Southern Oscillation (ENSO), and Pacific storm patterns.

41 **Projections to 2100**

42 Climate models indicate that global average surface temperature will increase at a rate of
 43 approximately 0.4°F (0.2°C) per decade for the period 2000–2020, and will increase by at least that
 44 amount per decade during the period 2020–2080. Based on a number of emissions scenarios, the
 45 IPCC projected an average increase in surface temperatures of 3.2 to 7.2°F (1.7 to 4°C) by 2100

1 compared to 1980 through 1999 levels, with a likely range of 2.0 to 11.5°F (1.1 to 2.2°C) when
 2 accounting for the uncertainty in climate science (Intergovernmental Panel on Climate Change
 3 2007a). Approximately half of this warming is the result of past GHG emissions and will occur even
 4 if GHG emissions were halted at 2000 levels. Some regions of the globe, particularly high latitudes,
 5 will experience much larger changes relative to Existing Conditions. Corresponding global average
 6 SLR levels during the period 2000–2100 are estimated to be between 7 inches (18 centimeters) and
 7 23 inches (58 centimeters) (Intergovernmental Panel on Climate Change 2007a). However, recent
 8 scientific data now strongly suggests that these SLR projections are likely too low and that actual
 9 SLR may be significantly greater than initially estimated (Rahmstorf 2007; National Research
 10 Council 2012).¹

11 The following additional changes to the global climate system are projected: increased ocean acidity
 12 due to increased carbon dioxide uptake by the oceans; reduced global snow cover; increased thaw
 13 depth in permafrost regions; decrease in sea ice with potential full disappearance in summer
 14 months; increased frequency in heat waves, droughts, and heavy precipitation events; increased
 15 intensity of tropical cyclone events; northward movement of extra-tropical storm tracks; increased
 16 precipitation at high latitudes and decreased precipitation in tropical and sub-tropical regions; and
 17 increased melting of the ice sheets (Intergovernmental Panel on Climate Change 2007a).

18 **29.5.1.2 Climate Change Effects on California**

19 **Recent Trends**

20 Scientific measurements and observations indicate that California’s climate is already changing in a
 21 manner consistent with what would be expected from global climate change. Since 1920, California’s
 22 average temperature has been increasing, although this change, or any climate change impact, is not
 23 uniform across California. Nighttime temperatures are rising across California and at a higher rate
 24 than day-time temperatures. Furthermore, daytime and nighttime heat wave events throughout
 25 California have increased in intensity, particularly the nighttime component (Moser et al. 2009).
 26 During the last century, sea level along the California coast has increased approximately 7 inches (18
 27 centimeters), with higher rates of increase occurring since 1993 (Cayan et al. 2009).

28 California’s water supply system is dependent on snowpack storage in the Sierra Nevada.
 29 Temperatures over the Sierra Nevada have increased during the last 100 years, resulting in less
 30 snowfall (and more rainfall) and an earlier snowmelt (Moser et al. 2009). The average early spring
 31 snowpack in the Sierra Nevada has decreased by about 10 percent during the last century, a loss of
 32 1.5 million acre-feet (MAF) of snowpack storage (California Department of Water Resources 2008).
 33 Reductions in water supply can adversely affect hydropower reserves, decreasing hydropower
 34 generation in the summer months when peak demand is highest (California Natural Resources
 35 Agency 2012).

¹ California agencies including the Bay Conservation and Development Commission (BCDC) and DWR are using the more recent data of Rahmstorf et al. 2007 in their SLR planning efforts in lieu of the estimates as reported by IPCC in the Fourth Assessment Report. As identified above, California Water Code Section 85320 identifies in order to be considered for inclusion in the Delta Plan, the BDCP must assess “[t]he potential effects of climate change, *possible sea level rise up to 55 inches* [140 centimeters], and possible changes in total precipitation and runoff patterns on the conveyance alternatives and habitat restoration activities.”

1 Data also show evidence for the following additional changes to California climate and conditions
 2 during the last 50 years: the warming of Lake Tahoe; decreasing chill hours and increased stresses
 3 on California agriculture; shifts and disturbances in managed landscapes; increased frequency of
 4 wildfire; changes in Santa Ana winds; increases in photochemical smog production in southern
 5 California; increased frequency and intensity of heat wave and drought events; changes in ENSO and
 6 the impact on California temperatures; and changes in extreme precipitation events and daily
 7 average precipitation (California Energy Commission 2011a).²

8 Plants and animals around the globe are already reacting to changes caused by increasing
 9 temperatures. In California, species are also reacting to extreme conditions, including heat waves
 10 (and increased fire frequency); cold snaps; droughts (and the saltwater intrusion that droughts
 11 often cause); floods; and coastal upwelling. Observed changes also include altered timing of animal
 12 and plant lifecycles (phenology), disruption of biotic interactions, changes in physiological
 13 performance, species range and abundance, increase in invasive species, altered migration patterns
 14 of fishes, aquatic-breeding amphibians, birds and mammals, changes in forage base, local extinction
 15 of plant and animal populations, and changes in habitat, vegetation structure, and plant and animal
 16 communities (California Department of Fish and Game 2010).

17 **Projections to 2100**

18 Average annual surface temperatures for California are projected to increase by between 2 and 5°F
 19 (1.1 and 2.8°C) by 2050 and between 4 and 9°F (2.2 and 5.0°C) by 2100, depending on the GHG
 20 emissions scenario assumed. Warming will not be uniform temporally or geographically across the
 21 state. Climate models project a greater amount of warming during summer months, especially
 22 during nighttime, and in the interior regions of California. Chill hours in the Central Valley are
 23 expected to decrease, but unprecedented extremes of cold weather are still possible (Gershunov
 24 2011). Changes in temperature and humidity have implications for agriculture in the Central Valley;
 25 as the climate warms and dries, crop diversity and production may slow (Jackson et al. 2011).
 26 Extreme events will also stress California's energy system (Auffhammer 2011).

27 Best available data indicate that California, as a whole, will experience changes in precipitation. It is
 28 likely that some areas in California will experience higher annual rainfall amounts whereas
 29 precipitation in other regions will decrease (Gershunov 2011). Cayan et al. (2009) estimates
 30 California, particularly southern California, will be 15–35% drier by 2100. Snowpack volumes are
 31 expected to diminish by 25% by 2050 (California Department of Water Resources 2010b).

32 Frequency and intensity of large storms and precipitation events may be influenced by changes in
 33 ARs. In California, nearly all major historic flood events have been associated with the presence of
 34 ARs along the Pacific coast. It is estimated that future changes in climate will increase the frequency
 35 of years with AR storms, but the number of storms per year is not likely to be affected. More
 36 importantly, occasional “much-larger-than-historical-range storm intensities” are projected to occur
 37 under most warming scenarios. Changes in the frequency and magnitude of ARs may result in
 38 increases in major flood and storm events (Dettinger 2011).

² The State of California under the auspices of the California Energy Commission (CEC) is conducting comprehensive and detailed research into a range of climate change impacts in California as well as research aimed at developing adaptation strategies to deal with impacts already underway and that can no longer be avoided. The majority of this research is available through the California Climate Change Portal. Available at: <<http://www.climatechange.ca.gov/>>.

1 Climate change and increasing temperatures are expected to increase energy demand in California,
 2 particularly during the summer months. The California Natural Resources Agency (2012) predicts
 3 that higher temperatures in the next decade could increase demand by up to 1 gigawatt. Increased
 4 energy demand would require additional generation resources or the purchase of costly peak power
 5 from external sources.

6 SLR along the California Coast is expected to accelerate during the 21st century. A recent study
 7 completed by the National Research Council (NRC) looked at both global (e.g., thermal expansion,
 8 land ice melting) and local (e.g., tectonic land movement, localized subsidence) factors effecting sea
 9 level relative to land surface. Table 29-2 below shows the projection and the range of uncertainty for
 10 expected sea level rise at San Francisco and the Delta at 2030, 2050, and 2100.

11 **Table 29-2. Sea Level Rise Projections and Ranges for San Francisco, California 2030, 2050, and 2100**

	2030		2050		2100		
	Projection	Range	Projection	Range	Projection	Range	
Projected Sea Level Rise at San Francisco	cm	14.4 ± 5.0	4.3–29.7	28.0 ± 9.2	12.3–60.8	91.9 ± 25.5	42.4–166.4
	in	5.7±2	1.7–11.7	11±3.6	4.84–23.9	36.2±10	16.7–65.5

Source: National Research Council 2012

12
 13 SLR will continue to threaten coastal lands and infrastructure, increase flooding at the mouths of
 14 rivers, place additional stress on levees in the Delta, and will intensify the difficulty of managing the
 15 Delta as the heart of the state's water supply system (California Department of Water Resources
 16 2010a). The effects of SLR, combined with large waves generated during El Niño events will have the
 17 greatest potential for impacts (Griggs 2011).

18 These changes in temperature, precipitation and sea level may have substantial effects on other
 19 resources areas. Potential effects of climate change anticipated in California (and discussed in this
 20 document) are listed below (California Natural Resources Agency 2009, 2012).

- 21 • Increased average temperatures (air, water, and soil).
- 22 • Changes in annual precipitation amounts.
- 23 • Change from snowfall (and spring snowmelt) to rainfall.
- 24 • Decreased Sierra snowpack (earlier runoff, reduced maximum storage).
- 25 • Changes in evapotranspiration.
- 26 • Increased frequency and intensity of Pacific storms (flood events).
- 27 • Increased severity of droughts.
- 28 • Increased frequency and severity of extreme heat events.
- 29 • Increased energy demand (particularly during peak summer periods).
- 30 • Increased frequency and severity of wildfire events.
- 31 • SLR (with increased salt water intrusion in the Delta).
- 32 • Changes in species distribution and ranges.
- 33 • Decreased number of species.

- 1 • Increased number of vector-borne diseases and pests (including impacts to agriculture).
- 2 • Altered timing of animal and plant lifecycles (phenology).
- 3 • Disruption of biotic interactions.
- 4 • Changes in physiological performance, including reproductive success and survival of plants and
- 5 animals.
- 6 • Changes in invasive species.
- 7 • Altered migration patterns of fishes, aquatic-breeding amphibians, birds and mammals.
- 8 • Changes in food (forage) base.
- 9 • Changes in habitat, vegetation structure, and plant and animal communities.

10 These changes have significant implications for water quality, water supply, flooding, aquatic
 11 ecosystems, energy generation, and recreation throughout the state. Several guidance documents
 12 have been drafted or have been published to discuss strategies to protect resources from climate
 13 change in California such as the 2009 California Climate Adaptation Strategy (California Natural
 14 Resources Agency 2009).

15 **29.5.1.3 Climate Change Effects on the Plan Area**

16 **Recent Trends**

17 Average annual temperatures in the Plan Area have increased approximately 0.9°F (0.53°C) during
 18 the period 1920–2003 (see Table 29-3). Local annual precipitation has increased by an average of
 19 approximately 1.7 inches (4.3 centimeters) during this same period. As discussed above, sea level in
 20 San Francisco Bay has risen approximately 7 inches (18 centimeters) over the last 100 years,
 21 affecting high tide events and salinity levels in the Delta. Hydrologic conditions in the Delta are
 22 largely determined by precipitation (amount, form, and timing) in the Sierra Nevada, as well as
 23 water management upstream (reservoir releases, diversions, operation of weirs, etc.), as opposed to
 24 local conditions. The average early spring snowpack in the Sierra Nevada has decreased by about
 25 10% during the last century, a loss of 1.5 MAF of snowpack storage (California Department of Water
 26 Resources 2008).

27 **Projections to 2100**

28 As shown in Table 29-3, by 2060, average annual temperatures in the Plan Area are projected to
 29 increase by 3°F (1.6°C), relative to current conditions. Average annual precipitation is projected to
 30 decrease slightly (approximately 0.16 inch [0.40 centimeter] during this same period).

31 It is important to note that, while the mean-annual amount of precipitation may only change slightly,
 32 the character of precipitation within the Sacramento and San Joaquin River basins is expected to
 33 change under warming conditions, resulting in more frequent rainfall events and less frequent
 34 snowfall events. Increased warming is expected to diminish the accumulation of snow during the
 35 cool season (i.e., late autumn through early spring) and the availability of snowmelt to sustain runoff
 36 during the warm season (i.e., late spring through early autumn). Warming may lead to more rainfall-
 37 runoff during the cool season rather than snowpack accumulation. This conceptually leads to
 38 increases in December–March runoff and decreases in April–July runoff.

1 **Table 29-3. Temperature, Precipitation, and Runoff Statistics for the Plan Area**

	Historical (1922– 1970)	Current (1970– 2003)	2025 Conditions	2060 Conditions
Average Temperature, Sacramento Basin	10.6°C	10.9°C	11.6°C	12.7°C
Average Temperature, San Joaquin Basin	11.5°C	12.0°C	12.7°C	13.7 °C
Average Temperature, Delta area	14.8°C	15.6°C	16.0°C	17.0°C
Average Precipitation, Sacramento Basin	86.3 cm	92.9 cm	88.6 cm	88.5 cm
Average Precipitation, San Joaquin Basin	63.1 cm	66.5 cm	63.2 cm	62.0 cm
Average Precipitation, Delta area	33.2 cm	35.9 cm	33.4 cm	32.9 cm
% of Runoff already arrived at Res. April 1 Sac Basin	73%	75%	80%	85%
% of Runoff already arrived at Res. April 1 SJ Basin	44%	45%	49%	55%

°C = degrees Celsius
cm = centimeters
Based on climate change scenarios and runoff data modeled for the BDCP alternatives (see Appendix 5A, *BDCP EIR/EIS Modeling Technical Appendix*)

2
3 Recent modeling indicates that sea level along the San Francisco Coast is expected to increase by 5
4 to 24 inches (12 to 61 centimeters) by 2050 and by as much as 17 to 66 inches (42 to 167
5 centimeters) by 2100 (National Research Council 2012). It is expected that more land in the Plan
6 Area will be subject to inundation by 2100 in comparison to current conditions. Potential changes in
7 inundation zones (tidal regime) will affect the salinity and suitable habitat for species in the Delta.

8 **Water Temperatures**

9 Reservoir operations may change temperatures below reservoirs, but will not affect temperatures in
10 the Delta. Increased water temperatures may have adverse effects on fish spawning (reduced egg
11 survival) and may reduce the habitat zone (reduced abundance) of fish that are sensitive to higher
12 temperatures (i.e., delta smelt). The projected effects of climate change on habitat and egg mortality
13 for the early long-term (2025) and late long-term (2060) timeframes were evaluated using three
14 water temperature models (BOR Temperature model, Sacramento River water quality model and
15 Delta temperature model). Specific modeling procedures and assumptions are further described in
16 Appendix 29C, *Climate Change and the Effects of Reservoir Operations on Water Temperatures in the*
17 *Study Area*.

18 **Precipitation and Runoff**

19 The projected effects of climate change on precipitation in the Central Valley were estimated using
20 general circulation model (GCM) results that were processed with a watershed hydrology model,
21 Variable Infiltration Capacity (VIC), to provide monthly runoff estimates for the CALSIM II planning
22 model. Two projections were developed with separate inflow sequences representing the early long-
23 term (2025) climate assumptions that included 5.9 inches (15 centimeters) of SLR, more variable
24 precipitation, and warmer temperatures, and late long-term (2060) climate assumptions that
25 included 17.7 inches (45 centimeters) of sea level rise, more variable precipitation than in the early
26 long-term, and warmer temperatures than in the early long-term. These potential climate conditions
27 were used to simulate the reservoir operations and Delta operations (export pumping) for each
28 BDCP alternative. The differences in these anticipated changes in the runoff sequences are fully

1 described in Appendix 29B, *Climate Change Effects on Hydrology in the Plan Area Used for CALSIM*
 2 *Modeling Analysis*.

3 **Sea Level Rise**

4 The likely effects of anticipated SLR on the Plan Area were evaluated based on detailed modeling
 5 simulations. When considering potential SLR impacts, special consideration must be given to the
 6 following three interrelated elements.

- 7 • **Inundation:** Changes in sea level have the potential to inundate previously dry areas. The extent
 8 of inundation in the Delta is sensitive to the magnitude of SLR. As discussed below, Figure 29-1
 9 depicts the changes in inundation at high tide assuming a 55 inch (140 centimeters) SLR.
- 10 • **Salinity Gradient:** The location of the gradient between saline, brackish, and freshwater in the
 11 San Francisco estuary will be affected by SLR. As sea levels rise, the salinity gradient will shift
 12 further upriver. The position of the daily average salinity gradient in the estuary is called “X2”,
 13 which is the distance in kilometers upstream of the Golden Gate Bridge of the 2 parts per
 14 thousand (ppt) isohaline, (1995 Bay-Delta Water Quality Control Plan [WQCP]). The X2 position
 15 is highly variable due to daily tidal movement. Outflow objectives identified in the WQCP
 16 manage the X2 position to control salinity intrusion into the Delta. The daily average X2 position
 17 provides a good index of the upstream extent of saltwater intrusion as a consequence of SLR.
 18 Maintaining the existing X2 position under future SLR scenarios will likely require increased
 19 outflows from the Delta.
- 20 • **Tidal Variations:** Changes in sea level will influence natural tidal variations along the California
 21 coast and within the San Francisco Bay and Delta. Edge species that rely on existing variations
 22 between wet and dry conditions may become permanently inundated or otherwise experience
 23 inhospitable environmental changes.

24 Best available information suggests a range of potential SLR from 17 to 66 inches (42 to 167
 25 centimeters) by 2100 (National Research Council 2012). Given the inherent variability in anticipated
 26 future scenarios, a broad range of potential sea level changes (from 6 to 55 inches) was analyzed.
 27 The projections from the NRC study were not used directly in the BDCP analysis for two reasons. 1)
 28 the study was published in June 2012, well after the modeling analysis for BDCP had been designed
 29 and performed, and 2) the projection years are not directly aligned with the 2025 and 2060 analysis
 30 periods used for BDCP. SLR projections for 2025 and 2060 were developed based on research
 31 available during the analysis design and based on the requirements of Water Code Section 85320,
 32 which required that BDCP evaluate a sea level rise of 55 inches (well in excess of the expected sea
 33 level described by any major study for 2060). The SLR projections used in the BDCP analysis at 2025
 34 and 2060 are consistent with the findings of the NRC and fall within the range of expected SLR that
 35 could be extrapolated from the NRC analyses at each analysis time period. The inclusion of
 36 additional analysis for 55 inches (140 centimeters) of SLR provides a conservative analysis of
 37 potential SLR late in the 21st century.

38 As discussed in Appendix 5A, *BDCP EIR/EIS Modeling Technical Appendix*, several models were used
 39 to assess and quantify the effects of SLR on the BDCP alternatives. Figure 29-2 identifies the three
 40 primary models used in the analysis, as well as how these models interact to predict tidal variations
 41 and other corresponding SLR effects in the Plan Area.

42 Climate and sea level change are global phenomena that can have unique impacts on local systems.
 43 As shown in Figure 29-2, the UnTRIM Bay-Delta Model (MacWilliams et al., 2009), a three

1 dimensional hydrodynamics and water quality model, was used to simulate localized impacts on
2 hydrodynamics and salinity transport in the Delta for a range of selected sea-level scenarios (6 to 55
3 inches [15 to 140 centimeters]). The results from the UnTRIM model were used to corroborate
4 (adjust coefficients to match) the RMA Bay-Delta Model (RMA 2005) and Delta Simulation Model
5 (DSM2) to correctly simulate tidal marsh restoration effects with and without SLR. Finally, the
6 DWR/Reclamation CALSIM II planning model was adjusted to match the salinity effects from SLR to
7 simulate CVP and SWP operation over the range of projected hydrologic conditions. Higher Delta
8 outflows were calculated to be required to meet the existing salinity objectives. Please refer to
9 Appendix 29A, *Effects of Sea-Level Rise on Delta Tidal Flows and Salinity*, for additional information
10 on modeling procedures and assumptions.

11 Potential changes in inundation at high tide as a consequence of 55 inches (140 centimeters) of SLR
12 are shown in Figure 29-1. Figure 29-1 is based on tidal elevation data developed as part of the Delta
13 Risk Management Strategy, Phase 1 (Phase 1 datasets) (California Department of Water Resources).
14 The Phase 1 datasets are projections of floodplain depths as a function of SLR scenarios (including
15 55 inches [140 centimeters]). Areas shaded in light yellow are at or below the high tide elevation
16 based on the current sea level. Areas shaded in orange are additional areas at or below high tide
17 elevation when a 55 inch (140 centimeters) rise in sea level is considered. Note that the yellow and
18 orange areas are not necessarily inundated due to control structures such as levees. Figure 29-1
19 provides insight as to which additional areas in the Delta may need to introduce or augment control
20 structures to avoid inundation should mean SLR increase by 55 inches (140 centimeters).

21 As shown in Figure 29-1, several communities with elevations greater than 17 feet (e.g., Fairfield,
22 Manteca, Tracy, and Brentwood) (5.2 meters) will likely not be directly affected by a 55 inch (140
23 centimeters)SLR. However, some of the Delta islands and other low lying areas may incur additional
24 inundation risk if 55 inches of SLR were to occur, especially if levees or other control structures
25 were to fail.

26 29.6 Resiliency and Adaptation Analysis

27 As described in Chapter 2, *Project Objectives and Purpose and Need*, the action alternatives seek to
28 make physical improvements to the SWP/CVP system which will serve to provide resiliency and
29 adaptability to rising sea levels and other reasonably foreseeable consequences of climate change.
30 The analysis below seeks to describe the manner in which the alternatives would achieve the stated
31 objective of increasing resiliency and adaptability to climate change over the No Action/No Project
32 Alternative. BDCP components that could affect the resilience and adaptability of the Plan Area to
33 climate changes consist of water diversion and conveyance facilities combined with differing
34 operational scenarios (collectively CM1), measures focused on the protection, restoration, and
35 enhancement of natural communities (CM2—CM11), and measures related to reducing other
36 stressors (CM12—CM22). These conservation measures and the components they comprise are
37 described in detail in Chapter 3, *Description of Alternatives*. Depending on the alternative, the water
38 facility components would create a new conveyance mechanism and operational guidelines to divert

1 water from the north Delta to existing SWP and CVP export facilities in the south Delta to achieve the
2 planning goals outlined in the BDCP³.

3 To the extent possible, detailed project specific analysis done for BDCP is reported to provide
4 evidence of the expected changes in resiliency and adaptability. Where no detailed project specific
5 analysis was available, references and or qualitative descriptions are included that provide evidence
6 that the described effect would provide a resiliency or adaptation benefit.

7 **29.6.1 Resiliency and Adaptability to Sea Level Rise and** 8 **Hydrology Changes**

9 **29.6.1.1 Water Supply Reliability and Aquatic Species in the Delta**

10 **Impacts**

11 Appendix 3E, *Potential Seismic and Climate Change Risks to SWP/CVP Water Supplies*, describes the
12 existing and future risks to the Plan Area and specifically to the Delta as a result of climate changes
13 described above. The appendix highlights how increased sea level and changes in upstream
14 hydrology will affect the Plan Area. For the BDCP analyses, potential sea level increases of 6 inches
15 (15 centimeters) in 2025 and 18 inches (46 centimeters) in 2060 were evaluated as was a sea level
16 rise of 55 inches (which is not projected to occur until 2099, but is evaluated consistent with the
17 requirements of California Water Code Section 85320). Expected changes in precipitation and
18 hydrology were also evaluated including earlier runoff as a result of warmer temperatures causing
19 more precipitation to fall as rain instead of snow and the remaining snow melting earlier. Additional
20 information about the analysis methodology and modeling assumptions can be found in Appendices
21 29A, *Effects of Sea-Level Rise on Delta Tidal Flows and Salinity*, and 29B, *Climate Change Effects on*
22 *Hydrology in the Study Area Used for CALSIM Modeling Analysis*.

23 Modeling results for the BDCP 2060 period indicate a shift in runoff from snowmelt months (April–
24 June) to snow/rainfall months (January–March) of about 5–10% for the Sacramento River Basin and
25 of about 5–7% for the San Joaquin River Basin. The total runoff was increased (over historical
26 conditions) slightly (2%) for the Sacramento River Basin and decreased (6–10%) for the San
27 Joaquin River Basin. While these change metrics represent long-term averages, modeling results for
28 the BDCP 2060 period also indicate that droughts will increase in severity and duration—resulting
29 in periods of critical dryness.

30 All of these climate changes may result in less water flowing into the Delta between March and
31 October. At the same time, higher sea levels, in the absence of intervention, will increase the
32 penetration of salinity into the Delta. This increased Delta salinity would have a myriad of impacts
33 on in-Delta and Delta export water users whose water quality would be diminished. Aquatic species
34 such as Delta smelt would also be affected by these changes as their habitats would shrink or move
35 to less productive areas as discussed in Chapter 11, *Fish and Aquatic Resources*, Section 11.3.4.1.
36 Interventions that could be taken to counteract additional salinity intrusion would likely include the
37 release of additional water from upstream storage reservoirs. These actions would have
38 corresponding tradeoffs as less water would be left in the reservoirs for other actions. Reduced

³ As described in Chapter 1, *Introduction*, Section 1.1, the full Draft EIR/EIS should be understood to include not only the EIR/EIS itself and its appendices but also the proposed BDCP documentation including all appendices.

1 water available for agricultural, municipal, and industrial water supplies would reduce reliability
2 and have economic costs. Reduced water available for instream and other ecological uses would
3 result in negative effects on upstream aquatic species including cold water pool resources, critical
4 for salmonid rearing.

5 All of the hydrologic changes discussed above will make water management more challenging and
6 more constrained in the future and are expected to result in more years of critical dryness. DWR's
7 modeling of future conditions suggests that with current management and operations, level of
8 demand, and current climate, major CVP and SWP reservoirs could reach dead storage levels (the
9 level below which water cannot be released) and that the likelihood of these critical conditions will
10 increase substantially as the climate warms. In these instances, there would be critical water
11 shortages leading to potentially extreme impacts to agriculture, municipal, industrial, and ecological
12 water uses.

13 **Resilience/Adaptation**

14 BDCP Alternatives 1A, 1B, 1C, 2A, 2B, 2C, 3, 4, and 5 would provide substantial resiliency and
15 adaptation benefits over the No Action/No Project alternative for dealing with the combined effect
16 of increases in sea level rise and changes in upstream hydrology. Implementation of any of these 9
17 alternatives result in an increase in Delta exports and total SWP and CVP water deliveries over the
18 No Action Alternative (Table 29-4). These alternatives have dual conveyance facilities, allowing
19 water to be moved through the Delta when conditions permit and allowing water to be diverted
20 from the Sacramento River in the northern Delta when conditions do not permit through Delta
21 conveyance. The location of the north Delta diversion facility is further inland making it less
22 vulnerable to salinity intrusion. Even with substantial sea level rise and critically dry upstream
23 conditions, salinity could be repelled from this location. By establishing an alternative diversion
24 point for Delta exports, a great deal of Delta management flexibility is added. Currently,
25 management of the Delta is constrained by requirements to maintain X2 at specific locations during
26 certain times of the year to ensure water diversions have low salinity and to ensure that critical fish
27 populations stay outside of the entrapment zone. Alternatives 1A thru 5 would allow the Delta to be
28 managed in a number of different ways, including maintaining salinity as it is currently managed or
29 allowing salinity to fluctuate more freely in the Delta as it did prior to the development of upstream
30 reservoirs. This added flexibility would allow managers more options for adaptively managing the
31 Delta so that conditions can be optimized to provide the greatest benefits across all Delta water uses
32 and habitat conditions.

33 As shown in Table 29-4, Alternatives 6 through 8 would slightly decrease Delta exports and total
34 SWP and CVP water deliveries over the No Action Alternative. Accordingly, these alternatives may
35 not add resiliency to existing water supplies. However, similar to Alternatives 1 through 5,
36 Alternatives 7 and 8 would have dual conveyance facilities, which could improve management
37 flexibility. The location of the north Delta diversion facility proposed under these alternatives, as
38 well as under Alternative 6, would also be further inland, making it less vulnerable to salinity
39 intrusion.

1 **Table 29-4. Delta Exports and CVP/SWP Deliveries**

Alternative	No Action/ No Project	1	2	3	4	5	6	7	8	9
Delta Exports	4,441	5,456	5,068	5,371	4,945	4,786	3,758	3,754	3,098	4,377
Change in Exports		1,015	627	930	504	345	(683)	(687)	(1,343)	(64)
Total CVP/SWP deliveries	6,553	7,371	7,060	7,321	6,995	6,868	5,902	5,942	5,273	6,496
Change in CVP/SWP deliveries (Alt-NAA)		818	507	767	442	315	(652)	(612)	(1,281)	(57)

2

3 For the analysis of BDCP alternatives, operation of the CVP and SWP systems are modeled using
4 current regulatory conditions and a set of operational strategies. While this provides a good
5 reference point for evaluating the potential operational benefits and impacts of the BDCP
6 alternatives, additional infrastructure constructed and ecological restoration implemented as
7 elements of the alternatives could also open up additional operational possibilities that could be
8 explored through the BDCP adaptive management process, thus allowing other operational
9 alternatives that could provide potentially larger benefits to Delta resources. There is currently a
10 high level of uncertainty about how different Delta conditions, including salinity, tidal habitat, Delta
11 outflow, water temperature, Delta water quality, and level of Delta exports would affect critical
12 aquatic species, which of these variables has the greatest effect on these species, and what the best
13 combination of management practices would be. Climate change responses add more uncertainty to
14 these variables and tighten the constraints within which the Delta can be managed. Alternatives 1A
15 thru 5 would also increase resilience and adaptability to this uncertainty by providing additional
16 management flexibility for in-Delta conditions.

17 In addition to added water management flexibility created by CM1, CM2–CM22 provide for actions
18 that will improve habitat and reduce the effects of other stressors on the Delta ecosystem. By
19 improving and expanding available habitat, the BDCP alternatives increase resilience and
20 adaptability to the climate changes described above by increasing the amount of alternative habitat
21 that is available during periods of high stress such as very high or low freshwater inflow or very
22 high salinity intrusion. By reducing other stressors on the Delta ecosystem, the BDCP alternatives
23 will improve the health of the ecosystem and of individual species populations making them
24 stronger and more resilient to the potential variability and extremes caused by climate change.
25 Below are some of the key ways in which CM2–CM11 improve resiliency and adaptability of aquatic
26 resources in the Delta to climate change.

- 27 • Restoration of wetlands, floodplains, and riparian habitats will restore ecosystem services,
28 including flow regulation, nutrient cycling, and sediment processes that enhance the functioning
29 of aquatic habitats (Mitsch and Gosselink 2000).
- 30 • Increased wetland plant biomass, including belowground production, helps to promote
31 accretion and the ability of the marsh to keep pace with sea-level rise (Callaway et al. 2011;
32 Parker et al. 2011).
- 33 • A wider and more extensive marsh plain in tidal wetlands and a wider floodplain in river
34 systems increase protection of upland habitat from flooding and storm surges, which are
35 projected to get worse with climate change (Cayan et al. 2008).

- 1 • Improved floodplain connections to rivers will restore the ability of floodplains to absorb flood
2 flows and provide a reservoir of water to help aquatic species withstand droughts.
- 3 • Seasonally inundated floodplains provide more resilience from invasive species by increasing
4 numbers and health of native species and excluding invasive species (Moyle et al. 2007).
- 5 • Restoration supports species diversity by providing a mosaic of habitats that can be used by
6 different species that have evolved to use specific habitats.
- 7 • Wetland restoration will include networks of channels within marshes that are used by fish for
8 foraging, refuge, and movement into and out of the marsh. Currently, such channels are rare
9 (Parker et al. 2011).

10 29.6.1.2 Terrestrial Habitat and Species

11 Impacts

12 SLR and hydrologic changes will also have potentially detrimental effects on important terrestrial
13 habitat and species in the Delta. In addition to sea level changes, changes in average precipitation,
14 and runoff timing (discussed above), there is one additional hydrologic effect of climate change that
15 could result in impacts to terrestrial species in the Plan Area: increased incidence and magnitude of
16 extreme precipitation events. This additional impact has not been included in the quantitative
17 modeling analysis done for BDCP because there remains high levels of uncertainty about the scale of
18 the effects and because BDCP hydrologic and operations modeling was not conducted at a time step
19 conducive to evaluating short duration extreme precipitation events⁴. Other analyses done for other
20 purposes suggest that extreme precipitation events may become more frequent and larger in the
21 future (Climate Action Team 2010; Dettinger 2011). While the effects of more extreme precipitation
22 events are not included in the quantitative analysis, the effects on terrestrial species and potential
23 for the BDCP alternatives to improve or reduce resilience and adaptability to increased frequency
24 and magnitude of extreme precipitation events are described here.

25 The remnant marshes of the Delta are habitat for several species listed under the Federal and State
26 Endangered Species Acts such as California black rail and Mason's lilaepsis. The Plan Area lies in a
27 central portion of the Pacific Flyway and continues to provide vital migratory, wintering, and
28 breeding habitat for migratory birds, especially in designated wildlife management areas
29 (e.g., Suisun Marsh and Yolo Bypass), where habitat management is optimized for managed species,
30 including waterfowl, shorebirds, and wading birds. Although waterfowl have been reduced in
31 numbers, the Delta still provides habitat for 26 species of wintering waterfowl (Bay Institute 1998).
32 The Pacific Flyway is also particularly important for shorebirds and neotropical migratory birds.
33 Although fragmented, limited riparian habitat remains in the Delta. Remnant patches of tall riparian
34 trees (e.g., Fremont cottonwood, western sycamore, Goodding's black willow) occur, but the

⁴ The hydrologic modeling done for BDCP was done on a monthly time step, as is typical for analysis of water management projects in California where flood protection is not a primary objective. However, extreme precipitation events often unfold over much shorter periods of time (usually 2–7 days). At a monthly time step, important details about how streamflows move through the system can be masked. Because flood protection is not a primary objective of the BDCP, analysis at a more detailed time step in order to evaluate these effects in detail is not necessary. Nonetheless, qualitatively, high flow events becoming more frequent or larger in the future could increase the vulnerability of terrestrial species in the Plan Area under the No Action/No Project Alternative and under the action alternatives.

1 reproduction of these species is greatly impaired by lack of active floodplain habitat and hydrologic
 2 modifications. Grasslands with vernal pools support high levels of endemic biodiversity in the
 3 Central Valley (Witham et al. 1998). This habitat type occurs in the northeast and southwest areas of
 4 the Delta.

5 Higher sea levels will inundate existing low lying terrestrial habitats described above, potentially
 6 destroying marshy and tidal habit and increasing species mortality or changes in distributions
 7 (California Department of Fish and Game 2010). Current Delta land use patterns, which devote most
 8 of the land to agricultural uses, provide habitat value for some species, but provide limited
 9 opportunities for migration of low lying terrestrial habitat as sea level rises. Terrestrial habit in the
 10 Delta is also likely to face higher risk of inundation or desiccation due to more extreme fluctuations
 11 in precipitation.

12 Resilience/Adaptation

13 The BDCP alternatives include measures to restore between 43,000 and 94,000 acres of new
 14 seasonally inundated floodplain, tidal wetland, valley/foothill riparian, grassland, vernal pool
 15 complex, and nontidal marsh habitat. Additionally, approximately 69,000 acres of natural
 16 communities would be protected and 20 or 40 miles of channel margin habitat would be enhanced.
 17 While the locations and specific characteristics of each of these restoration, enhancement, and
 18 protection activities are not yet fully known, the comprehensive analysis, selection, and
 19 implementation of these actions will allow resource managers to plan for habitat migration and
 20 transformation, providing greater resilience and adaptability to changing future conditions.
 21 Protection and restoration of a variety of natural communities will increase the patch size and
 22 connectivity of these habitats. Increasing patch size will tend to increase population sizes of native
 23 species, which provides more resilience against a changing climate. Increasing connectivity allows
 24 more genetic exchange among populations and movement to more suitable habitats as
 25 environmental conditions change. The expansion of habitat will also provide greater resilience and
 26 adaptability ensuring that alternate habitats exist if habitats in some locations are destroyed or
 27 degraded by expected or unforeseen climate changes or catastrophic events. BDCP measures that
 28 restore and protect habitat will also assist in protecting and restoring upland refuges for terrestrial
 29 species affected by changes in tidal influence thereby increasing resiliency. These upland refuges
 30 may also provide added resiliency and adaptability to more extreme precipitation events such as
 31 droughts and floods. The additional habitat will likely provide more possibilities for alternative
 32 habitat locations that are less impacted during temporary inundation or desiccation periods.
 33 Restoration activities can also provide opportunities to contribute to climate change mitigation by
 34 increasing the carbon sequestration potential of these habitats.

35 29.6.1.3 Delta Levee Stability and Reliability

36 Impacts

37 Whether increased sea levels are counteracted by increased outflows for salinity purposes or not,
 38 water levels in the Delta will rise as sea levels rise, placing additional stress on fragile Delta levees.
 39 In addition, increased likelihood and magnitude of extreme precipitation events, as described above,
 40 could also increase the vulnerability of Delta levees. This impact is described in greater detail in
 41 Appendix 3E, *Potential Seismic and Climate Change Risks to SWP/CVP Water Supplies*. These levees
 42 not only protect farmland but maintain hydrodynamic conditions in the Delta. Western Delta levees
 43 serve a critical function of restricting the flow of saline water into the interior Delta, central Delta

1 levees serve to direct freshwater inflows toward the south Delta pumping plants (reducing the
 2 amount of salinity that mixes with fresh water inflows). The additional stresses placed on these
 3 levees will increase the likelihood of levee failures, most notably from seepage and potentially result
 4 in catastrophic levee collapse. Depending on the location of the levee failure and hydrologic
 5 conditions at the time of the failure, a levee collapse could change the hydrodynamic balance in the
 6 Delta and lead to substantial salinity intrusion. Because the Delta serves as the conveyance system
 7 for SWP, CVP and local system exports and as the water source for in-Delta water users, a
 8 catastrophic levee collapse leading to salinity intrusion could interrupt water supplies to all of these
 9 water users for weeks or months while the levees are repaired and the salinity is flushed from the
 10 system. A catastrophic salinity intrusion could also have significant impacts on aquatic species as
 11 their habitat would also be affected.

12 **Resilience/Adaptation**

13 The BDCP alternatives, with the exception of Alternative 9, would not add resiliency to existing
 14 levees; levee fragility would remain high and increase with time as in the No Action/No Project
 15 Alternative. However, BDCP Alternatives 1A–8 would provide additional adaptability to catastrophic
 16 failure of Delta levees. By providing an alternate conveyance route around the Delta, Alternatives
 17 1A–8 provide a mechanism to continue making water deliveries to SWP/CVP contractors and local
 18 and in-Delta water users with conveyance interties even if the Delta were temporarily disrupted by
 19 a catastrophic levee failure. Alternative 9 adds additional resiliency to the Delta by strengthening
 20 and reinforcing levees critical to the through-Delta conveyance route, however, this alternative does
 21 not increase the adaptive capacity of the system.

22 **29.6.2 Resiliency and Adaptability to Increased Temperature**

23 **29.6.2.1 Water Demand**

24 **Impacts**

25 Increased air temperatures associated with climate change will lead to increased evapotranspiration
 26 that will increase the water demand for crops and vegetation (Anderson, et al, 2008). While
 27 additional factors such as increased CO₂, humidity, cloudiness, etc. will also influence water demand,
 28 agricultural water demand is expected to increase as a result of climate change (Climate Action
 29 Team 2010). Increased evaporation may also reduce water supplies in open water supply and
 30 conveyance facilities, such as canals and reservoirs.

31 **Resilience/Adaptation**

32 As shown in Table 29-5 below, modeling analysis of the BDCP alternatives indicates that
 33 Alternatives 1A, 1B, 1C, 2A, 2B, 2C, 3, 4, and 5 improve water supply reliability (i.e., increase the
 34 long-term average of Delta exports), and will therefore, provide more reliable water supplies which
 35 will provide additional resilience and adaptability to increases in water demand as a result of higher
 36 temperatures and increased evapotranspiration and evaporation. Alternatives 6A, 6B, 6C, 7, 8, and 9
 37 actually result in reduced water supply reliability and therefore provide reduced resilience and
 38 adaptability to the impacts of climate change.

1 **Table 29-5. Long-Term Average Exports (TAF)**

Alternative	Total Delta Exports
No Action/No Project	4,441
Alternatives 1A, 1B, 1C	5,456
Alternatives 2A, 2B, 2C	5,068
Alternative 3	5,371
Alternative 4	--
Scenario H1	5,255
Scenario H2	4,710
Scenario H3	4,945
Scenario H4	4,414
Alternative 5	4,786
Alternatives 6A, 6B, 6C	3,758
Alternative 7	3,754
Alternative 8	3,098
Alternative 9	4,377

2

3 **29.6.2.2 Water Temperatures**4 **Impacts**

5 Warmer water temperatures are expected to decrease suitable summer habitat of delta smelt, a
6 federally listed endangered species, because waters in the lower Delta may be too saline and lack
7 food, and fresh water in the upper Delta may be too warm (California Department of Water
8 Resources 2009). Warming of streams and rivers also facilitates colonization by invasive species
9 that will compete for native species' habitat (Kaushal et al. 2010).

10 **Resilience/Adaptation**

11 By creating a wider variety of water management options and restoring habitat on a large scale, the
12 BDCP can help buffer potential negative effects of increased water temperatures thereby adding
13 resiliency to increased water temperatures. More detail on existing temperature conditions in
14 watersheds within the Plan Area and water temperature effects on aquatic habitat as well as
15 biological and biochemical processes, and how managed flows influence water temperatures can be
16 found in Chapter 11, *Fish and Aquatic Resources*. Additional information about the analysis
17 methodology and modeling assumptions used in the analysis can be found in Appendix 29C, *Climate
18 Change and the Effects of Reservoir Operations on Water Temperatures in the Study Area*.

19 **29.7 Compatibility with Applicable Plans and Policies**

20 This section provides an overview of federal and statewide efforts to prepare for and adapt to
21 climate change. Regulations associated with the mitigation of GHG emissions (e.g., AB-32) are
22 discussed in Chapter 22, *Air Quality and Greenhouse Gases*, Section 22.2, and are not repeated here.

1 Constructing the proposed water conveyance facilities (CM1) and implementing CM2–CM22 could potentially
 2 result in incompatibilities with these plans and policies related to climate change. A number of plans and
 3 policies establish plans or guidance for resource protection, adaptation, and enhancement activities related to
 4 resources in the study area. This overview of plan and policy compatibility evaluates whether Alternatives
 5 1A-9 are compatible or incompatible with such enactments. This analysis is not required by NEPA or CEQA,
 6 but is instead performed here to provide full disclosure regarding the potential impacts the proposed project
 7 could have on the Plan Area in the future. Note that as discussed in Chapter 13, *Land Use*, Section 13.2.3, state
 8 and federal agencies are not generally subject to local land use regulations; incompatibilities with plans and
 9 policies are not, by themselves, physical consequences to the environment.

10 **29.7.1 Applicable Plans and Policies**

11 **29.7.1.1 Federal**

12 **Council on Environmental Quality**

13 **CEQ’s Draft NEPA Guidance on Consideration of the Effects of Climate Change and Greenhouse** 14 **Gas Emissions (2010)**

15 On February 19, 2010, the Council on Environmental Quality (CEQ) issued draft National
 16 Environmental Policy Act (NEPA) guidance on the consideration of the effects of climate change and
 17 GHG emissions. This guidance advises federal agencies that they should consider opportunities to
 18 reduce GHG emissions caused by federal actions, adapt their actions to climate change effects
 19 throughout the NEPA process, and address these issues in their agency NEPA procedures. Where
 20 applicable, the scope of the NEPA analysis should cover the GHG emissions effects of a proposed action
 21 and alternative actions, as well as the relationship of climate change effects on a proposed action or
 22 alternatives. The CEQ guidance is still considered draft as of the writing of this document and is not an
 23 official CEQ policy document (Council on Environmental Quality 2010).

24 **National Oceanic and Atmospheric Administration**

25 **Global Sea Level Rise Scenarios for the United States National Climate Assessment (2012)**

26 This report was produced in response to a request from the U.S. National Climate Assessment
 27 Development and Advisory Committee. It provides a synthesis of the scientific literature on global
 28 sea level rise, and a set of four scenarios of future global sea level rise. The report includes input
 29 from national experts in climate science, physical coastal processes, and coastal management.

30 **U.S. Fish and Wildlife Service**

31 **Rising to the Urgent Challenge: Strategic Plan for Responding to Accelerating Climate Change** 32 **(2010)**

33 This report establishes a 5-year framework to analyze fish and wildlife conservation strategies
 34 associated with climate change. Adaptation, which the U.S. Fish and Wildlife Service (USFWS) defines
 35 as “planned, science-based management actions,” forms the core of the Strategic Plan. The primary
 36 purpose of the Strategic Plan is to identify adaptive responses to climate change through the
 37 strategic conservation of terrestrial, freshwater, and marine habitats. USFWS will implement the
 38 Strategic Plan during the next five years. To the extent that USFWS actions target ecosystems in the
 39 Delta, climate change resiliency in the Plan Area may be improved.

1 **National Fish, Wildlife, and Plants Climate Adaptation Strategy (2013)**

2 The overarching goal of the National Fish, Wildlife, and Plants Climate Adaptation Strategy is to “inspire,
3 enable, and increase meaningful action that helps safeguard the nation’s natural resources in a changing
4 climate.” The strategy describes current and expected impacts of climate change on major ecosystems in
5 the United States, and describes steps that can be taken to reduce these impacts. The actions proposed by
6 the strategy address the following seven goals: 1) conserve and connect habitat, 2) manage species and
7 habitats, 3) enhance management capacity, 4) support adaptive management, 5) increase knowledge and
8 information, 6) increase awareness and motive action, and 7) reduce non-climate stressors.

9 **U.S. Environmental Protection Agency**

10 **Climate Ready Estuaries Program (ongoing)**

11 U.S. Environmental Protection Agency’s (EPA’s) Climate Ready Estuaries program has four primary
12 objectives related to climate change adaptation: (1) assess climate change vulnerabilities, (2) develop
13 and implement adaptation strategies, (3) engage and educate stakeholders, and (4) share the lessons
14 learned with other coastal managers. The program provides information and tools for managers to
15 develop adaptation plans for estuaries and coastal communities. Resources published by the Climate
16 Ready Estuaries program can provide guidance for adaptively managing estuaries in the Plan Area.

17 **U.S. Forest Service**

18 **Re-Framing Forest and Resource Management Strategies for a Climate Change Context (2008)**

19 This report provides a high-level, preliminary framework for developing forest and natural resource
20 management strategies in response to climate change in western mountainous environments. The report
21 summarizes an approach for developing adaptation and mitigation strategies using a “5-R strategy:”
22 increase resistance, promote resilience, enable response, encourage realignment, and implement
23 practices to reduce the human influence on climate. Strategies outlined in the document could provide
24 potential approaches for responding to climate change impacts on forests in the Plan Area.

25 **Climate Change Considerations in Project-Level NEPA Analysis (2009)**

26 This guidance document provides initial Forest Service guidance on how to consider climate change
27 in project-level NEPA analysis and documentation. This guidance document addresses how Forest
28 Service management may influence climate change mainly through incremental changes to global
29 pools of GHGs. This guidance will be revised as more scientific literature is published, climate
30 change management experience is gained, and government policies are established.⁵

31 **U.S. Department of Agriculture**

32 **2010–2015 Strategic Plan (2010)**

33 The 2010–2015 Strategic Plan outlines future initiatives the U.S. Department of Agriculture will
34 undertake to achieve its overall mission. Four strategic goals are outlined, of which one is to ensure
35 that national forests and private working lands are made more resilient to climate change.
36 Performance measures and strategies for meeting this objective are summarized in the Strategic
37 Plan, and may be applicable to land management in the delta.

⁵ Source: http://www.fs.fed.us/emc/nepa/climate_change/includes/cc_nepa_guidance.pdf

1 **U.S. Department of the Interior**

2 **National Fish, Wildlife and Plants Climate Adaptation Strategy (ongoing)**

3 The U.S. Fish and Wildlife Service, the National Oceanic and Atmospheric Administration, and the
 4 New York Division of Fish, Wildlife, & Marine Resources are developing a unified approach to
 5 maintaining the key terrestrial, freshwater and marine ecosystems needed to sustain fish, wildlife
 6 and plant resources and the services they provide in the face of accelerating climate change. This
 7 strategy will provide a unified approach—reflecting shared principles and science-based practices—
 8 for reducing the negative impacts of climate change on fish, wildlife, plants, and the natural systems
 9 upon which they depend.⁶

10 **WaterSMART (Sustain and Manage America’s Resources for Tomorrow) (ongoing)**

11 The WaterSMART program was established in February 2010 following passage of the SECURE
 12 Water Act, which authorizes federal water and science agencies to work together with state and
 13 local water managers to plan for climate change and the other threats to our water supplies.
 14 WaterSMART provides a framework for federal leadership and assistance on efficient water use,
 15 integration of water and energy policies, and coordination of water conservation activities. As the
 16 Department’s main water management agency, Reclamation plays a key role in implementing the
 17 WaterSMART program. Improving water management and supplies through the BDCP is a priority
 18 goal for the WaterSMART program. (Bureau of Reclamation 2013)

19 **U.S. Army Corps of Engineers**

20 **Climate Change Adaptation Plan and Report (2011)**

21 The purpose of this report is to develop practical and cost effective measures to reduce the
 22 vulnerability of national water conveyance infrastructure to climate change. Strategies for adaptive
 23 planning, design, construction, and maintenance are identified. The document also provides a
 24 framework for performing a vulnerability assessment.

25 **29.7.1.2 State**

26 **Relevant State Executive Order and California Water Code Section**

27 **Executive Order S-13-08**

28 This Executive Order requests that the National Academy of Sciences (NAS) convene an independent
 29 panel to complete the first California SLR Assessment Report and initiate an independent SLR
 30 science and policy committee made up of state, national and international experts. It requires that
 31 all state agencies that are planning construction projects in areas vulnerable to future SLR shall
 32 consider a range of SLR scenarios for the years 2050 and 2100 in order to assess project
 33 vulnerability and, to the extent feasible, reduce expected risks and increase resiliency to SLR. The
 34 order does not specify SLR scenarios, but it is worth noting that SLR projections for California of 16
 35 inches (41 centimeters) by 2050 and 55 inches (140 centimeters) by 2100 have been considered
 36 and/or used by multiple state agencies in impacts analyses and policy development. The executive

⁶ Source: <http://www.wildlifeadaptationstrategy.gov/>.

1 order also tasks the CA Natural Resource Agency to coordinate the development of a statewide
 2 climate change adaptation strategy which resulted in the 2009 California Climate Adaptation
 3 Strategy.

4 **California Water Code, Section 85320 (b)(2)(C)**

5 As noted earlier, Water Code, Section 85320 (b)(2)(C) requires that, to be a part of the Delta Plan,
 6 the BDCP EIR must analyze the potential effects of climate change, possible SLR up to 55 inches (140
 7 centimeters) (the high end of the projected range in SLR), and possible changes in total precipitation
 8 and runoff patterns on the conveyance alternatives and habitat restoration activities considered in
 9 the Plan Area. Please refer to Section 29.4.2.1 for additional detail on the quantitative modeling
 10 performed to evaluate these potential effects.

11 **California Department of Water Resources**

12 **Managing an Uncertain Future; Climate Adaptation Strategies for Water (2008)**

13 This report summarizes adaptation strategies that can be used by state and local water managers to
 14 improve the resiliency of the California water resources. The strategies are organized at the regional
 15 and state level and focus on investment planning and decision-making. The report was developed by
 16 DWR as part of the process of updating the California Water Plan, and was the basis for the water
 17 section of the 2009 California Climate Adaptation Strategy.

18 **California Water Plan Update (2009)**

19 Chapter 4 of Volume 2 of the California Water Plan (California Department of Water Resources
 20 2010c) focuses on the Delta. In this chapter, DWR highlights the need for adaptation strategies to
 21 improve the flexibility of water conveyance and storage. Improving water management will enable
 22 operators to store large volumes of water during periods of high flow for use during periods of low
 23 flow when water supply allocations are more competitive. In addition, streams and channels
 24 enlarged for conveyance and flood passage may incorporate riparian habitat improvements that are
 25 designed for varying hydrology and water management operations. Delta conveyance
 26 improvements incorporate flexibility that allows for increased water supply reliability of Delta
 27 supplies in light of climate change.

28 **Central Valley Flood Protection Plan (2012)**

29 Due to recent legislation, DWR is currently implementing the Central Valley Flood Management
 30 Program (CVFMP), which involves the preparation of a Central Valley Flood Protection Plan
 31 (CVFPP). The CVFPP outlines an approach for addressing climate change considerations for flood
 32 management in the Central Valley (California Department of Water Resources 2011). The following
 33 are relevant documents developed as part of the 2012 CVFMP that present climate change
 34 adaptation strategies for flood management.

- 35 • *2012 Central Valley Flood Protection Plan, Climate Change Scope Definition Work Group Summary*
 36 *Report.*
- 37 • *2012 Central Valley Flood Protection Plan, Climate Change Threshold Analysis Work Plan—Draft*
 38 *Technical Memorandum.*

39 Future management measures developed as part of the 2012 CVFMP and associated documents will
 40 have direct effects on the hydrology of the Bay-Delta system within the Plan Area.

1 **California Department of Fish and Wildlife**

2 **Unity, Integration, and Action: CDFW's Vision for Confronting Climate Change in California (2011)**

3 The California Department of Fish and Wildlife's (CDFW's) framework for addressing climate change
4 adaptation seeks to protect California's natural resources without compromising the economy. The
5 framework embodies CDFW's commitment to minimizing negative effects of climate change on the
6 state ecosystems through the development of adaptation measures that provide clear benefits to
7 terrestrial and marine ecosystems. CDFW is acutely aware of the uncertainties associated with
8 emerging climate science and is taking an approach that will allow CDFW to be both proactive and
9 adaptive through the use of a variety of planning tools and strategic initiatives. Specifically, their
10 adaptive management framework will allow for the continual improvement and adjustment of
11 management practices based on new information.

12 **California Department of Food and Agriculture**

13 **California Agricultural Vision: Strategies for Sustainability (2010)**

14 California Agricultural Vision (Ag Vision) was created by the California Department of Food and
15 Agriculture (CDFA) and the State Board to address challenges associated with sustainable
16 agriculture. The State Board has endorsed several actions to assure agricultural adaptation to
17 climate change. In particular, research is currently being conducted to determine the most likely
18 impacts of climate change on agriculture, and to propose strategies to help agriculture adapt to and
19 benefit from these changes. Strategies developed as part of Ag Vision's research may benefit
20 agriculture in the delta.

21 **California Department of Public Health**

22 **Climate Change and Public Health: Building Healthy Communities and a Healthy Planet (ongoing)**

23 California Department of Public Health (DPH) has developed a four-part webinar series to educate
24 health professionals on how climate change will impact health across California, especially within
25 vulnerable communities. The webinar provides tools to communicate the need for action at the local
26 level. The webinars also share how cities and counties throughout California are planning for
27 climate change and how health and equity can be integrated into those efforts. Assistance provided
28 by DPH may help buffer delta communities from changes in climate.

29 **California Natural Resources Agency**

30 **California Climate Change Adaptation Strategy (2009)**

31 In cooperation and partnership with multiple state agencies, the 2009 California Climate Adaptation
32 Strategy summarizes the best known science on climate change and provides recommendations on
33 how to manage against those threats. The report provides an update on the expected climate risks to
34 California, prioritizes solutions to addressing these risks, and develops an implementation plan for
35 minimizing risks. The adaptation strategy will reduce California's vulnerability to known and
36 projected climate change impacts. The California Natural Resources Agency is currently in the
37 process of updating the 2009 California Climate Adaptation Strategy. The update will be released for
38 public comment by the end of 2013.

1 **California Department of Forestry and Fire Projection**

2 **CAL FIRE Report on Adaption Strategies for Forestry (2008)**

3 This report reviews many of the observed and forecasted impacts to California forests and
4 rangelands as a result of climate change. The document proposes a framework for developing
5 adaptation strategies. More specifically, the report identifies an initial strategy for integrating
6 adaptation into future forest management. Strategies outlined in the document could provide
7 potential approaches for responding to climate change impacts on forests in the Plan Area.

8 **CALFED Bay-Delta Program**

9 **Independent Science Board Memorandum (2007)**

10 CALFED Independent Science Board (ISB) is a multidisciplinary panel that provides guidance on
11 climate change and water issues. The CALFED ISB recently prepared a memo recommending
12 appropriate SLR projections for ongoing delta planning. In addition, the CALFED Science Program
13 has funded an effort to develop a model-based approach for evaluating plausible future scenarios of
14 the Bay-Delta-River-Watershed system. The outcome is intended to be a strategic planning tool to
15 CALFED agency managers and decision-makers in meeting future delta resource management goals.

16 **Delta Protection Commission**

17 **2006–2011 Strategic Plan (2006)**

18 The Delta Protection Commission's (DPC's) Strategic Plan is intended to protect and enhance the
19 Delta's resources. The document summarizes current and future threats to the Delta, including
20 changes in climate. In particular, the document identifies SLR as a central threat facing the future
21 integrity of the Delta.

22 **Ocean Protection Council**

23 **SLR Task Force Interim Guidance Document**

24 This document provides guidance for incorporating sea-level rise projections into planning and
25 decision making for projects in California. Its underlying premise is that SLR potentially will cause
26 many harmful economic, ecological, physical and social impacts and that incorporating SLR into
27 agency decisions can help mitigate some of these potential impacts. For example, SLR will threaten
28 water supplies, coastal development, and infrastructure, but early integration of projected SLR into
29 project designs will lessen these potential impacts.⁷

30 **California Department of Transportation**

31 **Climate Change Adaptation Hot Spot Map (ongoing)**

32 The Climate Change Adaptation Hot Spot Map is a GIS-based assessment of transportation
33 infrastructure vulnerabilities using available data and studies and to identify critical transportation

⁷ Source: http://opc.ca.gov/webmaster/ftp/pdf/agenda_items/20110311/12.SLR_Resolution/SLR-Guidance-Document.pdf.

1 hotspots. Such hotspots are areas of increased vulnerability to the effects of climate change due to
 2 their location near population centers that depend on transportation infrastructure for essential
 3 services; are heavily traveled, so a compromise in infrastructure would affect large numbers of
 4 individuals; or are particularly situated geographically to be heavily impacted by climate effects
 5 (e.g., on the coast in an area that could be inundated by rising sea levels). This research will also
 6 result in the development of a climate vulnerability plan that will assess the level and type of
 7 transportation infrastructure vulnerability, the adaptation options and strategies, and a framework
 8 for prioritizing implementation efforts. California Department of Transportation (Caltrans) will
 9 develop a framework for prioritizing implementation efforts throughout the Delta and California.

10 **Addressing Climate Change Adaptation in Regional Transportation Plans: A Guide for California**
 11 **MPOs [Metropolitan Planning Organizations] and Regional Transportation Plans Agencies**
 12 **(ongoing)**

13 The 2010 Regional Transportation Plans Guidelines currently provide little direction for regions to
 14 analyze and address climate change adaptation. This effort will provide the data to develop a clear
 15 methodology for regional agencies to address climate change impacts through adaptation of
 16 transportation infrastructure. The purpose of this manual is to expand knowledge and develop tools
 17 that will assist California MPOs and Regional Transportation Plans. As with incorporating climate
 18 change impacts into planning, design, engineering, and operational decisions. The final product will
 19 be a literature review of adaptation, best practices for regional agencies, and available adaptation
 20 measures for transportation infrastructure.

21 **29.7.2 Compatibility Evaluation**

22 The USFWS, EPA, U.S. Forest Service, U.S. Department of Agriculture, U.S. Department of the Interior,
 23 U.S. Army Corp of Engineers, CVFMP, DWR, CDFW, CDFA, DPH, California Natural Resources Agency,
 24 California Department of Forestry and Fire Protection, Delta Protection Commission, and Caltrans
 25 have developed frameworks or initiatives to ensure their respective resources are made more
 26 resilient to climate change. Construction and operation of the proposed water conveyance facilities
 27 and implementation of other conservation measures would not affect the ability of these agencies to
 28 implement these plans and proactive measures. Accordingly, the project would be compatible with
 29 these federal and state plans to address climate change.

30 The CEQ has prepared draft guidance on how federal agencies should consider the effects of climate
 31 change in their evaluation proposals. Consistent with the draft guidance, this chapter evaluates the
 32 relationship of climate change effects to the proposed project and alternatives. BDCP is therefore
 33 compatible with the CEQA guidance on climate change.

34 Executive Order S-13-8, California Water Code, Section 85320 (b)(2)(C), CALFED *Independent*
 35 *Science Board Memorandum*, and the OPC SLR Task Force *Interim Guidance Document* address
 36 expected risk and vulnerability to future SLR in California. The Water Code specifically requires that,
 37 to be a part of the Delta Plan, the BDCP EIR must analyze the potential effects of climate change,
 38 possible SLR up to 55 inches (140 centimeters) (the high end of the projected range in SLR), and
 39 possible changes in total precipitation and runoff patterns on the conveyance alternatives and
 40 habitat restoration activities considered in the Plan Area. Given the inherent variability in
 41 anticipated future scenarios, a broad range of potential sea level changes (from 6 to 55 inches [15 to
 42 140 centimeters]) was analyzed (see Appendix 5A, *BDCP Modeling Technical Appendix*). Because

1 potential effects of SLR on BDCP were analyzed as part of this analysis, the project is considered
 2 compatible with applicable SLR guidance documents.

3 **29.8 References**

4 **29.8.1 Printed References**

- 5 Anderson, J., F. Chung, M. Anderson, L. Brekke, D. Easton, M. Ejeta, R. Peterson, and R. Snyder, 2008.
 6 Progress on Incorporating Climate Change into Management of California's Water Resources.
 7 *Climatic Change* 89, Supplement 1 (March):91-108. Netherlands: Springer.
- 8 Auffhammer, M. 2011. Potential Impacts of Extreme Events on Electrical Energy Demand and
 9 Infrastructure. In *Proceedings from Vulnerability and Adaptation to Extreme Events in California*
 10 *in the Context of a Changing Climate: New Scientific Findings*. Scripps Institute of Oceanography.
 11 Tuesday, December 13, 2011.
- 12 Bay Institute. 1998. *From the Sierra to the Sea: An Ecological History of the San Francisco Bay-Delta*
 13 *Watershed*. Novato, CA: The Bay Institute of San Francisco.
- 14 Bureau of Reclamation. 2011. *SECURE Water Act Section 9503(c) – Reclamation Climate Change and*
 15 *Water 2011*. April. Available: <[http://www.usbr.gov/climate/SECURE/docs/](http://www.usbr.gov/climate/SECURE/docs/SECUREWaterReport.pdf)
 16 [SECUREWaterReport.pdf](http://www.usbr.gov/climate/SECURE/docs/SECUREWaterReport.pdf)>. Accessed: March 8, 2012>.
- 17 Bureau of Reclamation. 2013. *WaterSMART*. Last Revised: June 12, 2013. Available:
 18 <<http://www.usbr.gov/WaterSMART/index.cfm>>. Accessed: July 25, 2013.
- 19 California Department of Fish and Game. 2010. *Climate Change: Confronting the Challenge*. Fall.
 20 Available at <<http://nrm.dfg.ca.gov/FileHandler.ashx?DocumentID=31839&inline=true>>.
 21 Accessed: March 8, 2012.
- 22 California Department of Water Resources. 2008. *Managing an Uncertain Future. Climate Change*
 23 *Adaptation Strategies for California's Water*. October.
- 24 ———. 2009. *2012 Central Valley Flood Protection Plan. Climate Change Scope Definition Work Group*
 25 *Summary Report*. Draft. December.
- 26 ———. 2010a. *Model CEQA Climate Change Discussion and Impact Analysis Section*. Prepared by the
 27 California Department of Water Resources CEQA Climate Change Committee. January.
- 28 ———. 2010b. *Climate Change Characterization and Analysis in California Water Resources Planning*
 29 *Studies*. Final Report. December.
- 30 ———. 2010c. *California Water Plan Update 2009*. Available:
 31 <<http://www.waterplan.water.ca.gov/cwpu2009/>>. Accessed: March 8, 2012.
- 32 ———. 2011. *2012 Central Valley Flood Protection Plan*. Executive Summary. Public Draft. December.
- 33 California Energy Commission. 2011a. *California Climate Change Portal*. Available:
 34 <<http://www.climatechange.ca.gov/background/index.html>>. Accessed: November 4, 2011.
 35 Last Modified: May 2, 2008.

- 1 California Natural Resources Agency. 2009. *2009 California Climate Adaptation Strategy: A Report to*
2 *the Governor and Legislature in Response to Executive Order S-13-08*. December 2. Available:
3 <<http://www.climatechange.ca.gov/adaptation/>>. Accessed: March 8, 2012.
- 4 California Natural Resources Agency. 2013. *Our Changing Climate 2012: Vulnerability & Adaptation*
5 *to increasing Risks from Climate Change in California*. Available: <[http://www.energy.ca.gov/](http://www.energy.ca.gov/2012publications/CEC-500-2012-007/CEC-500-2012-007.pdf)
6 [2012publications/CEC-500-2012-007/CEC-500-2012-007.pdf](http://www.energy.ca.gov/2012publications/CEC-500-2012-007/CEC-500-2012-007.pdf)>. Accessed: August 1, 2013.
- 7 Callaway, J. C., V. T. Parker, M. C. Vasey, L. M. Schile, and E. R. Herbert. 2011. Tidal Wetland
8 Restoration in San Francisco Bay: History and Current Issues. *San Francisco Estuary and*
9 *Watershed Science* 9(3):1-12. Available: <<http://www.escholarship.org/uc/item/5dd3n9x3>>.
- 10 Cayan, D., M. Tyree, M. Dettinger, H. Hidalgo, T. Das, E. Maurer, P. Bromirski, N. Graham and R. Flick.
11 2009. *Climate Change Scenarios and SLR Estimates for the California 2008 Climate Change*
12 *Scenarios Assessment*. Prepared by the California Climate Change Center for the California
13 Energy Commission. CEC-500-2009-014-D. Sacramento, CA. Available:
14 <<http://www.energy.ca.gov/2009publications/CEC-500-2009-014/CEC-500-2009-014-D.PDF>>.
15 Accessed: March 8, 2012.
- 16 Climate Action Team. 2010. *Climate Action Team Biennial Report*. April.
- 17 Dettinger, M. 2011. Climate Change, Atmospheric Rivers, and Floods in California – A Multimodel
18 Analysis of Storm Frequency and Magnitude Changes. *Journal of the American Water Resources*
19 *Association (JAWRA)* 47(3):514–523.
- 20 Dettinger, M. and Ralph, M. 2011. Storms, Floods, and the Science of Atmospheric Rivers. *Eos* 92(32).
21 August 9.
- 22 Gershunov, A. 2011. Expected Changes in Key Weather-Related Extreme Events in California. In
23 *Proceedings from Vulnerability and Adaptation to Extreme Events in California in the Context of a*
24 *Changing Climate: New Scientific Findings*. Scripps Institute of Oceanography. Tuesday,
25 December 13, 2011.
- 26 Griggs, G. 2011. Coastal Hazards, SLR and Extreme Events. In *Proceedings from Vulnerability and*
27 *Adaptation to Extreme Events in California in the Context of a Changing Climate: New Scientific*
28 *Findings*. Scripps Institute of Oceanography. Tuesday, December 13, 2011.
- 29 Huber-Lee, A., Yates, D., Purkey, D., Yu, W., and Runkle, B. 2003. *Water, Climate, Food, and*
30 *Environment in the Sacramento Basin. Contribution to the Project ADAPT: Adaptation Strategies*
31 *to Changing Environments*. Available: <[http://www.weap21.org/downloads/](http://www.weap21.org/downloads/adapt_sacramento.pdf)
32 [adapt_sacramento.pdf](http://www.weap21.org/downloads/adapt_sacramento.pdf)>. Accessed: May 21, 2012.
- 33 Intergovernmental Panel on Climate Change. 2007a. *Climate Change 2007: The Physical Science Basis.*
34 *Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel*
35 *on Climate Change*. S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor
36 and H. L. Miller (eds.). New York, NY: Cambridge University Press. Available:
37 <http://www.ipcc.ch/publications_and_data/publications_and_data_reports.shtml>. Accessed:
38 March 8, 2012.
- 39 ———. 2007b. *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working*
40 *Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. M. L.
41 Parry, O. F. Canziani, J. P. Palutikof, P. J. van der Linden and C. E. Hanson (eds.). New York, NY:

- 1 Cambridge University Press. Available: <[http://www.ipcc.ch/publications_and_data/
2 publications_and_data_reports.shtml](http://www.ipcc.ch/publications_and_data/publications_and_data_reports.shtml)>. Accessed: March 8, 2012.
- 3 ———. 2011. *Organization*. Available: <<http://www.ipcc.ch/organization/organization.shtml>>.
4 Accessed: September 29, 2011.
- 5 Jackson, L., D. Lobell, C. Field, M. Hanemann, R. Howitt, and A. Gunasekara. 2011. Extreme Events and
6 Agriculture Production in California. In *Proceedings from Vulnerability and Adaptation to
7 Extreme Events in California in the Context of a Changing Climate: New Scientific Findings*. Scripps
8 Institute of Oceanography. Tuesday, December 13, 2011.
- 9 Kaushal S. S., G. E. Likens, N. A. Jaworski, M. L. Pace, A. M. Sides, D. Seekell, K. T. Belt, D. H. Secor, and
10 R. L. Wingate. 2010. Rising Stream and River Temperatures in the United States. *Frontiers in
11 Ecology and the Environment* 8:461–466.
- 12 MacWilliams, M. L., F. G. Salcedo, and E. S. Gross. 2009. *Draft San Francisco Bay-Delta UnTRIM Model
13 Calibration Report, Sacramento and Stockton Deep Water Ship Channel 3-D Hydrodynamic and
14 Salinity Modeling Study*. Prepared for US. Army Corps of Engineers, San Francisco District. July.
- 15 Mitsch, W. J. and J. G. Gosselink. 2000. *Wetlands 3*. Canada: John Wiley and Sons.
- 16 Moser, S., G. Franco, S. Pittiglio, W. Chou, and D. Cayan. 2009. *The Future is Now: An Update on
17 Climate Change Science Impacts and Response Options for California (Special Report California
18 Climate Change Center)*. Prepared for the California Energy Commission. CEC-500-2008-071.
19 Sacramento, CA.
- 20 Moyle, P. B., P. K. Crain, and K. Whitener. 2007. Patterns in the Use of a Restored California
21 Floodplain by Native and Alien Fishes. *San Francisco Estuary and Watershed Science* 5(3):1–27.
- 22 National Research Council. 2012. *Sea-Level Rise for the Coasts of California, Oregon, and Washington:
23 Past, Present, and Future*. National Academies Press.
- 24 Parker, V. T., J. C. Calloway, L. M. Schile, M. C. Vasey, and E. R. Herbert. 2011. Climate Change and San
25 Francisco Bay–Delta Tidal Wetlands. *San Francisco Estuary and Watershed Science* 9(3).
- 26 Rahmstorf, S. 2007. A Semi-Empirical Approach to Projecting Future Sea-Level Rise. *Science*
27 315:368–370. DOI: 10.1126/science.1141283.
- 28 Resource Management Association, Inc. (RMA). 2005. *Flooded Island Pre-feasibility Study: RMA Delta
29 Model Calibration Report*. June.
- 30 San Francisco Bay Conservation and Development Commission. 2009. *Living with a Rising Bay:
31 Vulnerability and Adaptation in San Francisco Bay and on the Shoreline*. Draft Staff Report. San
32 Francisco, CA. Available: <http://www.bcdc.ca.gov/proposed_bay_plan/bp_amend_1-08.shtml>.
33 Accessed: November 22, 2011.
- 34 U.S. Army Corps of Engineers. 2002. *Comprehensive Study, Sacramento and San Joaquin River Basins*.
35 Sacramento, CA.
- 36 U.S. Geological Survey. 2000. *Sacramento–San Joaquin Delta: The Sinking Heart of the State*. Prepared
37 by S. E. Ingebritsen, M. E. Ikehara, D. L. Galloway, and D. R. Jones. USGS Fact Sheet 005-00.
38 Available: <<http://pubs.usgs.gov/fs/2000/fs00500/>>. Accessed: March 8, 2012.

- 1 Western Regional Climate Center. 2012. *Climate Summary*. Available:
- 2 <<http://www.wrcc.dri.edu/cgi-bin/cliMAIN.pl?ca7633>>. Accessed: June 5, 2012.
- 3 Witham, C., E. T. Bauder, D. Belk, W. R. Ferren Jr., and R. Ornduff (eds.). *Ecology, Conservation, and*
- 4 *Management of Vernal Pool Ecosystems. Proceedings from a 1996 Conference*. Sacramento, CA:
- 5 California Native Plant Society.