

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/232669266>

Use of artificial basking substrate to detect and monitor pacific pond turtles (Emys marmorata)

Article *in* Western North American Naturalist · January 2006

DOI: 10.3398/1527-0904(2006)66[129:UOABST]2.0.CO;2

CITATION

1

READS

865

1 author:



[Jeff A. Alvarez](#)

The Wildlife Project

82 PUBLICATIONS 128 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



California tiger salamander natural history investigations [View project](#)



Western Pond Turtle Natural History Investigations [View project](#)

USE OF ARTIFICIAL BASKING SUBSTRATE TO DETECT AND MONITOR PACIFIC POND TURTLES (*Emys marmorata*)

Alvarez, Jeff A. The Wildlife Project, P.O. Box 579805, Modesto, CA 95357, azoologist@SBCglobal.net

Key words: artificial, basking, construction, emydid, monitor, platform, turtle.

Many species of turtles, lizards, snakes, and crocodylians are known to bask diurnally. Cagle (1950) suggested that basking in emydid turtles is done in response to the need to thermoregulate, to condition the skin or shell, and to retard epizootic and epiphytic infestations. However, Boyer (1965) determined through field and laboratory study, that the basking response in turtles is driven primarily by the need to thermoregulate. Many semi-aquatic turtles are known to bask on rocks, logs, vegetation mats, and floating debris (Lindeman 1999, Bury and Wolfheim 1973, Boyer 1965, Cagle 1950). Petokas and Alexander (1979) developed a trap that utilized the basking behavior in semi-aquatic turtles in order to facilitate trapping them. Bury and Wolfheim reported on the aggressive interactions of basking Pacific pond turtles (*Emys marmorata*).

Because Pacific pond turtles are declining in number in the state of California, it has become increasingly important to detect them within their habitat (Jennings and Hayes 1994). The purpose of this study was to demonstrate that an artificial basking substrate (basking platform) could be easily constructed and deployed, and with this technique presence of aquatic turtles could be detected and monitored within simple and complex aquatic habitats.

The Los Vaqueros Watershed (watershed), which is part of the upper Kellogg Creek drainage, was located ca. 57 km east of San Francisco, California. Habitat consisted primarily of annual and perennial grassland, oak (*Quercus* spp.) woodlands, riparian woodland, perennial drainages, ephemeral and seasonal wetlands, chaparral, open water, and rock outcrops. Los Vaqueros Reservoir, Kellogg Creek and its tributaries, and 40 perennial stock ponds provided suitable habitat for aquatic turtles. Fifty additional ephemeral stock ponds and wetlands provided refuge for dispersing turtles.

Fourteen rectangular basking platforms were constructed in two different designs (Figure 1). One platform type was made primarily of wood with added floatation (n = 8). The second design was constructed using a foam panel and also included additional floatation, but to a lesser extent (n = 6). Platforms were floated at an oblique angle to the water surface by attaching one or more 60 cm capped polyvinyl chloride tubes (Figure 1).

Baseline surveys (prior to the placement of basking platforms) were conducted throughout the watershed in 1998 for southwestern pond turtles (*E. m. pallida*). Biologists surveyed ponds and creeks on foot by initially scanning the water surface and shoreline using binoculars from a distance. They also walked along the perimeter of ponds and creeks and noted where southwestern pond turtles were observed.

In early spring 1999, basking platforms were placed in nine ponds and two creeks where aquatic emydid turtles were known to occur historically, or where habitat appeared suitable but

where no turtles had previously been found. Platforms were placed in open water and each platform was anchored to the bottom with a length of nylon cord and a concrete weight. Turtles were given 2-7 days to acclimate to the new structures.

Surveys for turtles were made 5-6 times per year for up to five years beginning several days after basking platforms had been deployed. Additional observations of turtles using basking platforms were recorded coincident with mitigation monitoring survey efforts for California red-legged frogs (*Rana draytonii*) in ponds and creeks in the watershed.

Baseline surveys in 1998 indicated southwestern pond turtles occupied two ponds and two creeks in the watershed. A maximum of 11 southwestern pond turtles were observed basking on the bank of one of the two ponds; two southwestern pond turtles were detected in the second pond. Although southwestern pond turtles were observed in the two creeks, no attempt was made to determine their numbers.

Within 2-7 days after deployment of the basking platforms, southwestern pond turtles were detected in eight of the nine ponds and confirmed in both creeks. The maximum number of turtles detected in ponds was 49 (34 turtles in one pond). Moreover, southwestern pond turtles were detected in six ponds where they were not previously known.

Within occupied habitat, observations of basking emydid turtles are relatively common when appropriate basking substrate is present (Reese and Welsh 1998, Bury and Wolfheim 1973, pers. obs). However, complex aquatic habitats (i.e., ponds or creeks with dense emergent and/or riparian vegetation) may reduce or eliminate the ability of an observer to detect turtles. Prior to this study, emergent vegetation in five of the nine ponds where basking platforms were deployed was so dense that those ponds were considered impractical to survey and no southwestern pond turtles were believed to occur. However, after basking platforms were deployed, 3-11 southwestern pond turtles were observed in each of those eight ponds with a highly complex habitat.

In the State of Washington, the northwestern pond turtles (*E. m. marmorata*.) could be detected in lakes and ponds in Washington State by supplying the turtles with basking substrate (Nordby, unpublished report, Stringer, unpublished report). In my study, detection and observations of southwestern pond turtles increased dramatically through the use of basking platforms. Observational studies and surveys of basking southwestern pond turtles at the Los Vaqueros watershed were performed more efficiently after deployment of basking platforms. Timing and temperature relationships of basking (Alvarez, unpublished report), and length of basking time among different age classes (Wilkerson, unpublished report) were both studied using the basking platforms. Further, ongoing monitoring of the presence, distribution, and abundance of this population of southwestern pond turtles is conducted at the Los Vaqueros Watershed using the basking platforms described in this study. These platforms were also deployed in Los Vaqueros Reservoir as a means to attract and monitor non-native species of emydid turtle that may be introduced into the watershed. Species using basking platforms within the reservoir were identified using a spotting scope, and trapping was used to remove non-native species within areas where they were detected.

The basking platform designs described above have proven durable; platforms have lasted up to six years in the field. The designs offered here are inexpensive and easily transportable. Costs for materials used to construct basking platforms ranged from \$9.00 to \$19.00 depending upon the style selected. This technique can be used as part of a short-term presence-absence survey by deploying the basking platforms at least seven days prior to a survey

and returning to observe animals that may be basking during the day. Behavioral studies of basking turtles can also be facilitated through the use of this technique.

I thank M. A. Shea for assistance in the field setting up platforms and observing turtles, and for comments on the manuscript. I also thank J. L. Alvarez for constructive discussions and helpful comments on the manuscript. Valuable editorial comments were received from R. W. Baumann and two anonymous reviewers. The Contra Costa Water District provided access to the habitats within the Los Vaqueros watershed for which I am grateful.

LITERATURE CITED

- BOYER, D. R. 1965. Ecology of the basking habit in turtles. *Ecology* 46:99-118.
- BURY, R. B. AND J. H. WOLFHEIM. 1973. Aggression in free-living pond turtles (*Clemmys marmorata*). *BioScience* 23:659-662.
- CAGLE, F. R. 1950. The life history of the slider turtle, *Pseudemys scripta troostii* (Holbrook). *Ecological Monographs* 20:31-54.
- JENNINGS, M. R. AND M. P. HAYES. 1994. Amphibian and reptile species of special concern in California. Report prepared for the California Department of Fish and Game, Inland Fisheries Division, Rancho Cordova, California. 255 pp.
- LINDEMAN, P. V. 1999. Aggressive interactions during basking among four species of emydid turtles. *Journal of Herpetology* 33:214-219.
- PETOKAS, P. J., AND M. M. ALEXANDER. 1973. A new trap for basking turtles. *Herpetological Review* 10:90.
- REESE, D. A. AND H. H. WELSH. 1998. Habitat use by western pond turtles in the Trinity River, California. *Journal of Wildlife Management* 62:843-853.

Submitted for publication to *Western North American Naturalist*

Submitted: 20 July 2004


Accepted: 01 February 2005

Publication date: fall 2005

Alvarez, J. A. *In press*. Use of artificial basking substrate to detect and monitor Pacific pond turtle (*Emys marmorata*). *Western North American Naturalist*.

Figure 1. Design and floating position of 2 styles of basking platform used to detect and monitor Pacific pond turtles at the Los Vaqueros Watershed, east Contra Costa County, California. Design B was constructed from a foam panel that was covered by artificial grass (i.e. "astro-turf"). The artificial grass is glued to the top and 4 sides, and is folded over and glued along the bottom edge of the foam panel.

Design A.



Design B.

