



United States
Department of
Agriculture


NRCS

Natural
Resources
Conservation
Service

A product of the National
Cooperative Soil Survey,
a joint effort of the United
States Department of
Agriculture and other
Federal agencies, State
agencies including the
Agricultural Experiment
Stations, and local
participants

Custom Soil Resource Report for Sutter County, California, and Yolo County, California

Sites-South



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (<http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/>) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (<https://offices.sc.egov.usda.gov/locator/app?agency=nrcs>) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
How Soil Surveys Are Made	5
Soil Map	8
Soil Map.....	9
Legend.....	10
Map Unit Legend.....	11
Map Unit Descriptions.....	12
Sutter County, California.....	14
121—Columbia fine sandy loam, frequently flooded, 0 to 2 percent slopes.....	14
122—Columbia loam, 0 to 2 percent slopes.....	15
177—Water.....	16
Yolo County, California.....	17
AaA—Arbuckle gravelly loam, 0 to 2 percent slopes, MLRA 17.....	17
BrA—Brentwood silty clay loam, 0 to 2 percent slopes.....	18
Ca—Capy silty clay, 0 percent slopes, MLRA 17.....	19
Cb—Capay silty clay, 0 percent slopes, frequently flooded, MLRA 17.....	21
Ck—Clear Lake clay, 0 to 1 percent slopes, MLRA 17.....	22
Mf—Marvin silty clay loam.....	23
Rg—Rincon silty clay loam.....	25
Sc—Sacramento clay, 0 to 2 percent slopes, MLRA 17.....	26
SmD—Sehorn-Balcom complex, 2 to 15 percent slopes.....	28
SmE2—Sehorn-Balcom complex, 15 to 30 percent slopes, eroded.....	30
So—Sycamore silt loam, 0 to 1 percent slopes, MLRA 17.....	32
Ss—Sycamore silty clay loam, 0 to 1 percent slopes, MLRA 17.....	33
Su—Sycamore complex.....	34
Sv—Sycamore complex, drained.....	36
TaA—Tehama loam, 0 to 2 percent slopes, loamy substratum, MLRA 17..	38
Tb—Tyndall very fine sandy loam.....	40
W—Water.....	41
Ya—Yolo silt loam, 0 to 2 percent slopes, MLRA 17.....	41
References	44

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

Custom Soil Resource Report

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.

Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and


Custom Soil Resource Report

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.






































Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.

Custom Soil Resource Report Soil Map



MAP LEGEND

 Area of Interest (AOI)	 Spoil Area
 Soil Map Unit Polygons	 Stony Spot
 Soil Map Unit Lines	 Very Stony Spot
 Soil Map Unit Points	 Wet Spot
 Special Point Features	 Other
 Blowout	 Special Line Features
 Borrow Pit	 Streams and Canals
 Clay Spot	 Rails
 Closed Depression	 Interstate Highways
 Gravel Pit	 US Routes
 Gravelly Spot	 Major Roads
 Landfill	 Local Roads
 Lava Flow	 Aerial Photography
 Marsh or swamp	
 Mine or Quarry	
 Miscellaneous Water	
 Perennial Water	
 Rock Outcrop	
 Saline Spot	
 Sandy Spot	
 Severely Eroded Spot	
 Sinkhole	
 Slide or Slip	
 Sodic Spot	

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at scales ranging from 1:20,000 to 1:24,000.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service
 Web Soil Survey URL:
 Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Sutter County, California
 Survey Area Data: Version 17, Jun 1, 2020

Soil Survey Area: Yolo County, California
 Survey Area Data: Version 16, Jun 1, 2020

Your area of interest (AOI) includes more than one soil survey area. These survey areas may have been mapped at different scales, with a different land use in mind, at different times, or at different levels of detail. This may result in map unit symbols, soil properties, and interpretations that do not completely agree across soil survey area boundaries.

Soil map units are labeled (as space allows) for map scales 1:50,000 or larger.

Date(s) aerial images were photographed: Data not available.

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
121	Columbia fine sandy loam, frequently flooded, 0 to 2 percent slopes	30.2	0.8%
122	Columbia loam, 0 to 2 percent slopes	0.3	0.0%
177	Water	20.9	0.5%
Subtotals for Soil Survey Area		51.4	1.3%
Totals for Area of Interest		3,936.4	100.0%

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
AaA	Arbuckle gravelly loam, 0 to 2 percent slopes, MLRA 17	115.1	2.9%
BrA	Brentwood silty clay loam, 0 to 2 percent slopes	34.8	0.9%
Ca	Capay silty clay, 0 percent slopes, MLRA 17	213.4	5.4%
Cb	Capay silty clay, 0 percent slopes, frequently flooded, MLRA 17	218.6	5.6%
Ck	Clear Lake clay, 0 to 1 percent slopes, MLRA 17	158.3	4.0%
Mf	Marvin silty clay loam	87.6	2.2%
Rg	Rincon silty clay loam	470.6	12.0%
Sc	Sacramento clay, 0 to 2 percent slopes, MLRA 17	1,952.3	49.6%
SmD	Sehorn-Balcom complex, 2 to 15 percent slopes	97.1	2.5%
SmE2	Sehorn-Balcom complex, 15 to 30 percent slopes, eroded	20.3	0.5%
So	Sycamore silt loam, 0 to 1 percent slopes, MLRA 17	7.6	0.2%
Ss	Sycamore silty clay loam, 0 to 1 percent slopes, MLRA 17	42.4	1.1%
Su	Sycamore complex	30.1	0.8%
Sv	Sycamore complex, drained	40.7	1.0%
TaA	Tehama loam, 0 to 2 percent slopes, loamy substratum, MLRA 17	186.2	4.7%
Tb	Tyndall very fine sandy loam	147.8	3.8%
W	Water	35.4	0.9%
Ya	Yolo silt loam, 0 to 2 percent slopes, MLRA 17	26.8	0.7%

Custom Soil Resource Report

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
Subtotals for Soil Survey Area		3,885.0	98.7%
Totals for Area of Interest		3,936.4	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Custom Soil Resource Report

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Sutter County, California

121—Columbia fine sandy loam, frequently flooded, 0 to 2 percent slopes

Map Unit Setting

National map unit symbol: hdz4
Elevation: 20 to 80 feet
Mean annual precipitation: 17 to 22 inches
Mean annual air temperature: 60 to 64 degrees F
Frost-free period: 260 to 280 days
Farmland classification: Not prime farmland

Map Unit Composition

Columbia, fine sandy loam, frequently flooded, and similar soils: 80 percent
Minor components: 20 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Columbia, Fine Sandy Loam, Frequently Flooded

Setting

Landform: Flood plains
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Mixed sandy alluvium

Typical profile

Ap or A - 0 to 14 inches: fine sandy loam
C1 to C4 - 14 to 60 inches: stratified fine sandy loam to very fine sandy loam

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat poorly drained
Runoff class: Very low
Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95 in/hr)
Depth to water table: About 36 to 60 inches
Frequency of flooding: FrequentNone
Frequency of ponding: None
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water capacity: Moderate (about 6.6 inches)

Interpretive groups

Land capability classification (irrigated): 4w
Land capability classification (nonirrigated): 4w
Hydrologic Soil Group: A
Hydric soil rating: Yes

Minor Components

Tisdale

Percent of map unit: 7 percent
Landform: Flood plains

Custom Soil Resource Report

Hydric soil rating: No

Holillipah

Percent of map unit: 7 percent

Landform: Flood plains

Hydric soil rating: Yes

Shanghai

Percent of map unit: 6 percent

Landform: Flood plains

Hydric soil rating: Yes

122—Columbia loam, 0 to 2 percent slopes

Map Unit Setting

National map unit symbol: hdz5

Elevation: 20 to 80 feet

Mean annual precipitation: 14 to 17 inches

Mean annual air temperature: 61 to 64 degrees F

Frost-free period: 260 to 280 days

Farmland classification: Prime farmland if irrigated

Map Unit Composition

Columbia, loam, and similar soils: 80 percent

Minor components: 20 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Columbia, Loam

Setting

Landform: Flood plains

Landform position (two-dimensional): Toeslope

Landform position (three-dimensional): Tread

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Mixed alluvium

Typical profile

H1 - 0 to 25 inches: loam

H2 - 25 to 60 inches: stratified sand to silt loam

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high
(0.57 to 1.98 in/hr)

Depth to water table: About 36 to 60 inches

Frequency of flooding: Rare

Frequency of ponding: None

Custom Soil Resource Report

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water capacity: Moderate (about 7.2 inches)

Interpretive groups

Land capability classification (irrigated): 2w

Land capability classification (nonirrigated): 4w

Hydrologic Soil Group: B

Hydric soil rating: Yes

Minor Components

Shanghai

Percent of map unit: 7 percent

Landform: Flood plains

Hydric soil rating: Yes

Holillipah

Percent of map unit: 7 percent

Landform: Flood plains

Hydric soil rating: Yes

Byington

Percent of map unit: 6 percent

Landform: Flood plains

Hydric soil rating: Yes

177—Water

Map Unit Composition

Water: 100 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Yolo County, California

AaA—Arbuckle gravelly loam, 0 to 2 percent slopes, MLRA 17

Map Unit Setting

National map unit symbol: 2t7r8
Elevation: 30 to 1,420 feet
Mean annual precipitation: 20 to 32 inches
Mean annual air temperature: 61 to 63 degrees F
Frost-free period: 200 to 280 days
Farmland classification: Prime farmland if irrigated

Map Unit Composition

Arbuckle and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Arbuckle

Setting

Landform: Stream terraces
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Alluvium derived from metamorphic and sedimentary rock

Typical profile

A1 - 0 to 2 inches: gravelly loam
A2 - 2 to 14 inches: gravelly loam
Bt1 - 14 to 25 inches: gravelly loam
Bt2 - 25 to 59 inches: gravelly sandy clay loam
Bt3 - 59 to 72 inches: very gravelly loam

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.28 to 1.28 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline (0.3 to 0.5 mmhos/cm)
Available water capacity: Moderate (about 8.0 inches)

Interpretive groups

Land capability classification (irrigated): 2s
Land capability classification (nonirrigated): 3s
Hydrologic Soil Group: B
Hydric soil rating: No

Minor Components

Maywood

Percent of map unit: 5 percent
Hydric soil rating: No

Cortina

Percent of map unit: 5 percent
Hydric soil rating: No

Hillgate

Percent of map unit: 5 percent
Hydric soil rating: No

BrA—Brentwood silty clay loam, 0 to 2 percent slopes

Map Unit Setting

National map unit symbol: hdvb
Elevation: 50 to 400 feet
Mean annual precipitation: 12 to 20 inches
Mean annual air temperature: 61 to 63 degrees F
Frost-free period: 280 days
Farmland classification: Prime farmland if irrigated

Map Unit Composition

Brentwood and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Brentwood

Setting

Landform: Alluvial fans
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Base slope
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Alluvium derived from sedimentary rock

Typical profile

H1 - 0 to 10 inches: silty clay loam
H2 - 10 to 35 inches: silty clay loam
H3 - 35 to 60 inches: silty clay loam

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)

Custom Soil Resource Report

Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water capacity: High (about 10.8 inches)

Interpretive groups

Land capability classification (irrigated): 1
Land capability classification (nonirrigated): 4c
Hydrologic Soil Group: C
Hydric soil rating: No

Minor Components

Yolo

Percent of map unit: 5 percent
Hydric soil rating: No

Zamora

Percent of map unit: 5 percent
Hydric soil rating: No

Rincon

Percent of map unit: 3 percent
Hydric soil rating: No

Myers

Percent of map unit: 2 percent
Hydric soil rating: No

Ca—Capay silty clay, 0 percent slopes, MLRA 17

Map Unit Setting

National map unit symbol: 2xc8z
Elevation: 20 to 180 feet
Mean annual precipitation: 20 to 24 inches
Mean annual air temperature: 61 to 62 degrees F
Frost-free period: 317 to 326 days
Farmland classification: Prime farmland if irrigated

Map Unit Composition

Capay and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Capay

Setting

Landform: Basin floors
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Tread
Down-slope shape: Linear

Custom Soil Resource Report

Across-slope shape: Linear

Parent material: Silty and clayey alluvium derived from igneous, metamorphic and sedimentary rock

Typical profile

Ap - 0 to 11 inches: silty clay

A - 11 to 18 inches: silty clay

Bss1 - 18 to 36 inches: silty clay

Bkss - 36 to 49 inches: silty clay

B'ss2 - 49 to 64 inches: silty clay

Properties and qualities

Slope: 0 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Moderately well drained

Runoff class: High

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: RareNone

Frequency of ponding: Frequent

Calcium carbonate, maximum content: 1 percent

Maximum salinity: Nonsaline (0.2 to 1.0 mmhos/cm)

Sodium adsorption ratio, maximum: 10.0

Available water capacity: Moderate (about 7.8 inches)

Interpretive groups

Land capability classification (irrigated): 2s

Land capability classification (nonirrigated): 4s

Hydrologic Soil Group: C

Hydric soil rating: No

Minor Components

Willows

Percent of map unit: 4 percent

Landform: Basin floors

Hydric soil rating: Yes

Clear lake

Percent of map unit: 4 percent

Landform: Basin floors

Hydric soil rating: Yes

Marvin

Percent of map unit: 4 percent

Hydric soil rating: No

Myers

Percent of map unit: 3 percent

Hydric soil rating: No

Cb—Capay silty clay, 0 percent slopes, frequently flooded, MLRA 17

Map Unit Setting

National map unit symbol: 2y0f5
Elevation: 30 feet
Mean annual precipitation: 20 to 20 inches
Mean annual air temperature: 62 to 62 degrees F
Frost-free period: 322 to 330 days
Farmland classification: Not prime farmland

Map Unit Composition

Capay and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Capay

Setting

Landform: Basin floors
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Silty and clayey alluvium derived from igneous, metamorphic and sedimentary rock

Typical profile

Ap - 0 to 11 inches: silty clay
A - 11 to 18 inches: silty clay
Bss1 - 18 to 36 inches: silty clay
Bkss - 36 to 49 inches: silty clay
B'ss2 - 49 to 64 inches: silty clay

Properties and qualities

Slope: 0 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Moderately well drained
Runoff class: High
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)
Depth to water table: About 48 to 72 inches
Frequency of flooding: FrequentNone
Frequency of ponding: Frequent
Calcium carbonate, maximum content: 1 percent
Maximum salinity: Nonsaline (0.2 to 1.0 mmhos/cm)
Sodium adsorption ratio, maximum: 10.0
Available water capacity: Moderate (about 7.8 inches)

Interpretive groups

Land capability classification (irrigated): 4w

Custom Soil Resource Report

Land capability classification (nonirrigated): 4w
Hydrologic Soil Group: C
Hydric soil rating: Yes

Minor Components

Marvin

Percent of map unit: 7 percent
Hydric soil rating: No

Clear lake

Percent of map unit: 4 percent
Landform: Basin floors
Hydric soil rating: Yes

Willows

Percent of map unit: 4 percent
Landform: Basin floors
Hydric soil rating: Yes

Ck—Clear Lake clay, 0 to 1 percent slopes, MLRA 17

Map Unit Setting

National map unit symbol: 2vbsz
Elevation: 20 to 400 feet
Mean annual precipitation: 17 to 19 inches
Mean annual air temperature: 61 to 63 degrees F
Frost-free period: 260 to 280 days
Farmland classification: Prime farmland if irrigated

Map Unit Composition

Clear lake and similar soils: 90 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Clear Lake

Setting

Landform: Basin floors
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Basin alluvium derived from igneous, metamorphic and sedimentary rock

Typical profile

Ap - 0 to 10 inches: clay
Ag - 10 to 25 inches: clay
Bssg1 - 25 to 42 inches: clay
Bssg2 - 42 to 68 inches: clay

Custom Soil Resource Report

Properties and qualities

Slope: 0 to 1 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Poorly drained
Runoff class: Very low
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)
Depth to water table: About 0 to 35 inches
Frequency of flooding: Rare
Frequency of ponding: Frequent
Calcium carbonate, maximum content: 4 percent
Maximum salinity: Nonsaline to very slightly saline (1.0 to 2.0 mmhos/cm)
Sodium adsorption ratio, maximum: 14.0
Available water capacity: Moderate (about 8.5 inches)

Interpretive groups

Land capability classification (irrigated): 2w
Land capability classification (nonirrigated): 4w
Hydrologic Soil Group: C/D
Hydric soil rating: Yes

Minor Components

Capay

Percent of map unit: 4 percent
Landform: Basin floors
Hydric soil rating: Yes

Subaco

Percent of map unit: 3 percent
Landform: Flood plains
Hydric soil rating: Yes

Oswald

Percent of map unit: 3 percent
Landform: Basin floors
Hydric soil rating: Yes

Mf—Marvin silty clay loam

Map Unit Setting

National map unit symbol: hdwb
Elevation: 20 to 100 feet
Mean annual precipitation: 20 inches
Mean annual air temperature: 63 degrees F
Frost-free period: 280 days
Farmland classification: Prime farmland if irrigated

Map Unit Composition

Marvin and similar soils: 85 percent

Custom Soil Resource Report

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Marvin

Setting

Landform: Rims on basin floors

Landform position (two-dimensional): Toeslope

Landform position (three-dimensional): Rise, tal

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Mixed silty and clayey alluvium

Typical profile

H1 - 0 to 12 inches: silty clay loam

H2 - 12 to 41 inches: silty clay

H3 - 41 to 60 inches: silty clay loam

Properties and qualities

Slope: 0 to 1 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Runoff class: High

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: RareNone

Frequency of ponding: None

Calcium carbonate, maximum content: 5 percent

Maximum salinity: Nonsaline to slightly saline (0.0 to 4.0 mmhos/cm)

Available water capacity: High (about 10.5 inches)

Interpretive groups

Land capability classification (irrigated): 2s

Land capability classification (nonirrigated): 4s

Hydrologic Soil Group: C

Hydric soil rating: No

Minor Components

Capay

Percent of map unit: 5 percent

Landform: Rims

Hydric soil rating: Yes

Rincon

Percent of map unit: 5 percent

Hydric soil rating: No

Pescadero

Percent of map unit: 3 percent

Hydric soil rating: No

Unnamed

Percent of map unit: 2 percent

Hydric soil rating: No

Rg—Rincon silty clay loam

Map Unit Setting

National map unit symbol: hdww
Elevation: 50 to 350 feet
Mean annual precipitation: 20 inches
Mean annual air temperature: 61 degrees F
Frost-free period: 275 days
Farmland classification: Prime farmland if irrigated

Map Unit Composition

Rincon and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Rincon

Setting

Landform: Alluvial fans
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Base slope
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Alluvium derived from sedimentary rock

Typical profile

H1 - 0 to 15 inches: silty clay loam
H2 - 15 to 56 inches: silty clay loam
H3 - 56 to 72 inches: silty clay loam

Properties and qualities

Slope: 0 to 1 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: High
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water capacity: High (about 9.4 inches)

Interpretive groups

Land capability classification (irrigated): 2s
Land capability classification (nonirrigated): 3s
Hydrologic Soil Group: C
Hydric soil rating: No

Minor Components

Brentwood

Percent of map unit: 5 percent
Hydric soil rating: No

Marvin

Percent of map unit: 3 percent
Hydric soil rating: No

Tehama

Percent of map unit: 3 percent
Hydric soil rating: No

Zamora

Percent of map unit: 2 percent
Hydric soil rating: No

Yolo

Percent of map unit: 2 percent
Hydric soil rating: No

Sc—Sacramento clay, 0 to 2 percent slopes, MLRA 17

Map Unit Setting

National map unit symbol: 2w8b4
Elevation: 0 to 80 feet
Mean annual precipitation: 15 to 23 inches
Mean annual air temperature: 61 to 62 degrees F
Frost-free period: 250 to 300 days
Farmland classification: Prime farmland if irrigated and drained

Map Unit Composition

Sacramento and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Sacramento

Setting

Landform: Basin floors
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Clayey alluvium derived from igneous, metamorphic and sedimentary rock

Typical profile

Apg - 0 to 7 inches: clay
Ag1 - 7 to 16 inches: silty clay
Ag2 - 16 to 31 inches: clay

Custom Soil Resource Report

Ag3 - 31 to 38 inches: clay
Ag4 - 38 to 53 inches: clay
Cg1 - 53 to 60 inches: clay
Cg2 - 60 to 69 inches: silty clay
Cg3 - 69 to 77 inches: silty clay loam

Properties and qualities

Slope: 0 to 2 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Poorly drained
Runoff class: High
Capacity of the most limiting layer to transmit water (K_{sat}): Moderately low (0.04 to 0.06 in/hr)
Depth to water table: About 0 to 35 inches
Frequency of flooding: Rare
Frequency of ponding: Frequent
Calcium carbonate, maximum content: 3 percent
Gypsum, maximum content: 1 percent
Maximum salinity: Nonsaline to slightly saline (0.2 to 4.0 mmhos/cm)
Sodium adsorption ratio, maximum: 4.0
Available water capacity: Moderate (about 8.5 inches)

Interpretive groups

Land capability classification (irrigated): 3w
Land capability classification (nonirrigated): 4w
Hydrologic Soil Group: C/D
Hydric soil rating: Yes

Minor Components

Sycamore

Percent of map unit: 3 percent
Landform: Alluvial fans
Hydric soil rating: Yes

Willows

Percent of map unit: 3 percent
Landform: Basin floors
Hydric soil rating: Yes

Omni

Percent of map unit: 3 percent
Landform: Basin floors
Hydric soil rating: Yes

Merritt

Percent of map unit: 3 percent
Landform: Alluvial fans
Hydric soil rating: Yes

Clear lake

Percent of map unit: 3 percent
Landform: Basin floors
Hydric soil rating: Yes

SmD—Sehorn-Balcom complex, 2 to 15 percent slopes

Map Unit Setting

National map unit symbol: hdx
Elevation: 100 to 2,000 feet
Mean annual precipitation: 15 to 35 inches
Mean annual air temperature: 57 to 64 degrees F
Frost-free period: 200 to 340 days
Farmland classification: Farmland of statewide importance

Map Unit Composition

Sehorn and similar soils: 60 percent
Balcom and similar soils: 30 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Sehorn

Setting

Landform: Hills
Landform position (two-dimensional): Footslope
Landform position (three-dimensional): Side slope
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Calcareous residuum weathered from sedimentary rock

Typical profile

H1 - 0 to 10 inches: clay
H2 - 10 to 40 inches: clay
H3 - 40 to 60 inches: weathered bedrock

Properties and qualities

Slope: 2 to 15 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Very high
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water capacity: Moderate (about 6.0 inches)

Interpretive groups

Land capability classification (irrigated): 3e
Land capability classification (nonirrigated): 4e
Hydrologic Soil Group: C
Ecological site: R015XE001CA - Clayey Hills 10-14" p.z.
Hydric soil rating: No

Custom Soil Resource Report

Description of Balcom

Setting

Landform: Hills
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Concave
Across-slope shape: Linear
Parent material: Residuum weathered from calcareous sandstone

Typical profile

H1 - 0 to 20 inches: silty clay loam
H2 - 20 to 37 inches: silty clay loam
H3 - 37 to 60 inches: weathered bedrock

Properties and qualities

Slope: 9 to 15 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: High
Capacity of the most limiting layer to transmit water (Ksat): Very low to low (0.00 to 0.01 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum content: 10 percent
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water capacity: Moderate (about 7.0 inches)

Interpretive groups

Land capability classification (irrigated): 3e
Land capability classification (nonirrigated): 4e
Hydrologic Soil Group: C
Ecological site: R015XE001CA - Clayey Hills 10-14" p.z.
Hydric soil rating: No

Minor Components

Positas

Percent of map unit: 3 percent
Hydric soil rating: No

Unnamed, in swales

Percent of map unit: 3 percent
Hydric soil rating: No

Myers

Percent of map unit: 2 percent
Hydric soil rating: No

Corning

Percent of map unit: 2 percent
Hydric soil rating: No

SmE2—Sehorn-Balcom complex, 15 to 30 percent slopes, eroded

Map Unit Setting

National map unit symbol: hdxg
Elevation: 70 to 2,000 feet
Mean annual precipitation: 15 to 35 inches
Mean annual air temperature: 57 to 64 degrees F
Frost-free period: 200 to 340 days
Farmland classification: Not prime farmland

Map Unit Composition

Sehorn and similar soils: 50 percent
Balcom and similar soils: 40 percent
Minor components: 10 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Sehorn

Setting

Landform: Hills
Landform position (two-dimensional): Backslope
Landform position (three-dimensional): Side slope
Down-slope shape: Concave
Across-slope shape: Convex
Parent material: Calcareous residuum weathered from sedimentary rock

Typical profile

H1 - 0 to 8 inches: clay
H2 - 8 to 38 inches: clay
H3 - 38 to 60 inches: weathered bedrock

Properties and qualities

Slope: 15 to 30 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Very high
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water capacity: Low (about 5.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified
Land capability classification (nonirrigated): 4e
Hydrologic Soil Group: D
Ecological site: R015XE001CA - Clayey Hills 10-14" p.z.

Custom Soil Resource Report

Hydric soil rating: No

Description of Balcom

Setting

Landform: Hills

Landform position (two-dimensional): Backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Concave

Across-slope shape: Concave

Parent material: Residuum weathered from calcareous sandstone

Typical profile

H1 - 0 to 20 inches: silty clay loam

H2 - 20 to 37 inches: silty clay loam

H3 - 37 to 60 inches: weathered bedrock

Properties and qualities

Slope: 15 to 30 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Very low to low (0.00 to 0.01 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Calcium carbonate, maximum content: 10 percent

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water capacity: Moderate (about 7.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: C

Ecological site: R015XE001CA - Clayey Hills 10-14" p.z.

Hydric soil rating: No

Minor Components

Corning

Percent of map unit: 5 percent

Hydric soil rating: No

Positas

Percent of map unit: 5 percent

Hydric soil rating: No

So—Sycamore silt loam, 0 to 1 percent slopes, MLRA 17

Map Unit Setting

National map unit symbol: 2xcbj

Elevation: 10 to 130 feet

Mean annual precipitation: 20 to 21 inches

Mean annual air temperature: 61 to 62 degrees F

Frost-free period: 322 to 330 days

Farmland classification: Prime farmland if irrigated and drained

Map Unit Composition

Sycamore and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Sycamore

Setting

Landform: Natural levees, flood-plain splays

Landform position (three-dimensional): Tread

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Alluvium derived from igneous, metamorphic and sedimentary rock

Typical profile

Ap - 0 to 4 inches: silt loam

A - 4 to 14 inches: silt loam

Bw - 14 to 26 inches: silt loam

Bwk - 26 to 44 inches: silt loam

C - 44 to 60 inches: silt loam

Properties and qualities

Slope: 0 to 1 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.60 to 2.00 in/hr)

Depth to water table: About 36 to 60 inches

Frequency of flooding: NoneOccasional

Frequency of ponding: None

Calcium carbonate, maximum content: 1 percent

Maximum salinity: Nonsaline (0.2 to 0.7 mmhos/cm)

Sodium adsorption ratio, maximum: 1.0

Available water capacity: Very high (about 12.4 inches)

Interpretive groups

Land capability classification (irrigated): 2w

Custom Soil Resource Report

Land capability classification (nonirrigated): 4w
Hydrologic Soil Group: C
Hydric soil rating: No

Minor Components

Maria

Percent of map unit: 4 percent
Landform: Alluvial fans
Hydric soil rating: Yes

Tyndall

Percent of map unit: 4 percent

Merritt

Percent of map unit: 4 percent

Yolo

Percent of map unit: 3 percent

Ss—Sycamore silty clay loam, 0 to 1 percent slopes, MLRA 17

Map Unit Setting

National map unit symbol: 2xcbm
Elevation: 10 to 30 feet
Mean annual precipitation: 20 to 20 inches
Mean annual air temperature: 61 to 62 degrees F
Frost-free period: 322 to 329 days
Farmland classification: Prime farmland if irrigated and drained

Map Unit Composition

Sycamore and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Sycamore

Setting

Landform: Natural levees
Landform position (three-dimensional): Tread
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Alluvium derived from igneous, metamorphic and sedimentary rock

Typical profile

Ap - 0 to 4 inches: silty clay loam
A - 4 to 14 inches: silty clay loam
Bw - 14 to 26 inches: silty clay loam
Bwk - 26 to 44 inches: silty clay loam
C - 44 to 60 inches: loam

Custom Soil Resource Report

Properties and qualities

Slope: 0 to 1 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat poorly drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.60 in/hr)
Depth to water table: About 36 to 60 inches
Frequency of flooding: RareNone
Frequency of ponding: None
Calcium carbonate, maximum content: 1 percent
Maximum salinity: Nonsaline (0.2 to 0.7 mmhos/cm)
Sodium adsorption ratio, maximum: 1.0
Available water capacity: High (about 10.9 inches)

Interpretive groups

Land capability classification (irrigated): 2w
Land capability classification (nonirrigated): 4w
Hydrologic Soil Group: C
Hydric soil rating: No

Minor Components

Maria

Percent of map unit: 4 percent
Landform: Alluvial fans
Hydric soil rating: Yes

Tyndall

Percent of map unit: 4 percent
Hydric soil rating: No

Merritt

Percent of map unit: 4 percent
Hydric soil rating: No

Marvin

Percent of map unit: 3 percent
Hydric soil rating: No

Su—Sycamore complex

Map Unit Setting

National map unit symbol: hdxq
Elevation: 0 to 60 feet
Mean annual precipitation: 15 to 20 inches
Mean annual air temperature: 61 degrees F
Frost-free period: 280 days
Farmland classification: Prime farmland if irrigated and drained

Map Unit Composition

Sycamore and similar soils: 60 percent

Sycamore and similar soils: 30 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Sycamore

Setting

Landform: Alluvial fans

Landform position (two-dimensional): Toeslope

Landform position (three-dimensional): Base slope

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Mixed alluvium derived from sedimentary rock

Typical profile

H1 - 0 to 14 inches: silty clay loam

H2 - 14 to 44 inches: silty clay loam

H3 - 44 to 60 inches: silty clay

Properties and qualities

Slope: 0 to 1 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)

Depth to water table: About 36 to 60 inches

Frequency of flooding: RareNone

Frequency of ponding: None

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water capacity: High (about 10.4 inches)

Interpretive groups

Land capability classification (irrigated): 2w

Land capability classification (nonirrigated): 4w

Hydrologic Soil Group: C

Hydric soil rating: Yes

Description of Sycamore

Setting

Landform: Alluvial fans

Landform position (two-dimensional): Toeslope

Landform position (three-dimensional): Base slope

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Mixed alluvium derived from sedimentary rock

Typical profile

H1 - 0 to 14 inches: silt loam

H2 - 14 to 44 inches: silt loam

H3 - 44 to 60 inches: silty clay

Properties and qualities

Slope: 0 to 1 percent

Custom Soil Resource Report

Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat poorly drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)
Depth to water table: About 36 to 60 inches
Frequency of flooding: RareNone
Frequency of ponding: None
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water capacity: High (about 10.7 inches)

Interpretive groups

Land capability classification (irrigated): 2w
Land capability classification (nonirrigated): 4w
Hydrologic Soil Group: C
Hydric soil rating: Yes

Minor Components

Merritt

Percent of map unit: 3 percent
Landform: Alluvial fans
Hydric soil rating: Yes

Marvin

Percent of map unit: 3 percent
Hydric soil rating: No

Sacramento

Percent of map unit: 3 percent
Landform: Basin floors
Hydric soil rating: Yes

Unnamed

Percent of map unit: 1 percent
Hydric soil rating: No

Sv—Sycamore complex, drained

Map Unit Setting

National map unit symbol: hdxr
Elevation: 0 to 60 feet
Mean annual precipitation: 15 to 20 inches
Mean annual air temperature: 61 degrees F
Frost-free period: 280 days
Farmland classification: Prime farmland if irrigated

Map Unit Composition

Sycamore and similar soils: 60 percent
Sycamore and similar soils: 25 percent

Custom Soil Resource Report

Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Sycamore

Setting

Landform: Alluvial fans
Landform position (two-dimensional): Toeslope
Landform position (three-dimensional): Base slope
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Mixed alluvium derived from sedimentary rock

Typical profile

H1 - 0 to 14 inches: silty clay loam
H2 - 14 to 44 inches: silty clay loam
H3 - 44 to 60 inches: silty clay

Properties and qualities

Slope: 0 to 1 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat poorly drained
Runoff class: Medium
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water capacity: High (about 10.7 inches)

Interpretive groups

Land capability classification (irrigated): 1
Land capability classification (nonirrigated): 4c
Hydrologic Soil Group: C
Hydric soil rating: Yes

Description of Sycamore

Setting

Landform: Alluvial fans
Landform position (three-dimensional): Base slope
Down-slope shape: Linear
Across-slope shape: Linear
Parent material: Mixed alluvium derived from sedimentary rock

Typical profile

H1 - 0 to 14 inches: silt loam
H2 - 14 to 44 inches: silt loam
H3 - 44 to 60 inches: silty clay

Properties and qualities

Slope: 0 to 1 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat poorly drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20 to 0.57 in/hr)

Custom Soil Resource Report

Depth to water table: About 36 to 60 inches
Frequency of flooding: None
Frequency of ponding: None
Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)
Available water capacity: High (about 10.7 inches)

Interpretive groups

Land capability classification (irrigated): 1
Land capability classification (nonirrigated): 4c
Hydrologic Soil Group: C
Hydric soil rating: Yes

Minor Components

Marvin

Percent of map unit: 5 percent
Hydric soil rating: No

Sacramento

Percent of map unit: 4 percent
Landform: Alluvial fans
Hydric soil rating: Yes

Merritt

Percent of map unit: 3 percent
Landform: Alluvial fans
Hydric soil rating: Yes

Unnamed

Percent of map unit: 3 percent
Hydric soil rating: No

TaA—Tehama loam, 0 to 2 percent slopes, loamy substratum, MLRA 17

Map Unit Setting

National map unit symbol: 2srj5
Elevation: 50 to 580 feet
Mean annual precipitation: 19 to 27 inches
Mean annual air temperature: 63 degrees F
Frost-free period: 265 days
Farmland classification: Prime farmland if irrigated

Map Unit Composition

Tehama and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Tehama

Setting

Landform: Alluvial fans

Custom Soil Resource Report

Landform position (two-dimensional): Toeslope

Landform position (three-dimensional): Tread

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Mixed fine-loamy alluvium derived from sedimentary rock

Typical profile

Ap - 0 to 10 inches: loam

Bt - 10 to 40 inches: clay loam

BCt - 40 to 63 inches: gravelly loam

C - 63 to 75 inches: sandy loam

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: High

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to moderately high (0.06 to 0.20 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline to very slightly saline (0.0 to 2.0 mmhos/cm)

Available water capacity: High (about 10.3 inches)

Interpretive groups

Land capability classification (irrigated): 2s

Land capability classification (nonirrigated): 4s

Hydrologic Soil Group: C

Hydric soil rating: No

Minor Components

Yolo

Percent of map unit: 4 percent

Hydric soil rating: No

Zamora

Percent of map unit: 4 percent

Hydric soil rating: No

Brentwood

Percent of map unit: 4 percent

Hydric soil rating: No

Rincon

Percent of map unit: 3 percent

Hydric soil rating: No

Tb—Tyndall very fine sandy loam

Map Unit Setting

National map unit symbol: hdxw

Elevation: 0 to 70 feet

Mean annual precipitation: 17 inches

Mean annual air temperature: 63 degrees F

Frost-free period: 280 days

Farmland classification: Prime farmland if irrigated and drained

Map Unit Composition

Tyndall and similar soils: 85 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Tyndall

Setting

Landform: Alluvial fans

Landform position (two-dimensional): Toeslope

Landform position (three-dimensional): Base slope

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Alluvium derived from sedimentary rock

Typical profile

H1 - 0 to 16 inches: very fine sandy loam

H2 - 16 to 60 inches: very fine sandy loam

Properties and qualities

Slope: 0 to 1 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Somewhat poorly drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): High (1.98 to 5.95 in/hr)

Depth to water table: About 36 to 72 inches

Frequency of flooding: RareNone

Frequency of ponding: None

Calcium carbonate, maximum content: 10 percent

Maximum salinity: Slightly saline to moderately saline (4.0 to 8.0 mmhos/cm)

Available water capacity: High (about 9.6 inches)

Interpretive groups

Land capability classification (irrigated): 2w

Land capability classification (nonirrigated): 4w

Hydrologic Soil Group: B

Hydric soil rating: No

Minor Components

Reiff

Percent of map unit: 3 percent
Hydric soil rating: No

Lang

Percent of map unit: 3 percent
Ecological site: R016XA002CA - Freshwater, Stratified, Fluventic Sites (PROVISIONAL)
Hydric soil rating: No

Laugenour

Percent of map unit: 3 percent
Landform: Alluvial fans
Ecological site: R016XA002CA - Freshwater, Stratified, Fluventic Sites (PROVISIONAL)
Hydric soil rating: Yes

Sycamore

Percent of map unit: 3 percent
Landform: Alluvial fans
Hydric soil rating: Yes

Unnamed

Percent of map unit: 3 percent
Hydric soil rating: No

W—Water

Map Unit Composition

Water: 100 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Ya—Yolo silt loam, 0 to 2 percent slopes, MLRA 17

Map Unit Setting

National map unit symbol: 2w89y
Elevation: 20 to 2,020 feet
Mean annual precipitation: 19 to 35 inches
Mean annual air temperature: 59 to 62 degrees F
Frost-free period: 250 to 270 days
Farmland classification: Prime farmland if irrigated

Map Unit Composition

Yolo and similar soils: 85 percent
Minor components: 15 percent
Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Yolo

Setting

Landform: Alluvial fans, flood plains

Landform position (two-dimensional): Toeslope

Landform position (three-dimensional): Tread

Down-slope shape: Linear

Across-slope shape: Linear

Parent material: Alluvium derived from igneous, metamorphic and sedimentary rock

Typical profile

Ap1 - 0 to 2 inches: silt loam

Ap2 - 2 to 8 inches: silt loam

A1 - 8 to 19 inches: silt loam

A2 - 19 to 26 inches: silt loam

C1 - 26 to 33 inches: silt loam

C2 - 33 to 41 inches: silt loam

Ab - 41 to 58 inches: silty clay loam

C'3 - 58 to 65 inches: silt loam

Properties and qualities

Slope: 0 to 2 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Well drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.20 to 2.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: Rare

Frequency of ponding: None

Calcium carbonate, maximum content: 1 percent

Maximum salinity: Nonsaline (0.2 to 1.0 mmhos/cm)

Available water capacity: Very high (about 12.2 inches)

Interpretive groups

Land capability classification (irrigated): 1

Land capability classification (nonirrigated): 4c

Hydrologic Soil Group: B

Hydric soil rating: No

Minor Components

Arbuckle

Percent of map unit: 3 percent

Hydric soil rating: No

Zamora

Percent of map unit: 2 percent

Hydric soil rating: No

Soboba

Percent of map unit: 2 percent

Hydric soil rating: No

Reiff

Percent of map unit: 2 percent

Custom Soil Resource Report

Hydric soil rating: No

Loamy alluvial land

Percent of map unit: 2 percent

Hydric soil rating: No

Brentwood

Percent of map unit: 2 percent

Hydric soil rating: No

Sycamore

Percent of map unit: 2 percent

Landform: Alluvial fans

Hydric soil rating: Yes

References

- American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.
- American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.
- Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.
- Federal Register. July 13, 1994. Changes in hydric soils of the United States.
- Federal Register. September 18, 2002. Hydric soils of the United States.
- Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.
- National Research Council. 1995. Wetlands: Characteristics and boundaries.
- Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_054262
- Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053577
- Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053580
- Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.
- United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.
- United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2_053374
- United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. <http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/landuse/rangepasture/?cid=stelprdb1043084>

Custom Soil Resource Report

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf